

Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: NKDA

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
N/A	Isotonic/ Hypotonic/ Hypertonic	N/A	N/A	N/A

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is Correct dose?	IVP - List solution to dilute and rate to push.  IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzinepamoate(vistaril)	Antihistamines	Anxiety	50mg PO Q6H PRN	Y  N	N/A	drowsiness  headache  dry mouth  Skin rash.	Drinking alcohol can increase certain side effects of Vistaril.  2.notify your provider about all your medications because it can cause serious heart problem  3. This medicine may impair your thinking or reactions. Be careful if you drive or do anything that requires you to be alert  4. You should not take Vistaril for longer than 4 months
Lorazepam	Benzodiazepine Anticonvulsants	Anxiety	0.5mg PO BID PRN	Y  N	N/A	Sedation dizziness fatigue impaired cognitive function	1.never discontinue abruptly can cause withdrawal  2.Take as directed because it is highly addictive  3.avoid alcohol potentiates effects of alcohol  4. Caution while driving due to slower reflexes and response time.
Haloperidol(Haldol)	Misallaneous antipsychotic agents	Psychotic disorder	5mg PO Q6h PRN	Y  N	N/A	Tachycardia  Orthostatic hypotension weight gain seizures	1.change positions slowly because of orthostatic hypotension  2.assees the vital signs B/P and HR  3.limit sun exposure wear sunscreen or sunglasses  4.may take 2 -4 weeks or if not months for full effect
Risperidone(Risperdal)	Atypical	Severe	0.5mg	Y	N/A	Anxiety	1.daily weights and exercise

Medication Worksheet - Current Medications & PRN for Last 24 Hours

Allergies: NKDA

sperdal)	Antipsychotics	Psychosis	PO Daily	N		Headache weight gain Sedation dyslipidemia	2. Adhere to the medication regime for effective work. 3.observe for signs of infection(high fever) and vital signs 4.use sugar free liquids and candies for dry mouth
Donepezil(HCL)	Cholinesterase inhibitors.	Mild to moderate Alzheimer's	10mg PO daily	Y N	N/A	Nausea diarrhea Not sleeping well Feeling tired not wanting to eat	1. Monitor electrolytes and hydrate well. 2. Be careful if you drive or do anything that requires you to be awake and alert. Donepezil can cause side effects related to impaired thinking 3. Do not swallow the tablet whole. Allow it to dissolve on your tongue. 4. Do not stop taking the medicine abruptly.