

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
D5NS+20KCl (115ml/hr)	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input checked="" type="checkbox"/>	Patient was NPO and on a clear liquid diet --> Hydration, electrolyte balance	Sodium, Potassium, Chloride, Blood Glucose	Fluid overload, Phlebitis, Warmth, redness, swelling at IV site, Electrolyte imbalance

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT		Date: 1/29/2025	Allergies: Adhesive tape, Cefedindir, Lactose Intolerant	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Dicyclomine	Gastrointestinal anticholinergic	Antispasmodic, Anticholinergic Relieves smooth muscle spasm of the GI tract	20mg capsule PO QID	Yes <a href="#">Click here to enter text.</a>	N/A	Dry mouth, thirst, dizziness, tachycardia, sedation, constipation, decreased sweating	1. Perform oral hygiene as this medication can cause dry mouth. Sucking on sugar-free hard candy may help alleviate dry mouth 2. Avoid hot environments and ensure hydration as this medication can decrease sweating 3. Empty bladder before taking if urinary retention occurs 4. May cause dizziness/ blurred vision Be careful getting OOB/ambulating, use call light for assistance.
Erythromycin base	Antiinfective	Tx of infections caused by suspected bacteria	250mg tab PO TID	No Choose an item. 30-50mg/kg/d for a total of	N/A	Seizures, Dysrhythmias, QT prolongation, Hepatotoxicity, esophagitis,	1. Report sore throat, fever, fatigue as this could indicate super infection, rhythm changes in the heart, hearing loss 2. Take medication without to improve absorption

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT	Date: 1/29/2025	Allergies: Adhesive tape, Cefedinir, Lactose Intolerant		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				2,145mg-3,575 --> Falls below range		anaphylaxis	3. Notify nurse of diarrhea, dark urine, pale stools, jaundice, mor severe abdominal pain 4. Assess for signs of infection, hepatic levels, I&O
Hydroxyzine hydrochloride	Antiemetic, antihistamine, anxiolytic	Relief of anxiety, allergies, antiemetic	25mg tab PO nightly	No Choose an item. 0.6mg/kg PO for a total of 42.9mg PO --> Falls below range	N/A	Constipation, Dizziness, Dry mouth, Drowsiness, prolong QT, Urine retention	1. Assess anticholinergic effects: dry mouth, dizziness, confusion, hypotension --> Monitor B/P 2. Rise slowly because fainting may occur, use call light for assistance 3. Enforce fall precautions as this medication can increase drowsiness and dizziness 4. Report difficulty urinating as this medication may cause urinary retention
Hyoscyamine	Anticholinergic	Decreases muscle spasms	0.125mg sublingual QID before meals nightly	Yes Choose an item. Click here to enter text.	N/A	Dry mouth, Blurred vision, constipation, dizziness, urinary retention	1. Perform oral hygiene as this medication can cause dry mouth. Sucking on sugar-free hard candy may help alleviate dry mouth 2. May cause dizziness/ blurred vision Be careful getting OOB/ambulating, use call light for assistance. 3. Increase fluids and fiber rich foods

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT	Date: 1/29/2025	Allergies: Adhesive tape, Cefedinir, Lactose Intolerant		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							(whole grain, apples) to prevent constipation  4. Avoid eating or drinking for at least 5-10 minutes after taking medication to ensure proper absorption
Lansoprazole DR	Antiulcer, PPI	Suppress secretion of gastric acid	30mg Capsule PO daily before meals	Yes Choose an item. Click here to enter text.	N/A	Diarrhea, headache	1. Report severe diarrhea, cramping, blood in stools, fever 2. May take 1-4 days to see full benefit 3. Increase intake of fluids as this medication may cause constipation and diarrhea 4. Decrease/avoid spicy, fatty, or acidic foods as this may worsen acid reflux
Lisinopril	ACE Inhibitor	Lowers blood pressure, treats HTN	5mg tab PO daily	Yes Choose an item. Click here to enter text.	N/A	Angioedema, hyperkalemia, hypoglycemia, hypotension	1. Monitor blood pressure 2. May cause orthostatic hypotension, change position slowly, use call light for assistance out of bed 3. Call HCP of mouth sores, sore throat, fever, swelling of hands feet, irregular pulse, SOB, chest pain 4. Caution patient to not withdraw medication abruptly to prevent rebound hypertension

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT	Date: 1/29/2025	Allergies: Adhesive tape, Cefedinir, Lactose Intolerant		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Propranolol	Beta blocker	Treat HTN	10mg tab BID	No Choose an item. Below range, 0.5-1mg/kg/d BID (35.8-71.5mg/day )	N/A	Bradycardia, laryngospasm, thrombocytopenia, hypotension	<ol style="list-style-type: none"> <li>1. Assess/monitor blood pressure &amp; apical pulse</li> <li>2. Do not abruptly stop taking this medication to prevent dysrhythmias, angina, MI</li> <li>3. Caution when changing positions or getting OOB, use call light</li> <li>4. Do not administer medication when heart rate is less than 60bpm</li> </ol>
Sucralfate	Antiulcer drug	Creates a protective barrier against gastric acids	1,000mg PO QID before meals	Yes Choose an item. Click here to enter text.	N/A	Dry mouth, constipation	<ol style="list-style-type: none"> <li>1. Take medication on an empty stomach one hour before meals and bedtime</li> <li>2. Avoid the use of antacids, alkaline water within 1 hour of administration</li> <li>3. Increase fluids and fiber rich foods (whole grain, apples) to prevent constipation</li> <li>4. Assess GI symptoms such as ABD pain, blood in stools</li> </ol>
Zonisamide	Anticonvulsant	Adjunctive therapy for spartial eizures	100mg cap BID	Yes Choose an item. Click here to enter text.	N/A	Seizures, suicidal ideation, irritation, aplastic anemia	<ol style="list-style-type: none"> <li>1. Do not discontinue medication abruptly as seizures may occur</li> <li>2. Notify of rash immediately, abdominal pain, back pain, blood in urine</li> </ol>

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT	Date: 1/29/2025	Allergies: Adhesive tape, Cefedinir, Lactose Intolerant		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							3. Increase fluid intake to reduce risk of kidney stones 4. Report suicidal thoughts and behaviors immediately
Ketorlac	Antiinflammatory NSAID	Antiinflammatory for use before or after surgery	10mg tab PRN every six hours PO	Yes Choose an item. Click here to enter text.	N/A	BBW: Bleeding, MI, Stroke, limit duration of use. Seizures, nephrotoxicity	1. Report blood in urine/tarry stools, bruising 2. Report blurred vision, ringing/roaring in ears (may indicate toxicity) 3. Will increase bleeding times. Use soft bristled toothbrush, electric razor, avoid any activities that may cause trauma 4. Avoid use with other anticoagulants such as aspirin and other NSAIDS
Diphenhydramine	Antihistamine	Treat allergy symptoms, decreases itching	25mg cap PO every six hours PRN	Yes Choose an item. Click here to enter text.	N/A	Dizziness, drowsiness, seizures, hypotension, thrombocytopenia, urinary retention	1. Use hard sugar free candy, gum, frequent rinsing of mouth for dry mouth 2. Use caution when getting out of bed or ambulating as dizziness may occur 3. Assess for urinary retention, dysuria 4. May increase CNS depression, report confusion, sedation, and hypotension
Acetaminophen	analgesic, antipyretic	Mild to moderate pain	1,000mg tab PO	Yes Choose an	N/A	BBW: May cause	1. Assess for fever/pain (type of pain, location, temp, duration)

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Jasmine Luna		Unit: Pediatric Floor 3	Patient Initials: NT	Date: 1/29/2025	Allergies: Adhesive tape, Cefedinir, Lactose Intolerant		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
		or fever reliever	every six hours PRN	item.Click here to enter text.		hepatotoxicity, GI bleeding, rash, thrombocytopenia	<ol style="list-style-type: none"> <li>2. Inform patients that urine may become dark brown d/t phenacetin (metabolite of acetaminophen)</li> <li>3. Do not exceed max dosage as liver damage may result, report s/s of N/V, ABD pain</li> <li>4. Report bleeding, bruising, malaise, fever, sore throat as these may be s/s of chronic overdose</li> </ol>
Polyethylene glycol solution	Laxative	Constipation	2L/hr per NG tube	Yes Choose an item.Click here to enter text.	Reconstituted in 4L of water	Nausea, abdominal fullness/bloating, dry mouth, dizziness	<ol style="list-style-type: none"> <li>1. Increase fluid intake d/t increase amount of diarrhea and to prevent dehydration</li> <li>2. Assess electrolytes of electrolyte imbalance</li> <li>3. Report signs of dry mouth, dark urine, reduced urination, and tachycardia as these s/s may indicate dehydration</li> <li>4. Increase hygiene measures to prevent skin breakdown</li> </ol>
Fluoxetine	Antidepressant, SSRI	Inhibits CNS neuron uptake of serotonin but not of	20mg cap PO daily	Yes Choose an item.Click here to	N/A	BBW: Children, suicidal ideation, seizures,	<ol style="list-style-type: none"> <li>1. Therapeutic effect may take up to 2-6 weeks</li> <li>2. May cause dizziness (orthostatic hypotension), Use call light for help OOB/</li> </ol>

## Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

<b>Student Name:</b> Jasmine Luna		<b>Unit:</b> Pediatric Floor 3	<b>Patient Initials:</b> NT		<b>Date:</b> 1/29/2025	<b>Allergies:</b> Adhesive tape, Cefedinir, Lactose Intolerant	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Is med in therapeutic range? If not, why?	IVP - List diluent solution, volume, and rate of administration  IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
		norepinephrine		enter text.		bradycardia, torsades de pointes, hemorrhage, serotonin syndrome	Rise or change positions slowly 3. Report suicidal ideation as suicidal thoughts/behavior may occur 4. Report fever, sweating, diarrhea, poor coordination, N/V, flushing as these are S/S of Serotonin syndrome