

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

Student Name: Click here to enter text.		Unit: Click here to enter text.		Patient Initials: Click here to enter text.		Date: Click here to enter a date.		Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)		
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	Choose an item. Click here to enter text.	N/A	- Fast or pounding HR, seizures, headaches with chest pain, and severe dizziness	1. Talk to your doctor if you become pregnant as this med may cause birth defects. 2. Immediately report the new onset of tremors, seizures, or restless muscles. 3. Do not drink with alcohol because it can increase your risks of side effects. 4. Avoid activities that require a high level of alertness while taking this med because it may cause impaired thinking.		
lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	Choose an item. Click here to enter text.	N/A	- insomnia, Respiratory arrest, vision changes, severe drowsiness, and jaundice	1. Do not take with other CNS depressants such as opioids or alcohol. 2. Do not stop taking abruptly without speaking to Dr. 3. This med is addictive and should be taken as prescribed. 4. Do not take while pregnant or it can cause birth defects or NB withdrawal symptoms.		

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: <small>Click here to enter text.</small>		Unit: <small>Click here to enter text.</small>	Patient Initials: <small>Click here to enter text.</small>	Date: <small>Click here to enter a date.</small>	<small>Click here to enter text.</small>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
haldol	antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	Choose an item. <small>Click here to enter text.</small>	N/A	PMS Symptoms (tardive dyskinesia), rapid mood swings, pounding heart beat	<ol style="list-style-type: none"> 1. Report having a pounding or fast HR and SOB 2. Do not quit taking abruptly, take as directed 3. It may take a few weeks to notice the benefits of this med. 4. Avoid alcohol because it can cause side effects. Reactions could be impaired while taking.
diphenhydramine	antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	Choose an item. <small>Click here to enter text.</small>	N/A	Painful or little urination, confusion, and tightness in neck.	<ol style="list-style-type: none"> 1. May cause dizziness or drowsiness so avoid driving or high alert activities. 2. Be cautious when drinking alcohol because it can increase drowsiness 3. Immediately report EPS symptoms to your HCP 4. Take this med as instructed and prescribed.
acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-4 on pain scale	Choose an item. <small>Click here to enter text.</small>	N/A	Overdose can lead to severe liver disease.	<ol style="list-style-type: none"> 1. Do not take more than 4000mg within 24 hours. 2. Immediately report if you develop upper right side stomach pain 3. Avoid taking other meds that have

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: <small>Click here to enter text.</small>		Unit: <small>Click here to enter text.</small>	Patient Initials: <small>Click here to enter text.</small>	Date: <small>Click here to enter a date.</small>	<small>Click here to enter text.</small>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							acetaminophen in them to prevent overdose. 4. Do not drink alcohol while taking acetaminophen, this will lead to increased risk of liver damage.
olanzapine	antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	Choose an item. <small>Click here to enter text.</small>	N/A	Increased risk of stroke or death in the elderly patients with dementia	<ol style="list-style-type: none"> 1. Immediately report thoughts of suicide, especially if taking with fluoxetine. 2. Be aware and report stiffness in the neck and uncontrolled muscle movements in face-tardive dyskinesia. 3. warn of dizziness and educate on getting up slowly from bed to prevent orthostatic hypotension. 4. Manage blood sugar to make sure it is not too high. Look for excessive thirst and hunger.
ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	Choose an item. <small>Click here to enter text.</small>	N/A	Overdose can lead to stomach or intestine damage. May increase risk of fatal heart attack or stroke.	<ol style="list-style-type: none"> 1. Immediately report chest pain, slurred speech, SOB, or weakness on one side of the body. 2. Immediately report bloody or tarry stools, this may look like coffee grounds and signifies intestinal bleeding 3. It is common to develop N&V and

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: <small>Click here to enter text.</small>		Unit: <small>Click here to enter text.</small>	Patient Initials: <small>Click here to enter text.</small>	Date: <small>Click here to enter a date.</small>	<small>Click here to enter text.</small>		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
							headache while taking ibuprophen. 4. Do not take with antidepressants because it may increase your bleeding risk.
risperdal	antipsychotic	Psychotic behavior/bipolar mania	1 mg tablet PO BID	Choose an item. <small>Click here to enter text.</small>	N/A	Low WBC's and platelets, and high blood sugar	1. Avoid moving too fast or dizziness may develop 2. Do not take with alcohol because it increases your risks of dangerous side effects 3. Drink plenty of fluids and avoid becoming too hot or too cold. 4. Report new onset of depression.
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	Choose an item. <small>Click here to enter text.</small>	N/A	Allergic reaction	1. Immediately report itching, hives, rash, or difficulty breathing. 2. Store these patches at room temperature and dispose of it properly. 3. The patch can sometimes cause flu-like signs, back pain, or trouble sleeping. Notify HCP 4. If an overdose occurs call poison control center!

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: <small>Click here to enter text.</small>		Unit: <small>Click here to enter text.</small>	Patient Initials: <small>Click here to enter text.</small>		Date: <small>Click here to enter a date.</small>	<small>Click here to enter text.</small>	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
ziprasidone	antipsychotic	Psychotic behavior/bipolar disorder/mania	40 mg PO BID	Choose an item. <small>Click here to enter text.</small>	N/A	Drug interactions and allergic reaction	<ol style="list-style-type: none"> 1. Stop taking and contact HCP if you have a fast pounding fluttering in your chest. 2. Do not use if you have a coexisting heart condition such a prolonged QT interval, or heart failure. 3. Check with HCP to make sure all your meds are compatible with ziprasidone. 4. Take as directed by prescriber.
prazosin	Alpha-blocker	Nightmares	1 mg PO QHS	Choose an item. <small>Click here to enter text.</small>	N/A	hypotension	<ol style="list-style-type: none"> 1. Store at room temperature and prevent it from being exposed to moisture or heat. 2. Do not take if under the age of 18 years old. 3. Try to stay on top of managing blood pressure and check it often to make sure all is good. 4. Immediately report extreme drowsiness or underactive reflexes.
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	Choose an item. <small>Click here to enter text.</small>	N/A	Hypertension leading to headache, passing out, or changes in	<ol style="list-style-type: none"> 1. Do not use if your are pregnant or it may cause harm to the baby. 2. provide resources such as counseling to stop smoking. 3. Use as instructed. Do not overdose- if

Adult/Geriatric Medication Worksheet – Current Medications & PRN for Last 24 Hours

Student Name: Click here to enter text.		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List diluent solution, volume, and rate of administration IVPB – List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						eyesight.	you do, call poison control center. 4. Speak to HCP about how to dispose of expired or unused drugs. (throw it away)
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.