

IM 7 Math Module

Complete the required math problems and check your answers.

Drop Factor Problems

1. Infuse 1.5 gram of a medication over 3 hours. The drug is supplied as 1.5 gram/250mL. The drip factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$\frac{250 \text{ mL} \times 15}{180 \text{ min}} = 20.83 \approx \boxed{21 \text{ gtt/min}}$$

2. An order has been written to give 1 gram of a medication over 30 minutes. The drug is supplied as 1 gram/50mL. The gtt factor is 60. How many gtt/min will you infuse?

$$\frac{50 \text{ mL} \times 60}{30} = \boxed{100 \text{ gtt/min}}$$

3. The nurse is to give 500mg IV of a medication over 1 hr. The drug is supplied as 1 gram/250mL. The gtt factor is 15. How many gtt/min will you infuse? (Round to the nearest whole number)

$$\frac{500 \text{ mg}}{1000} (250 \text{ mL}) = 125 \text{ mL} \quad \frac{125 \text{ mL} \times 15}{60 \text{ min}} = 31.25 \approx \boxed{31 \text{ gtt/min}}$$

4. The patient is to receive 400 mg IV of a medication over 1 hour. You receive an IV bag from the pharmacy labeled 400 mg in 100 mL D5W. The IV tubing delivers 12 gtt/mL. How many drops per minute (gtt/min) will the nurse deliver?

$$\frac{100 \text{ mL} \times 12}{60} = \boxed{20 \text{ gtt/min}}$$

mL/hr Infuse over time

5. The physician writes an order to give 1000mL of intravenous fluid over 8hrs. How many mL/hr will you infuse?

$$\frac{1000 \text{ mL}}{8 \text{ hr}} = \boxed{125 \text{ mL/hr}}$$

6. Infuse 1000 mLs of intravenous fluid over 4 hrs. How many mL/hr will you set on the pump?

$$\frac{1000 \text{ mL}}{4 \text{ hr}} = \boxed{250 \text{ mL/hr}}$$

7. A physician orders 1000 mg of a medication to be given every 6 hours over 1.5 hours. The medication is delivered with 1000 mg in 250 mL. How many mL/hr will you set the pump? (Round to the nearest whole number)

$$\frac{250 \text{ mL}}{1.5 \text{ hr}} = 166.66 \approx \boxed{167 \text{ mL/hr}}$$

IV Push

8. An order is received for 75mcg IV push of a medication now. The drug is supplied as 100mcg/2mL. How many mL will you give?

$$\frac{75 \text{ mcg}}{100 \text{ mcg}} (2 \text{ mL}) = \boxed{1.5 \text{ mL}}$$

9. The patient is to receive 5mg of a medication IV push. The drug is supplied as 20mg/5mL. How many mL will you give? (Do not round your final answer)

$$\frac{5 \text{ mg}}{20 \text{ mg}} (5 \text{ mL}) = \boxed{1.25 \text{ mL}}$$

10. The order is to give 2mg IV push of a medication now. The drug is supplied as 10mg/1mL. How many mL will you give?

$$\frac{2 \text{ mg}}{10 \text{ mg}} (1 \text{ mL}) = \boxed{0.2 \text{ mL}}$$

IM 7 Math Module

Complete the required math problems and check your answers.

Mcg/kg/min or Mcg/min

$$142 \text{ mL} \times 100 = 14200 \text{ mcg/hr}$$

11. The patient is receiving an intravenous medication currently infusing at 142 mL/hr. The IV bag of reads 50 mg in 500 mL D5W. The patient weighs 175 lbs. How many mcg/kg/min are infusing? (Round to the nearest tenth)

$$\div 60 \text{ min} \\ 236.666...$$

$$79.5 \text{ kg}$$

$$2.972$$

$$3 \text{ mcg/kg/min}$$

12. The physician has ordered a medication that states to start at 1 mcg/kg/min and titrate as needed. The IV bag of medication contains 250 mg in 500 mL D5W. The patient weighs 70 kg. How many mL/hr should the IV pump be set at to achieve the starting dose? (Round to the nearest whole number)

$$\frac{4200 \text{ mcg} (2 \text{ mL})}{1000 \text{ mcg}} = 8.4 \approx 8 \text{ mL/hr}$$

13. The patient is currently receiving a medication at 12 mL/hr. The bottle reads 100 mg in 250 mL D5W. How many mcg/min is the patient receiving?

$$0.2 \text{ mL/min}$$

$$400 \text{ mcg/mL}$$

$$400 \times 0.2 = 80 \text{ mcg/min}$$

14. The physician has ordered a medication to start at 2 mcg/kg/min. The patient weighs 165 lbs. The IV bag reads 800 mg in 500 mL D5W. What rate would the nurse set on the infusion pump? (Round to the nearest tenth)

[mL/hr]

$$1.6 \text{ mg/mL}$$

$$0.002 \text{ mg/kg/min} = 0.12 \text{ mg/hr} \times 75 \text{ kg} = 9 \text{ mg/hr}$$

$$\frac{9 \text{ mg (mL)}}{1.6 \text{ mg (hr)}} = 5.625 \approx 5.6 \text{ mL/hr}$$

15. The physician has written an order to increase the medication to 4 mcg/kg/min. The IV bag reads 64 mcg/mL. What rate would the nurse set on the IV pump? (Round to the nearest tenth)

[mL/hr]

$$\frac{240 \text{ mcg (mL)}}{64 \text{ mcg}} = 3.75 \approx 3.8 \text{ mL/hr}$$

16. The patient is on a medication drip infusing at 15 mL/hr. The label reads 50 mcg/mL. The patient weighs 65 kg. How many mcg/min is the patient receiving? (Do not round)

$$0.25 \text{ mL/min}$$

$$50 \text{ mcg} \times 0.25 \text{ mL/min} = 12.5 \text{ mcg/min}$$

Heparin/Insulin or mg/hr

17. The physician orders a heparin infusion at 500 units/hr. The IV bag of medication reads 25,000 units in 250 mL D5W. How many mL/hr should be showing on the IV pump?

$$(100 \text{ unit/mL}) \times 5 =$$

$$5 \text{ mL/hr}$$

$$0.4 \text{ u/mL}$$

18. The patient is on a regular insulin drip infusing at 5 units/hr. The bag is labeled 100 units in 250 mL NS. At what rate should the pump be infusing? Round to the nearest whole number.

$$0.4 / 5 \text{ units} = 12.5 \text{ mL/hr} \approx 13 \text{ mL/hr}$$

Burns (Parkland Formula) Do not round weights

$$4 \text{ mL} \times \% \text{ TBSA} \times \text{weight (kg)}$$

19. A 200-pound patient presents to the emergency department with 50% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$4 \text{ mL} \times 50 \times 90.9 = 18180 \text{ mL} = 18 \text{ L}$$

20. A 150-pound patient presents to the emergency department with 75% total body surface area (TBSA) burn. How many liters of fluid would be given in the first 24 hours? (Round to the nearest whole number)

$$(68.2 \text{ kg})$$

$$4 \text{ mL} \times 75 \times 68.2 = 20460 \text{ mL} = 20 \text{ L}$$