

Student Name: Emily Arismendez

Unit: \_\_\_\_\_

Pt. Initials: BO

Date: \_\_\_\_\_

**Medication Worksheet - Current Medications & PRN for Last 24 Hours**

Allergies: \_\_\_\_\_

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List solution to dilute and rate to push. IVPB - List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Paliperidone	antipsychotic	For command hallucinations	234mg IM Day 1  156mg Day 8	Y  N		Sedation, dizziness, akathisia, and extrapyramidal	1. Do not rub site of injection after application 2. warning: increased mortality for elderly patients with dementia 3. Make sure to report back in 8 days to receive 156mg on day 8 4. Make sure to take 117 mg maintenance dose
Lorazepam	Benzodiazepines	Anxiety	0.5mg 1 tab Q 12 hrs PO	Y  N		Dizziness  Confusion  Blurred vision  nausea	1. Report any unusual movements or tremors 2. Do not drive if med causes blurred vision and report this symptom immediately to HCP 3. Drink 3 L of water daily due to nausea and vomiting or diarrhea. 4. Do not drink alcohol since both will depress the CNS even more.
Haloperidol	antipsychotic	Schizophrenia	5mg 1 tab PO at bedtime	Y  N		Akathisia, tachycardia, shortness of breath, low white blood cells, cough with mucus	1. High dose or long term use can cause akathisia. 2. Get up or move slowly in order to not fall and hurt yourself. 3. Do not take if you have Parkinson's 4. Do not take with alcohol.

Student Name: Emily Arismendez

Unit: \_\_\_\_\_

Pt. Initials: BO

Date: \_\_\_\_\_

**Medication Worksheet - Current Medications & PRN for Last 24 Hours**

**Allergies:** \_\_\_\_\_

Tramadol	opioid	Withdrawal pain	50 mg 1 tab PO q 6 hrs	Y N		Headaches Dry mouth Constipation sweating	<ol style="list-style-type: none"> <li>DO not operate heavy machinery while taking this med.</li> <li>Contact HCP if respiratory depression occurs</li> <li>Change positions slowly due to orthostatic hypotension</li> <li>Take with a magnesium or other laxative to avoid constipation</li> </ol>
Trazodone	antidepressant	MDD	100mg 1 tab PO at bedtime	Y N		Mood behavior changes Trouble sleeping Slow heartbeats Unusual thoughts/behaviors	<ol style="list-style-type: none"> <li>Don't take with Methylene blue injection</li> <li>Don't take if you have taken an MAOI in the last 14 days</li> <li>report any mood changes such as panic attacks, agitation, and restlessness</li> <li>Stop if suicidal ideations occur</li> </ol>