

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: Codeine, Proton Pump Inhibitors, sumatriptan, Versed (Midazolam)

Primary IV fluid and rate: Lactated Ringers 100mL/hr.

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Hyoscyamine (LEVSIN)	Tab 0.125mg Sublingual every 4 hrs.	N/A	Peptic Ulcer	<ol style="list-style-type: none"> 1. Take tab let it dissolve under tongue; do not split, chew, or crush due to losing its effect 2. DO NOT take with antacids may interfere with hyoscyamine, making it less effective 3. Can cause dizziness & drowsiness, avoid standing up or sitting down too quickly due to med relaxing muscles in digestive tract
Ketorolac (TORADOL)	Injection 15mg IV every 6 hrs.	IVP Diluent: Sterile Water Volume: 15 mg Rate: slow push rate	Short-term relief of acute, moderately severe pain	<ol style="list-style-type: none"> 1. DO NOT lie down for about 15-30 mins after admin. helps to prevent irritation that may lead to trouble in swallowing 2. Contraindicated in pts with renal disease or renal failure because it may increase fluid retention and worsen renal function 3. NURSING: monitor BP due to med. increasing BP during treatment
Acetaminophen (OFIRMEV)	1,000 mg IV	IVPB Concentration: Lactated Ringers Rate: 400 mL/hr. (infused for 15mins)	Moderate Pain	<ol style="list-style-type: none"> 1. Taking too much can cause liver damage, report yellowing of skin & eyes 2. Can lead to constipation, increase in fiber & drink lots of water since med. Can have an effect to digestive system 3. Can cause N/V, abdominal pain, confusion due to intaking more doses than recommended amount causing toxin build up.

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: Codeine, Proton Pump Inhibitors, sumatriptan, Versed (Midazolam)

Primary IV fluid and rate: Lactated Ringers 100mL/hr.

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Diatrizoate (GASTROGRAFIN)	Liquid 120 mL PO daily	N/A	Radiographic examination of the gastrointestinal tract	<ol style="list-style-type: none"> 1. Inform physician if pregnant because it can cross the placenta and potentially affect the fetus 2. REPORT if allergic to iodine, any foods, or x-ray materials due to having the possibility to affect future thyroid tests 3. REPORT rash, hives, itching, SOB, any kind of swelling which determines an allergic reaction
Methocarbamol (ROBAXIN)	Injection 1,000 mg IV every 8 hrs.	IVP Diluent: Sterile Water Volume: 1 mL Rate: Slow push rate	Relax muscle & relieve pain from injury	<ol style="list-style-type: none"> 1. Can cause dizziness, drowsiness, alertness or coordination due to being a muscle relaxer 2. AVOID alcoholic drinks due to also causing drowsiness which increases pt to have more side effects 3. If have an upset stomach or nausea, take food with med. to help with absorption
				<ol style="list-style-type: none"> 1. 2. 3.

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: Codeine, Proton Pump Inhibitors, sumatriptan, Versed (Midazolam)

Primary IV fluid and rate: Lactated Ringers 100mL/hr.

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):