

64

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies: **Baclofen**

Primary IV fluid and rate: **NA**

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):
NA

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Amlodipine	5mg po daily 1 tab	NA	hypertension	<ol style="list-style-type: none"> It is important for you to take this medication as well as take other actions to help your HTN such as losing weight, adopting a low sodium diet, stop smoking and exercise regularly. This medication will not cure your HTN, only helps control it. It is important for your to maintain good oral hygiene while on this medication. Make sure you are keeping up with your dentist appointments as this medication can cause gum enlargement, bleeding and tenderness. This medication can cause irregular heart beats, shortness of breath, and swelling of your hands and feet. Maybe even dizziness, nausea or constipation. Please notify your HCP if you experience any of these symptoms.
Atorvastatin	10mg po daily 1 tab	NA	hyperlipidemia (hx)	<ol style="list-style-type: none"> Avoid foods high in fat/cholesterol as this will increase your already elevated levels of cholesterol. Avoid fried foods. This medication should be used with diet restriction, exercise, and smoking cessation for best results. Notify your HCP if you experiences any new muscle pain, tenderness, weakness. If fever, feeling sick/weak are also accompanied, This medication make take up to 2 weeks to improve your cholesterol. It is important to continue to take this medication and to keep you follow up appointments to review the effectiveness.
Ceftriaxone	1g IV push daily	1g/10ml reconstituted in sterile water IVP over 3-5min	abx	<ol style="list-style-type: none"> You are currently taking this antibiotic as a preventative measure while your urine/blood samples are being tested. Blood in urine, and yeast present does indicate an infection. It is important to report signs of continues or new infection such as vaginal it hits, discharge or diarrhea. This medication can cause C. Diff in some patients. Please notify your HCP if you notice any blood in your stool. If your stools are become loose and or foul smelling.
Clozapin	25mg po daily 1 cap	NA	antidepressant	<ol style="list-style-type: none"> This medication does increases your chances of becoming drowsy, and the sedation effects make it important to continue to take this medication as prescribed, nightly. This medication can cause sleep walking, which you may or may not remember when you wake up the next morning. Please inform your HCP if you experience any episodes similar immediately. This medication can cause dry mouth and or constipation. You can increase your fluid intake or eat sugarless candy to help. If you notice these symptoms with no relief please notify your healthcare provider.

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
<i>duloxetine</i>	<i>60mg po daily</i>	<i>NA</i>	<i>Anti depressant</i>	<ol style="list-style-type: none"> 1. This medication can cause your blood pressure to drop when you get up, please call for help so we can make sure you can ensure you have your walker available, and to prevent any falls. 2. This medication may cause drowsiness. It is important to not avoid driving or activities that requires her to be alert. 3. If she starts to feel any anxiety, agitation, insomnia, irritability, or worsening of her depression, to please inform your HCP.
<i>enoxaparin</i>	<i>40mg/aml SQ daily</i>	<i>NA</i>	<i>Anti coagulant</i>	<ol style="list-style-type: none"> 1. Notify HCP of unusual bleeding, gums, nose, stool, excess in urine. Call for help getting out of bed. We will make sure your walker is nearby to prevent any falls/injury. 2. Do not take with aspirin or ibuprofen without taking to your HCP first. These are also blood thinners and may cause complications. 3.
				<ol style="list-style-type: none"> 1. 2. 3.
				<ol style="list-style-type: none"> 1. 2. 3.

REFERENCES:
 update
 drugs.com
 Davis's Drug Guide

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 12pm prior day to 12pm current day

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
				1. 2. 3.
				1. 2. 3.

Medication reference: