

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

| | | | | |
|--|---|---------------------------|-------------------------------------|---------------------------------|
| Primary IV Fluid and Infusion Rate (ml/hr) | Circle IVF Type | Rationale for IVF | Lab Values to Assess Related to IVF | Contraindications/Complications |
| Click here to enter text. | Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/> | Click here to enter text. | Click here to enter text. | Click here to enter text. |

| Student Name: Click here to enter text. | | Unit: Click here to enter text. | Patient Initials: Click here to enter text. | | Date: Click here to enter a date. | Allergies: Click here to enter text. | |
|---|------------------------------|---|---|---|---|--|---|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| Hydroxyzine pamoate | Antihistamine | Anxiety | 25mg PO PRN q6 | Choose an item. | | Chest pain, tightness. Drowsy, dizzy. Fast or irregular heartbeat. | 1. Do not drive or use machinery if you feel drowsy. 2. Contraindicated in patients with long QT intervals. 3. Contraindicated in early pregnancy and do not take while breastfeeding. 4. Alcohol and cannabis can make you drowsier. |
| lorazepam | Benzodiazepine | Alcohol withdrawal / anxiety | 1-2 mg PO PRN q4 hours | Choose an item. Choose an item. | | Confusion, blurred vision, dizziness, weakness, drowsiness, dry mouth, restless. | 1. Do not drive or use machinery if you are not alert or able to think or see well. 2. Should only be prescribed lowest dose for for short term relief. 3. Report suicide thoughts. 4. May be dangerous for people with acute asthma, liver or kidney disease. |
| haldol | antipsychotic | for psychotic symptoms/agitation | 10 mg IM PRN q8 | Choose an item. | | Hypotension, constipation, dry mouth, blurred vision, | 1. Get up slowly to decrease feeling lightheaded. 2. Contraindicated in PD. |

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

| Student Name: Click here to enter text. | | Unit: Click here to enter text. | Patient Initials: Click here to enter text. | | Date: Click here to enter a date. | Allergies: Click here to enter text. | |
|--|------------------------------|--|--|--|---|---|--|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| | | | | Choose an item. | | insomnia, drowsy. | 3. Make sure you don't feel drowsy if blurred vision before driving. 4. Eat high fiber for constipation and drink plenty of fluids. |
| diphenhydramine | antihistamine | management of cholinergic effects of antipsychotic | 50 mg IM q8 PRN psychotic symptoms | Choose an item. | | Loss of appetite, N/V, dizzy, drowsiness, dry mouth, constipation. | 1. Contraindicated in HTN. 2. Use alcohol cautiously. 3. Caution when driving or performing hazardous activities. 4. Should not be used in older adults because it is not safe or as effective. |
| acetaminophen | Analgesic/antipyretic | Mild-moderate pain | 500 mg PO q6 for pain of 1-4 on pain scale | Choose an item. | | Rash, itching, hives, swelling of face, throat, lips, difficulty breathing or swallowing. | 1. Do not take more than 4000mg in 24hrs. 2. Can cause liver damage. 3. Take it as prescribed by your doctor. 4. Report any difficulty breathing or tightness of chest. |
| olanzapine | antipsychotic | Agitation/psychotic behavior | 5 mg PO q6 for agitation | Choose an item. | | Feeling sleepy during the day, increase in appetite, feeling dizzy, constipation. | 1. Do not drive feeling sleepy throughout the day. 2. Drinking alcohol can increase drowsiness. 3. Contraindicated in patients with known hypersensitivity to the product. |

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

| Student Name: Click here to enter text. | | Unit: Click here to enter text. | Patient Initials: Click here to enter text. | | Date: Click here to enter a date. | Allergies: Click here to enter text. | |
|--|------------------------------|------------------------------------|--|--|---|---|---|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| | | | | | | | 4. Avoid blood pressure medications. |
| ibuprofen | NSAID | Mild-moderate pain | 400 mg PO q6 for pain of 1-4 on pain scale | Choose an item. Choose an item. | | HA, N/V, dizziness. | <ol style="list-style-type: none"> 1. Drink plenty of fluids for HA. 2. Be careful standing up too fast. 3. Eat simple meals for nausea. 4. Can cause kidney damage. |
| risperdal | antipsychotic | Psychotic behavior/bipolar mania | 1 mg tablet PO BID | Choose an item. | | N/V, diarrhea, heartburn, constipation, dry mouth, increased appetite/saliva. | <ol style="list-style-type: none"> 1. Can cause toxicity in those with renal impairment. 2. Avoid drinking alcohol 3. Report motor function especially tardive dyskinesia because may be irreversible. 4. Drink fluids for dry mouth, eat fiber for constipation. |
| Nicotine patch | Stimulant | Nicotine replacement | 21 mg transdermal patch daily | Choose an item. Choose an item. | | HA, dizziness, N/V, diarrhea, redness or swelling at the patch site. | <ol style="list-style-type: none"> 1. Be cautious when applying and wash hands after to wash away any nicotine you may have gotten on your fingers. 2. Put patch on different area to avoid skin irritation. 3. Avoid wearing patch in the same place more than once per week. 4. When removing clean area with alcohol |

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

| Student Name: Click here to enter text. | | Unit: Click here to enter text. | Patient Initials: Click here to enter text. | | Date: Click here to enter a date. | Allergies: Click here to enter text. | |
|--|------------------------------|---|--|---|---|--|---|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| | | | | | | | wipe to remove any nicotine left. |
| ziprasidone | antipsychotic | Psychotic behavior/bipolar disorder/mania | 40 mg PO BID | Choose an item. | | Loss of balance, drooling, fear/nervousness, inability to sit still, muscle trembling/stiffness. | <ol style="list-style-type: none"> 1. Drink more water and stay out of the sun. 2. Be careful if you exercise often or exposed to high temps./humidity. 3. May cause some suicidal thoughts. 4. Contraindicated in heart conditions. |
| prazosin | Alpha-blocker | Nightmares | 1 mg PO QHS | Choose an item. Choose an item. | | Dizziness, drowsiness, less alert, low BP, heart palpitations. | <ol style="list-style-type: none"> 1. Do not drive if you feel drowsy. 2. Monitor BP regularly 3. Avoid use for HTN 4. Get up slowly or sit down for a minute if you feel dizzy. |
| Nicotine inhaler | Stimulant | Nicotine replacement | 10 mg cartridge for inhalation Q2 | Choose an item. | | Fast irregular heartbeat, HA, runny nose, nausea, tightness of chest, skin rash, itching, tearing of the eyes. | <ol style="list-style-type: none"> 1. Do not eat or drink for 15 mins before using or during use. 2. Never use more than the prescribed amount. 3. Keep out of reach of children. 4. Avoid smoking, tobacco products while using inhaler. |

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

| Student Name: Click here to enter text. | | Unit: Click here to enter text. | Patient Initials: Click here to enter text. | | Date: Click here to enter a date. | Allergies: Click here to enter text. | |
|---|------------------------------|---|---|---|---|--|--|
| Generic Name | Pharmacologic Classification | Therapeutic Reason | Dose, Route & Schedule | Correct Dose? If not, what is correct dose? | IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration | Adverse Effects | Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.) |
| Click here to enter text. | Click here to enter text. | Click here to enter text. | Click here to enter text. | Choose an item. | Click here to enter text. | Click here to enter text. | 1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text. |
| Click here to enter text. | Click here to enter text. | Click here to enter text. | Click here to enter text. | Choose an item. | Click here to enter text. | Click here to enter text. | 1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text. |