Student Name: Brianna Meyer Date: 5/6/25

IM6 (Acute Psychiatric) Critical Thinking Worksheet

1. DSM-5 Diagnosis and Brief Pathophysiology (include reference):	2. Psychosocial Stressors (i.e. Legal, Environmental, Relational, Developmental, Educational, Substance Use, etc.);	3. DSM-5 Criteria for Diagnosis (Asterisk or Highlight Symptoms Your Patient Exhibits and Include References)
· Schizophrenia · chronic psychiatric disorder involving dysregulation of dopamine pathways in brain causing hallucinations & delusions	· Living arrangements · charges	A. · delusions B. significant portion of time level of fxn is marked below level C. cont. signs of disturbances lasting <6 mo e. disturbances not r/t drug of abuse
4. Medical Diagnoses: ~~MDD~~ ~~Schizophrenia~~		
5. Diagnostic Tests Pertinent or Confirming of Diagnosis · QSPS: 5 yes; 3 no · Colombia: low risk	**6. Lab Values That May Be Affected:** N/A	**7. Current Treatment:** ~~group therapy~~ Meds ~~1:1 therapy~~ · Close obs. q15 · milieu therapy · V/S · nursing/group session · ADLS assist · ind. support · reality orientation · encourage disc. · wt monitoring

Adopted: August 2016, revised October 2018

Student Name: Brianna M Unit: Oceans Pt. Initials: D Date: 5·6·25

Allergies: NKDA

Medication Worksheet – Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type		Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
N/A	Isotonic/ Hypotonic/ Hypertonic		N/A	N/A	

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Quetiapine fumarate	atypical antipsychotic (2nd gen)	severe psychosis	100mg PO,D	(Y) N	N/A	Orthostatic hypotension · Wt gain · EPS · tardive dyskinesia	1. monitor wt 2. monitor glucose/lipid levels 3. watch for neuroleptic malignant S/S 4. take as prescribed even if symptoms improve
fluphenazine HCl	typical antipsychotic (1st gen)	severe psychosis	5mg	Y N		· EPS · sedation/drowsy · photosensitivity · orthostatic	1. monitor mental status (delusions) 2. assess for EPS/tardive 3. V/S esp. BP, SPF protection 4. eval LFT periodically ★ ↑ mortality in elderly
venlafaxine	antidepressant	depressive disorder (major)	37.5 mg	Y N		· serotonin syndrome · HTN · insomnia	1. take w/ food to prevent nausea 2. 2-4 wks for full effect 3. avoid alcohol or other depressants 4. monitor BP ★ suicidal thoughts
oxcarbazepine		BPD	150mg	Y N		· ataxia · hyponatremia · dizzy · fatigue	1. monitor Na+ levels 2. take as prescribed 3. don't stop abruptly 4. report any rashes/skin changes
				Y N			1. a sap 2. 3. 4.