

Reconstitution for IV Push

Medication	Reconstitution	Diluent	Comments	Rate of IV Push Administration
acetaZOLAMIDE (Diamox)	500 mg in 5 mL	Sterile Water	Shake vigorously	3-5 minutes
alteplase (Cathflo)	2 mg in 2 mL	2.2 mL Sterile Water	DO NOT SHAKE (gentle swirl)	Over 1 minute
aztreonam (Azactam)	1 gram in 10 mL 2 grams in 10 mL	Sterile Water or Normal Saline	Shake vigorously	2-4 minutes
calcium gluconate	1 gram/10 mL (liquid)	Undiluted	N/A	Over 10 minutes
ceFAZolin (Ancef)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water or Normal Saline	Shake vigorously	2-4 minutes
cefoTEtan (Cefotan)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water	Shake Vigorously	2-4 minutes
cefuroxime (Zinacef)	750 mg in 10 mL 1.5 grams in 20 mL	Sterile Water or Normal Saline	Shake Vigorously	2-4 minutes
cefTRIAxone (Rocephin)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water or Normal Saline	Shake Vigorously	2-4 minutes
cefTAZidime (Tazicef)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water	Shake Vigorously	2-4 minutes
cefepime (Maxipime)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water or Normal Saline	Shake Vigorously	2-4 minutes
methocarbamol (Robaxin)	1 gram/10 mL (liquid)	Undiluted	N/A	3-5 minutes
meropenem (Merrem)	1 gram in 10 mL 2 grams in 20 mL	Sterile Water or Normal Saline	Shake Vigorously	3-5 minutes
pantoprazole (Protonix)	40 mg in 10 mL	Normal Saline	Shake Vigorously	2-4 minutes