Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Vancomycin	N/A	N/A	Usually bacterial infections	1. BBW: risk of fetal toxicity. Do not take in pregnancy. 2. Risk of C. Diff infection. Monitor for foul smelling stool 3. Risk of nephrotoxocity. Monitor I&O's closely. Report any radiating back pain. 4. Can cause drug-induced immune thrombocytopenia. Monitor for unusual bleeding and hemorrhage.
Metoprolol	N/A	N/A	Chronic stable angina, rate control of atrial fibrillation/flutter, heart failure with reduced ejection fraction, chronic HTN, early treatment or prevention of MI, supraventricular tachycardia, ventricular arrhythmias	1. Use with caution with DM because it can mask signs of hypoglycemia. Will check blood glucose regularly. 2. Do not stop abruptly because of rebound hypertension. If having a procedure that requires NPO diet, contact Dr. 3. Use with caution in pts with asthma. Can cause bronchospasm 4. Can cause bradycardia. Inform nurse if feeling SOB or dyspnea. Will monitor HR closely.
Prednisone	N/A	N/A	Anti-inflammatory or immunosuppressant agent in the treatment of many diseases	1. Do not administer a live vaccine because of the medication's immunosuppressive quality

				2. Can cause adrenal suppression on higher doses for prolonged periods. Labs need to be taken regularly. 3. Can cause HTN so monitor BP closely. 4. Significant drug interactions so consult drug databases before usage.
Ondansetron	N/A	N/A		Used for nausea and vomiting prevention and treatment	1. Can cause constipation. Inform pt to notify nurse if having abdominal pain. Monitor for BM and abdominal distention 2. Can cause headache, so notify nurse if headache occurs within minutes of administration. 3. Can cause QT prolongation on an EKG. Monitor HR closely before and after admin. 4. Watch for signs of Serotonin Syndrome when taken with other serotonergic agents. Monitor for mental status changes.
Levofloxacin	N/A	N/A		Primarily used as an antibiotic	1. BBW: Tendinitis or tendon rupture, peripheral neuropathy, and CNS effects. Monitor tendon pain, PV circulation, and CNS impairment. 2. If pt has myasthenia gravis, avoid giving because it may exacerbate muscle weakness in pt's with that condition. 3. Can cause a C. Diff. Infection.

				Monitor for foul smelling stool 4. Can cause hepatotoxicity so monitor labs for liver failure

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference:

UpToDate. (2025). Uptodate.com. https://www.uptodate.com/contents/vancomycin-drug-information?

       sectionName=Adult&anchor=F233334&source=auto_suggest&selectedTitle=1~1---2~4---

       vanco&showDrugLabel=true&search=vancomycin#F54416571