

## Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Vancomycin	125 – 500mg PO QID	N/A	Antibiotic used for Bacterial Infection	<ol style="list-style-type: none"> <li>1. Do not take this medication if you are pregnant, planning to become pregnant, or breast feeding because of fetal toxicity</li> <li>2. Report any signs of excessive bruising or bleeding. (The medication can cause thrombocytopenia)</li> <li>3. Report any signs and symptoms of abdominal pain and n/v/d, especially watery or loose stool as this medication can cause or prolong c-diff</li> <li>4. I am going to stay in the room with you after the medication is given for 5 minutes and check your vital signs because it may cause an allergic reaction or symptoms related closely to that (pruritis, flushing of the skin or a rash might appear, dyspnea, and hypotension)</li> </ol>
Metoprolol	25mg PO BID	N/A	Antihypertensive drug for high blood pressure / control blood pressure to a normal range	<ol style="list-style-type: none"> <li>1. Always check your blood pressure before administering this drug as you could cause severe hypotension</li> <li>2. Never stop taking this medication abruptly as it can cause rebound hypertension, tachycardia, and/or ischemia</li> <li>3. You must be very cautious taking this with asthma as this medication can cause bronchospasms</li> <li>4. Diabetic patients should always check your blood sugar before and during treatment taking this medication as it is known to mask the signs and symptoms of hypoglycemia (low blood</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

				sugar)
Prednisone	10-60mg PO Daily	N/A	Corticosteroid used for inflammation	<ol style="list-style-type: none"> <li>1. Always check your blood pressure before administering this drug as it will raise your blood pressure, putting you at risk of hypertension and/or a cardiac event</li> <li>2. Do not get a live vaccine while taking this medication as you have a weakened immune system and it can result in further sickness</li> <li>3. Do not take the medication if you are currently taking antifungal drugs</li> <li>4. Do not stop taking the medication abruptly as it can cause the condition to come back, finish the medication as prescribed by the provider</li> </ol>
Ondansetron	4-8mg PO Daily	N/A	Antiemetic used for nausea and vomiting	<ol style="list-style-type: none"> <li>1. While taking this drug we want to up your fluid intake as well as fresh fruits and fiber to prevent constipation</li> <li>2. Report any signs or symptoms of irregular heartbeat and chest pain as the medication can cause prolonged QT intervals</li> <li>3. Report if you start to develop headache or headaches or head pain as this drug is known to cause this, also another reason to increase fluid intake</li> <li>4. I am going to stay in the room with you after the medication is given for 5 minutes and check your vital signs because it may cause an allergic reaction or symptoms related closely to that (hypotension, angioedema, dyspnea, and urticaria)</li> </ol>
Levofloxacin	250mg	N/A	Antibiotic used for Bacterial Infection	<ol style="list-style-type: none"> <li>1. Report any leg pain, pain near a joint, or muscle weakness as this medication is known</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

				<p>to cause tendonitis and tendon rupture</p> <p>2. Report any signs or symptoms of irregular heartbeat and chest pain as the medication can cause prolonged QT intervals</p> <p>3. Report any signs and symptoms of abdominal pain and n/v/d, especially watery or loose stool as this medication can cause or prolong c-diff</p> <p>4. Report any changes in your vision especially if light in the room or outside becomes intolerable as this medication can cause photosensitivity</p>
				<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>
				<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>
				<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>
				<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>

## Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

				1. 2. 3. 4.
--	--	--	--	----------------------

Allergies: None

Primary IV fluid and rate: N/A

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic): N/A

Medication reference: <https://www.uptodate.com/contents/search>

<https://www.drugs.com/>