

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Eliquis	5 mg/ 1 tab One daily		Direct acting by mouth anti-coagulant	<ul style="list-style-type: none"> • Apixaban is used to treat certain types of blood clots • Prevention of deep vein thrombosis (DVT) • Treatment of and reduction in risk of recurrence of Deep vein thrombosis or pulmonary embolism
Levothyroxine	100 mcg tablet		Mimic natural process of thyroid hormone production	<ul style="list-style-type: none"> • Swallow the capsule whole. Do not cut, chew, or crush it. • Hypothyroidism can sometimes cause infertility in men and women • Blood tests will be needed to check for unwanted effects.
Atorvastatin	10 mg tablet		Inhibit HMG-CoA reductase, controls cholesterol production in the liver	<ul style="list-style-type: none"> • Pt will need to have regular medical examinations to monitor the effectiveness of this drug • Avoid grapefruit juice while you are taking this drug. • Avoid driving or performing hazardous or delicate tasks that require concentration because of headache, blurred vision, nervousness.
Pantoprazole	40mg/10mL IVP Daily	IVP – 40mg /10mL NS 10ml over 2-4 minutes	This medication lowers stomach acid and treats peptic ulcers, as well as aiding in Gastroesophageal Reflux Disease.	<ul style="list-style-type: none"> • Monitor stool for diarrhea because of possibility of Clostridium difficile infection • Monitor for bleeding because of risk of low platelet level by ordering PLT labs • Report muscle cramps and tremors, may require magnesium supplementation • Report shortness of breath, chest pain, severe cough to monitor pneumonia
Nifedipine/	30 mg x1 tablet		Calcium channel blocker, results in dilation, reduced blood pressure,	<ul style="list-style-type: none"> • Used to treat hypertension (high blood pressure) and angina (chest pain).

Adult/Geriatric Medication Worksheet - Current Medications & PRN from 0700-1400 current day

			decreased peripheral vascular resistance	<ul style="list-style-type: none">• Nifedipine may also be used for purposes not listed in this medication guide.• You should not use nifedipine if you have severe coronary artery disease, or if you have had a heart attack.

Allergies: NKDA

Primary IV fluid and rate:

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference: UpToDate