

Student Name:   Rafaella Colmenero  

Unit:   Oceans  

Pt. Initials:   F.V.  

Date:   9/2/25  

**Medication Worksheet – Current Medications & PRN for Last 24 Hours**

Allergies:   NKDA  

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
	Isotonic/ Hypotonic/ Hypertonic			

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP – List solution to dilute and rate to push. IVPB – List mL/hr and time to give	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Oxcarbazepine	Anticonvulsant	Mood changes	300mg PO BID	Y N		Hyponatremia	<ol style="list-style-type: none"> <li>1. Dizziness and/or ataxia may occur—call before getting up to avoid falling</li> <li>2. Electrolytes will be monitored especially sodium—can cause hyponatremia</li> <li>3. Notify if you experience any leg cramping, cold/clammy skin, headache, N/V/D—signs and symptoms of hyponatremia</li> <li>4. Monitor for signs of seizures (changes in LOC, foaming at the mouth, etc.)—can increase risk of new onset of primary generalized seizures.</li> </ol>
Bupropion	Norepinephrine dopamine reuptake inhibitors	Severe depression	100mg PO daily	Y N		Seizures, hepatic impairment	<ol style="list-style-type: none"> <li>1. May take up to 4 weeks to reach full therapeutic effect—do not stop abruptly</li> <li>2. may cause decreased alertness and motor skills—avoid tasks that require increased alertness until response to drug is established</li> <li>3. do not take more than prescribed—more than 150mg/day increases risk of seizures and hepatic impairment</li> <li>4. Ask patient if they have taken an MAOI as Bupropion should not be started until 14 days after stopping the MAOI and vice versa.</li> </ol>

Student Name: \_Rafaella Colmenero\_\_\_\_\_

Unit: \_Oceans\_

Pt. Initials: F.V.\_\_\_\_

Date: \_9/2/25\_\_

**Medication Worksheet – Current Medications & PRN for Last 24 Hours**

Allergies: \_NKDA\_\_\_\_\_

Quetiapine Fumarate	Antipsychotic	Severe psychosis	25mg PO at bedtime	Y N		EPS, NMS	<ol style="list-style-type: none"> <li>1. Drink fluids often and continue regular physical activity—risk for constipation</li> <li>2. Drowsiness, decreased alertness, decreased motor skills and hypotensive effect can occur—change positions slowly or call for help before getting up</li> <li>3. Report any stiffness in the neck or any other changes to muscle and/or joints – increased risk for EPS</li> <li>4. avoid exposure to extreme heat – increases heat intolerance which interferes with ability to regulate temperature</li> </ol>
Hydroxyzine Pamoate	Antihistamine	Anxiety	50mg PO Q6 PRN	Y N		Anticholinergic effects	<ol style="list-style-type: none"> <li>1. CNS depression—call before getting up and use caution when driving</li> <li>2. Obtain EKG before starting med—contraindicated in patients with prolonged QT interval</li> <li>3. Can use sugar free candy/gum for dry mouth</li> <li>4. Do not consume with alcohol—can increase CNS depression</li> </ol>
				Y N			<ol style="list-style-type: none"> <li>1.</li> <li>2.</li> <li>3.</li> <li>4.</li> </ol>