

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
Click here to enter text.	Isotonic <input type="checkbox"/> Hypotonic <input type="checkbox"/> Hypertonic <input type="checkbox"/>	Click here to enter text.	Click here to enter text.	Click here to enter text.

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
Hydroxyzine pamoate	Antihistamine	Anxiety	25mg PO PRN q6	Yes	N/A	Long term = risk of tardive dyskinesia	1. Drowsiness - caution with driving 2. Potentiating action w/ CNS depressants (barbiturates, narcotics, non-narcotic analgesics) 3. Use sugar-free liquids and candies for dry mouth 4. QT Prolongation/Torsade de Pointes
Lorazepam	Benzodiazepine	Alcohol withdrawal / anxiety	1-2 mg PO PRN q4 hours	Yes Choose an item.	N/A	Abrupt or too-rapid withdrawal may result in pronounced restlessness, irritability, insomnia, hand tremor, abdominal cramping, muscle cramps, diaphoresis,	1. Drowsiness usually disappears during continued therapy 2. Avoid tasks that require alertness, motor skills until response to drug is established 3. Do not use alcohol, CNS depressants 4. Smoking reduces drug effectiveness (Contraception / Notify if pregnant)

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						vomiting	
Haldol	Antipsychotic	for psychotic symptoms/agitation	10 mg IM PRN q8	Yes Choose an item.	N/A	BLACK BOX WARNING: Increased risk of mortality in elderly pts with dementia-related psychosis with use of injections	<ol style="list-style-type: none"> 1. Inject slow, deep IM into upper outer quadrant of gluteus maximus (Max 3mL per site) 2. Should remain lying down 30-60 min to minimize hypotensive effect after admin 3. Anticholinergic effects 4. Tachycardia / Prolonged Qt intervals
Diphenhydramine	Antihistamine	management of cholinergic effects of antipsychotic	50 mg IM q8 PRN psychotic symptoms	Yes	N/A	Injection site reaction, Sedation, apnea, hypotension	<ol style="list-style-type: none"> 1. Avoid tasks that require alertness, motor skills until response to drug is established 2. Dry mouth, drowsiness, dizziness may occur - Use sugar-free liquids and candies for dry mouth 3. Slow movements - increased risk of orthostatic hypotension 4. Monitor BP
Acetaminophen	Analgesic/antipyretic	Mild-moderate pain	500 mg PO q6 for pain of 1-	Yes	N/A	EARLY SIGNS OF TOXICITY: Anorexia,	<ol style="list-style-type: none"> 1. Maximum 4000 mg per day 2. Do not take for longer than 10 days

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
			4 on pain scale			nausea, diaphoresis, fatigue in first 12-24 hrs	3. Assess baseline temperatue and/or pain type & location 4. Caution taking with preexisting liver disorder
Olanzapine	Antipsychotic	Agitation/psychotic behavior	5 mg PO q6 for agitation	Yes	N/A	Rare reactions include seizures, neuroleptic malignant syndrome	1. Drink plenty of h2o, use sugar-free liquids and candies for dry mouth 2. Assess for tremors, changes in gait, abnormal muscular movements 3. Monitor diet, exercise program to prevent weight gain 4. Monitor A1C & liver function
Ibuprofen	NSAID	Mild-moderate pain	400 mg PO q6 for pain of 1-4 on pain scale	Yes Choose an item.	N/A	BLACK BOX WARNING: Increased risk of severe GI reactions, including ulceration, bleeding, perforation. Increased risk of new onset or worsening of preexisting	1. N/V, dizziness, rash - take with food or milk for stomach upset 2. Avoid aspirin & ETOH with drug (increased risk of GI bleed) 3. Assess onset of pain - type & location 4. Assess renal function

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.	Date: Click here to enter a date.	Allergies: Click here to enter text.		
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						hypertension	
Risperdal	Antipsychotic	Psychotic behavior/bipolar mania	1 mg tablet PO BID	Yes	N/A	BLACK BOX WARNING: Increased risk of mortality in elderly pts with dementia-related psychosis with use of injections	<ol style="list-style-type: none"> 1. Agitation, anxiety, sedation, headache 2. Assess liver/renal function **A1C 3. Adherence to med regimen 4. Encourage use of sugar-free liquids and candies for dry mouth
Nicotine patch	Stimulant	Nicotine replacement	21 mg transdermal patch daily	Yes Choose an item.	N/A	OD: palpitations, tachyarrhythmia, seizures, hypotension, diaphoresis	<ol style="list-style-type: none"> 1. DO NOT CUT PATCH 2. Replace daily; rotate sites; do not use same site within 7 days; do not use same patch longer than 24 hrs 3. Wash hands with water alone after applying patch 4. Do not smoke while wearing patches
Ziprasidone	Antipsychotic	Psychotic behavior/bipolar disorder/maniac	40 mg PO BID	Yes	N/A	BLACK BOX WARNING: Increased risk of mortality in elderly pts with dementia-	<ol style="list-style-type: none"> 1. Agitation, anxiety, sedation, headache 2. Assess liver/renal function **A1C 3. Drink plenty of H₂O, use sugar-free liquids and candies for dry mouth 4. Monitor diet, exercise program to

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
						related psychosis, mainly due to pneumonia, heart failure	prevent weight gain
Prazosin	Alpha-blocker	Nightmares	1 mg PO QHS (everynight at bedtime)	Yes Choose an item.	N/A	upper stomach pain, loss of appetite, dark urine, clay-colored stools, jaundice / priapism (prolonged erection)	<ol style="list-style-type: none"> 1. Dizziness. Headache, fatigue 2. Hypotensive effects - slow movements 3. Do not administer if BP is low 4. Adhere to med regimen
Nicotine inhaler	Stimulant	Nicotine replacement	10 mg cartridge for inhalation Q2	Yes	N/A	OD: palpitations, tachyarrhythmia , seizures, hypotension, diaphoresis	<ol style="list-style-type: none"> 1. Insert into mouthpiece - Puff on nicotine cartridge mouthpiece for 20 min 2. Puff on nicotine cartridge mouthpiece for about 20 min as instructed 3. May cause nausea, hiccups 4. Reduced withdrawal symptoms
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item.	Click here to enter text.	Click here to enter text.	<ol style="list-style-type: none"> 1. Click here to enter text. 2. Click here to enter text.

Adult/Geriatric Medication Worksheet - Current Medications & PRN for Last 24 Hours

Student Name: Synthia Gomez		Unit: Click here to enter text.	Patient Initials: Click here to enter text.		Date: Click here to enter a date.	Allergies: Click here to enter text.	
Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Correct Dose? If not, what is correct dose?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Click here to enter text.			3. Click here to enter text. 4. Click here to enter text.
Click here to enter text.	Click here to enter text.	Click here to enter text.	Click here to enter text.	Choose an item. Click here to enter text.	Click here to enter text.	Click here to enter text.	1. Click here to enter text. 2. Click here to enter text. 3. Click here to enter text. 4. Click here to enter text.