

Question #

Student Name:

Kaitlyn Diaz

Based on the "Topic" and "Subtopic," I missed a question about:

Intrapartum Care



I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content

List five or more bullet points with your "take-aways" from this packet.

(What is most important for you to remember as you prepare for the NCLEX and future patient care?)

- Intrapartum Care is for early identification of fetal heart rate abnormalities and prevention of maternal complications.

- A woman with preeclampsia with severe features is maintained on bed rest with the side rails up.

- With mothers who have preeclampsia emergency drugs, oxygen and suction equipment should be available and at bedside.

- Important signs and symptoms to look for with preeclampsia with severe features: nausea in upper right quadrant, blurred vision, headaches that are unresponsive with meds., htn, proteinuria, seizure activity

- Pulmonary complications that occur with severe preeclampsia is dyspnea, pulmonary edema, hypoxemia.

Question #

Based on the "Topic" and "Subtopic," I missed a question about:

Assessment



I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content

List five or more bullet points with your "take-aways" from this packet.

(What is most important for you to remember as you prepare for the NCLEX and future patient care?)

- Accurate measurements of patient's blood pressure is essential in early detection for mother's with hypertensive disorders.

- BP assessments should focus on trends over time, rather than on a single measurement

- Dependent edema is edema of the lowest or most dependent parts of the body, where hydrostatic pressure is greatest.

- Proteinuria is defined as a concentration at or greater than 300 mg or a protein/creatinine ratio greater than 0.3.

- 24-hour collection measures the quantity of protein and creatinine clearance is more reflective of true renal status, it is preferred over dipstick testing, which should not be used for diagnosis of preeclampsia.