

Question #	Student Name: Kaitlyn Diaz
Based on the "Topic" and "Subtopic," I missed a question about: Prolapsed Umbilical Cord	
<input type="checkbox"/>	I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content
<p>List five or more bullet points with your "take-aways" from this packet. (What is most important for you to remember as you prepare for the NCLEX and future patient care?)</p> <ul style="list-style-type: none"> • A prolapse of the umbilical cord occurs when the cord lies below the fetus. • The four positions is A. occult prolapse of cord. B. complete prolapse of cord. C. cord presenting in front of the fetal head may be seen in the vagina. D. frank breech presentation with prolapsed cord. • Occlusion of blood flow from a prolapsed umbilical cord that last more than 5 minutes will result in CNS damage or death of fetus. • A vaginal birth can still be possible as long as the cervix is fully dilated and forceps or vacuum may be used for assistance. • A cesarean birth is most likely to be performed with a prolapsed umbilical cord. 	

Question #	
Based on the "Topic" and "Subtopic," I missed a question about: Therapeutic Management	
<input type="checkbox"/>	I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content
<p>List five or more bullet points with your "take-aways" from this packet. (What is most important for you to remember as you prepare for the NCLEX and future patient care?)</p> <ul style="list-style-type: none"> • The three positions that can be used to aid in delivery are a. knee-chest position b. Trendelenburg position c. Hips elevated with pillows, with side-lying position maintained. • The nurse must maintain vaginal elevation to minimize cord compression during transport. • During transportation avoid or minimize manual palpation or handling of the cord to minimize cord vessel vasospasm. • Ultrasound examination may be used to confirm presence of fetal heart activity before cesarean delivery. • While preparing for surgery, give the woman oxygen 8 to 10 (L/min) 	