

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Generic Name	Dosage with route and schedule	IVP-List diluent solution, volume, and rate of administration IVPB- List concentration and rate of administration	Patient specific therapeutic reasoning	Patient specific teaching with reasoning
Vancomycin	125mg PO QID 10 days	5mg/ml IVP over at least one minute	Treat infection of the intestines caused by Clostridium difficile, which can cause watery or bloody diarrhea.	<ol style="list-style-type: none"> 1. Tell doctor if you have 2. hearing problems. 3. Don't take more than prescribed, it may cause serious or life threatening side effects. 4. prescribed, it may cause
Metoprolol	50mg PO BID Immediate release 100mg PO BID extended release	1mg/ml IVP 1-2 minutes	Used to treat angina (chest pain) & hypertension (high blood pressure).	<ol style="list-style-type: none"> 1. Do not use two doses 2. at one time. 3. Avoid driving or hazardous activity. 4. until you know how med will affect you.
Prednisone	1mg-50mg PO daily		Treat allergic disorders, skin conditions, ulcerative colitis, & arthritis.	<ol style="list-style-type: none"> 1. Med can weaken your immune system. 2. Do not stop taking medication suddenly. 3. Do not stop taking 4. medication suddenly.
Ondansetron	8mg PO BID	2mg/ml IVP 2-5 minutes	Prevent nausea & vomiting	<ol style="list-style-type: none"> 1. Don't use if you have 2. prolonged QT syndrome, 3. can be fatal. 4. Don't take med if you're taking apomorphine.
Levofloxacin	750mg PO daily	5mg/ml IVPB	Treat bacterial infections	<ol style="list-style-type: none"> 1. Take liquid medication 2. on an empty stomach. 3. Monitor for tendon rupture. 4. Monitor for tendon

Look on bottom sheet for other teaching

Adult/Geriatric Medication Worksheet – Current Medications & PRN from 0700-1400 current day

Vancomycin				<ol style="list-style-type: none"> 1. Use medication for full 2. prescribed time, even if 3. symptoms quickly improve. 4. Give med slowly, can cause flushing syndrome.
Metoprolol				<ol style="list-style-type: none"> 1. Rebound HTN if 2. stopped abruptly. 3. Take with meal to 4. avoid GI upset.
Prednisone				<ol style="list-style-type: none"> 1. long time use can lead 2. to bone loss. 3. Do not use "live" vaccine 4. while using this medication.
Ondansetron				<ol style="list-style-type: none"> 1. Take with a full glass 2. of water. 3. May impair your reaction, 4. so avoid driving.
Levofloxacin				<ol style="list-style-type: none"> 1. Drink extra liquid to 2. keep kidneys working 3. properly. 4. skipping doses can increase risk of infection.

Allergies:

Primary IV fluid and rate:

Patient specific reasoning for IV fluids (including type isotonic, hypotonic, hypertonic):

Medication reference: