

Question #	Student Name: <u>Camryn Cowley</u>
Based on the "Topic" and "Subtopic," I missed a question about: <u>Newborn - Hyperbilirubinemia</u>	
<input checked="" type="checkbox"/>	I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content
<p>List five or more bullet points with your "take-aways" from this packet. (What is most important for you to remember as you prepare for the NCLEX and future patient care?)</p> <ul style="list-style-type: none"> • Most newborns have some jaundice after 24 hours • Assess jaundice every 8-12 hours - bony areas, conjunctiva, buccal mucosa • TcB is useful but levels > 15 or first 24 hours need serum check • Phototherapy helps breakdown bilirubin • Severe cases may require exchange transfusion 	

Question #	
Based on the "Topic" and "Subtopic," I missed a question about: <u>Ante-Partum - Adolescent Pregnancy needs</u>	
<input checked="" type="checkbox"/>	I have reviewed each of the excerpts/activities listed under the Packet > Remediation Content
<p>List five or more bullet points with your "take-aways" from this packet. (What is most important for you to remember as you prepare for the NCLEX and future patient care?)</p> <ul style="list-style-type: none"> • Teens are at an ↑ risk for being deficient in Calcium and Iron • Cephalopelvic disproportion is common in younger women due to growth of the pelvis is delayed • Recommended weight gain goals are no different • The adolescent needs careful teaching regarding nutritional intake and physical activity in the post partum period • Adolescents are less likely to seek pre-natal care - so promoting access to it is essential 	