

## Invega Sustenna® Review

1. You are preparing to administer Invega Sustenna® to a patient who weighs 146 pounds. What size needle will you use?

**1 Inch, 23 gauge**

2. Name the most common side effects a patient may experience when taking Invega Sustenna®.

**Injection site reactions, sleepiness, drowsiness, dizziness, feeling restless or needing to be constantly moving, abnormal muscle movements, shuffling, uncontrolled involuntary movements, abnormal movements of eyes**

3. What black box warning is associated with this medication?

**Increased mortality in elderly patients with dementia related psychosis**

4. What are the standard *recommended* Day 1 and Day 8 initiation dosages for a person with schizophrenia?

**Day 1 (234 mg) Day 8 (156 mg)**

5. What is the standard *recommended* maintenance dosage for a patient with schizophrenia?

**39mg - 234mg monthly**

6. What is the correct location for administration of the Day 1 and Day 8 initiation doses?

**Deltoid**

7. What are the *recommended* Day 1 and Day 8 initiation dosages for a person with a creatinine clearance of 65mL/min?

**Day 1 (156 mg) Day 8 (117 mg)**

Revised 09/2023

8. What would the *recommended* maintenance dosage be for a patient with mild renal impairment?

**78 mg monthly**

9. What is the rationale for a trial of oral paliperidone or oral or injectable risperidone before placing a patient on Invega Sustenna®?

**To establish tolerability to adverse effects**

10. Name three practices when preparing an intramuscular injection that ensure aseptic technique is maintained throughout the procedure.

**Clean medication vial's rubber stopper with alcohol before piercing with needle, clean patients skin with alcohol, keep needle covered and do not touch the uncovered needle with anything except when going into the patients skin**

11. Explain the rationale for obtaining consent for this psychoactive medication prior to preparing the medication for administration.

**It ensures patient understands the risk and benefits. It also allows the patient the opportunity to make decisions and ask questions before getting the medication**