

Pediatric Medication Worksheet - Current Medications & PRN for Last 24 Hours **Allergies: NKDA**

Primary IV Fluid and Infusion Rate (ml/hr)	Circle IVF Type	Rationale for IVF	Lab Values to Assess Related to IVF	Contraindications/Complications
0.9%NS 50ml/hr	ISOTONIC Isotonic/ Hypotonic/ Hypertonic	TO HELP CORRECT HYPERCALCEMIA BY INCREASING THE AMOUNT OF CALCIUM EXCRETED IN URINE	TOTAL SERUM CALCIUM, IONIZED FREE CALCIUM, AND RENAL FUNCTION SUCH AS CREATININE AND BUN	Contraindicated in pts. with uncontrolled HF, RF, and Pulmonary Edema as it can cause risk for fluid overload

Generic Name	Pharmacologic Classification	Therapeutic Reason	Dose, Route & Schedule	Therapeutic Range?	IVP - List diluent solution, volume, and rate of administration IVPB - List concentration and rate of administration	Adverse Effects	Appropriate Nursing Assessment, Teaching, Interventions (Precautions/Contraindications, Etc.)
				Is med in therapeutic range?			
				If not, why?			
CANNABIDIOL	PHYTOCANNABINOID	USED AS ANTICONVULSANT	100mg/mL liquid 200mg 2mL via G BUTTON BID	yes	N/A	FATIGUE, DIARRHEA, DRY MOUTH, REDUCED BLOOD PRESSURE	<ol style="list-style-type: none"> 1. MEDICATION CAN INERACT WITH OTHER MEDS SUCH AS BLOOD THINNERS, WHEN MIXED WITH THESE MEDS IT CAN CAUSE LIVER TOXICITY. WE WILL MONITOR LIVER ENZYME LEVELS AND REPORT TO PROVIDER IF OUT OF RANGE 2. AVOID ACTIVITIES THAT REQUIRE ALERTNESS OR OPERATING HEAVY MACHINES AS THIS MEDICATION CAN ALTER MENTAL AND PHYSICAL STATES. 3. PROPER MEDICATION STORAGE AND HANDLING AS MEDICATION CAN CAUSE MENTAL IMPAIRMENT, ALWAYS HANDLE MEDICATION WITH GLOVES OR OTHER PROTECTIVE BARRIER TO AVOID ACCIDENTAL SELF OR OVER MEDICATION. 4. THIS TYPE OF MEDICATION IS NOT TIGHTLY REGULATED AND COULD CONTAIN CONTAMINANTS SUCH AS

							PESTICIDES. RECOMMENDED TO ACQUIRE MEDICATION FROM REPUTABLE SOURCE TO AVOID CONTAMINANTS ISSUE.
CETIRIZINE	2 ND GEN ANTIHISTAMINE	ANTI-ALLERGIC AGENT TO TREAT SNEEZING, ITCHY EYES, AND HIVES.	1mg/mL liquid 5mg 5mL PO ONCE DAILY	YES	N/A	DROWSINESS, FATIGUE, HEADACHE, DRY MOUTH	<p>1. THOUGH IT SHOULD BE NON-DROWSY, IN CERTAIN PATIENTS IT CAN CAUSE FATIGUE/DROWSINESS, AVOID HEAVY MACHINERY OR ACTIVITY THAT REQUIRES ALERTNESS AND WAKFULNESS UNTIL YOU KNOW HOW THE MED WILL AFFECT YOU.</p> <p>2. AVOID ALCOHOL AND ANY OTHER CNS DEPRESSANTS AS IF THEY ARE MIXED WITH THIS MED, IT CAN CAUSE DANGEROUSLY SLOW BREATHING RATES</p> <p>3. DRY MOUTH IS A COMMON SIDE EFFECT, IF IT BECOMES UNBEARABLE WE CAN MANAGE DRY MOUTH WITH GOOD ORAL HYGIENE, MOUTH RINSING, AND ALSO CHEWING SUGAR FREE GUM</p> <p>4. IF MEDICATION IS USED FOR LONGER THAN 3 MONTHS, IT COMES WITH A RISK FOR WITHDRAWAL ESPECIALLY SEVERE INTENSE ITCHING. REPORT IMMEDIATELY TO PROVIDER.</p>
GLYCOPHYRROLATE	ANTICHOLINERGIC AGENT	ACETYLCHOLINE NEUROTRANSMITTER BLOCKER THAT REDUCES STOMACH ACID, SALIVA, AND SWEAT PRODUCTION	1mg tablet PO Q 8 HRS	YES	N/A	SEVERE DRY MOUTH, CONSTIPATION, URINARY RETENTION, BLURRED VISION	<p>1. MEDCATION CAUSES INTENSE DRY MOUTH, USE GOOD ORAL HYGIENE, FREQUENT MOUTH RINSES, AND DRINK PLENTY OF WATER TO HELP WITH DRY MOUTH.</p> <p>2. AVOID EXTREME HEAT OR HEAVY EXERCISE AS THIS MEDICATION CAN REDUCE SWEATING SIGNIFICANTLY. EXTREME HEAT AND EXERCISE CAN CAUSE RISK FOR OVERHEATING AND HEAT STROKE, STAY HYDRATED AND</p>

							<p>AVOID HOT WEATHER.</p> <p>3. AVOID ACTIVITY THAT REQUIRES CLEAR VISION AND A CLEAR HEAD AS THIS MEDICATION CAN CAUSE BLURRY VISION AND DIZZINESS.</p> <p>4. PLEASE REPORT IMMEDIATELY TO PROVIDER ANY SUDDEN ABNORMAL URINE CHANGES SUCH AS LESS URINE OR GOING LESS FREQUENTLY IN THE DAY AS THESE COULD BE SIGNS OF URINARY RETENTION.</p>
LACOSAMIDE	ANTICONVULSANT AND ANTIPILEPTIC DRUG	TO TREAT PARTIAL-ONSET AND PRIMARY GENERALIZED TONIC-CLONIC SEIZURES	50mg tablet via NG button Q 8 HRS	YES	N/A	<p>DIZZINESS, HEADACHE, NAUSEA, FATIGUE, AND VISION CHANGES</p>	<p>1. DO NOT STOP MEDICATION ABRUPTLY EVEN IF PT FEELS WELL AS SUDDEN CESSATION CAN CAUSE SEIZURES.</p> <p>2. AVOID HEAVY MACHINERY AND ACTIVITY THAT REQUIRES ALERTNESS OR CLEAR VISION AS MEDICATION CAN CAUSE DIZZINESS AND VISION CHANGES THAT ARE NORMAL</p> <p>3. PLEASE REPORT IMMEDIATELY TO PROVIDER ANY SUDDEN OR WORSENING DEPRESSION OR SUICIDAL THOUGHTS, OR ANY UNUSUAL BEHAVIORS.</p> <p>4. TAKE MEDICATION AS DIRECTED BY PROVIDER, WITH FOOD IF HAVING STOMACH UPSET AND DO NOT DOUBLE DOSE IF A DOSE IS MISSED.</p>
OXCARBAZEPINE	ANTICONVULSANT	TO TREAT PARTIAL-ONSET SEIZURES	225mg tablet via G BUTTON Q 8 HRS	YES	YES	<p>DOUBLE VISION, COORDINATION ISSUES, NAUSEA, DIZZINESS</p>	<p>1. DO NOT STOP ABRUPTLY AS IF STOPPED WITHOUT A PROVIDERS DIRECTIONS CAN CAUSE RISK FOR MORE SEIZURES</p> <p>2. MONITOR AND REPORT ANY SUDDEN SKIN CHANGES SUCH AS RASH, FEVER, BLISTERS, AND EVEN SWOLLEN LYMPH NODES AS IT COULD BE SIGNS OF SEVERE-FATAL</p>

								SKIN REACTIONS. 4. PLEASE REPORT NAUSEA, SEVERE HEADACHE, CONFUSION, FATIGUE, OR INCREASED SEIZURES IMMEDIATELY TO PROVIDER AS THESE CAN SYMPTOMS OF LOW SODIUM LEVELS WHEN TAKING THIS MEDICATION.
--	--	--	--	--	--	--	--	--