

Marisol Fick, Nadia Miracle, and Macy Schafer

**EBP Phase 1**

-

**H**ealth care organizations strive to minimize harm and provide safe environments for patients they serve. Despite a vast array of prevention strategies, hospital-acquired pressure injuries (HAPIs) continue to occur, especially in critical care. Rates of such injuries have been reported to range from 2.8% to 53.4% in critical care units, compared with 2.0% to 8.3% in medical-surgical units.<sup>1</sup>

Regulatory and quality organizations consider HAPI rates to be a measure of the quality of nursing care provided and assess financial penalties when they occur. However, organizations of experts such as the Centers for Medicare and Medicaid Services, the National Pressure Ulcer Advisory Panel (NPUAP), and the Wound, Ostomy, and Continence Nurses Society have acknowledged that some HAPIs may be unavoidable.<sup>2,4</sup> The NPUAP defined "unavoidable" pressure injuries as those that develop even when the provider (1) evaluated the individual's clinical condition and pressure injury risk factors; (2) defined and implemented interventions that were consistent with individual needs, goals, and recognized standards of practice; (3) monitored and evaluated the impact of the interventions; and (4) revised the approaches as appropriate.<sup>3</sup>

Risk factors have been associated with the development of HAPIs, yet those that best predict the development of HAPIs are not completely understood.<sup>1,5</sup> Nonmodifiable risk factors such as age<sup>3</sup> and history of pressure injuries<sup>6</sup> may tip the scale toward pressure injury development despite the best preventive interventions. Unavoidable pressure injuries may occur when the magnitude and severity of the risk factors are extremely high and preventive measures are either contraindicated or inadequate given the risk.<sup>7</sup> Although new technology is available to provide quantitative assessment of turning and patient mobility, currently this technology is not widespread or integrated into most acute care settings.

#### About the Authors

**Joyce Pittman** was a nurse practitioner and coordinator, Wound/Ostomy Department, Indiana University Health Academic Health Center, Indianapolis, Indiana, and is currently an associate professor, University of South Alabama, Mobile, Alabama. **Terrie Beeson** is a clinical nurse specialist for surgical intensive care, and **Jill Dillon** is a clinical nurse specialist for surgical trauma intensive care, Indiana University Health Academic Health Center. **Ziyi Yang** is a biostatistician, Indiana University School of Nursing, Indianapolis. **Janet Cuddigan** is a professor, College of Nursing, University of Nebraska, Omaha, Nebraska.

**Corresponding author:** Joyce Pittman, PhD, ANP-BC, FNP-BC, CNRN, Associate Professor, College of Nursing, University of South Alabama, 5721 USA Dr N, Mobile AL 36688 (email: joyce.pittman@comcast.net).

Until recently, research on unavoidable HAPIs has been limited because of the lack of valid and reliable tools for measuring and evaluating the provision of appropriate preventive interventions. The Pressure Ulcer Prevention Inventory (PUPI)<sup>7</sup> was designed using the NPUAP definition of unavoidable pressure injury<sup>3</sup> and Braden and Bergstrom's conceptual model of the etiology of pressure injuries.<sup>8</sup> The PUPI operationalized the 4 key concepts that are thought to capture the construct of unavoidable HAPIs and demonstrated acceptable validity (content validity index = 0.99) and reliability ( $\kappa = 1.0$ ,  $P = .02$ ; rater agreement 93%).<sup>1,2,7</sup>

The aims of this study were (1) to determine the proportion of HAPIs among patients in critical and progressive care units that are unavoidable, and (2) to identify the risk factors (or characteristics) of patients in critical and progressive care units that differentiate avoidable from unavoidable HAPIs.


#### Methods

##### Design

This study used a descriptive, retrospective, comparative design to examine rates of avoidable and unavoidable HAPIs in adult critical and progressive care patients in 6 acute care hospitals within a large academic health care system in the midwestern United States. The critical care areas included surgical, trauma, cardiovascular surgical, cardiac, neurologic, and medical intensive care and corresponding progressive care units. The study was approved by the Indiana University institutional review board before the start of data collection.

An investigator-developed conceptual framework for differentiating unavoidable from avoidable pressure ulcers (Figure 1) was used to guide this study. The aspects of this model that are new and unique incorporate (1) new epidemiological evidence on pressure injury risk factors,<sup>9,10</sup> (2) risk-based prevention strategies consistent with the 2014 Pressure Ulcer International Guideline,<sup>11</sup> and (3) guidance for determining whether the pressure injury was avoidable or

Despite prevention strategies, hospital-acquired pressure injuries continue to occur, especially in critical care.



**Figure 1** Conceptual framework for differentiating unavoidable from avoidable pressure ulcers.

Reproduced with permission from Janet Cuddigan. ©2014 Janet Cuddigan, PhD, RN, CWCN, FAAN.  
<sup>a</sup> Risk factors measured with Braden Scale subscales.

unavoidable based on the implementation of appropriate risk-based interventions.<sup>7</sup>

#### Sample/Setting

The sample consisted of patients who (1) had a HAPI develop while in critical and progressive care (NPUAP stage 2, 3, or 4, unstageable, or deep tissue pressure injury,<sup>12</sup> including pressure injuries from medical devices and on mucous membranes), (2) were hospitalized between 2012 and 2015, and (3) were aged 18 years or older. Patients were excluded if the HAPI developed outside the critical or progressive care unit. We used the National Database of Nursing Quality Indicators methods for identifying unit-acquired pressure injuries.<sup>13</sup> On the basis of a previous pilot study,<sup>7</sup> we determined potential sample size by estimating the number of HAPIs occurring during a 12-month period and the proportion of those estimated as unavoidable. Approximately 160 patients with a HAPI were estimated to provide an adequate sample to detect meaningful information.

#### Identification of Unavoidable HAPIs

To achieve the first study aim, eligible patients were identified by using monthly pressure injury prevalence surveys, medical record reporting mechanisms, medical coding procedures, and/or quality reporting processes. Patients were divided into 2 groups (avoidable and unavoidable) using the PUPI tool.

The PUPI contains 13 items (Figure 2). If all items in the PUPI are answered "yes" (all interventions were appropriately performed and documented), the HAPI was identified as unavoidable.<sup>7</sup>

The Braden Inventory Worksheet, an investigator-developed data collection tool specific to our electronic medical record (EMR) documentation, was used to collect Braden Scale subscale and total scores<sup>14</sup> for the 3 days before the first documentation of the HAPI and the interventions documented that correspond to each Braden Scale subscale (Figure 3). These data were then used to complete the PUPI.

#### Identification of Risk Factors Differentiating Avoidable From Unavoidable HAPIs

Data were collected from the EMR upon admission and 3 days before first documentation of the HAPI, including (1) demographic and clinical information, (2) Braden Scale subscale and total scores, (3) additional risk factors identified as significant in epidemiological studies (eg, poor perfusion, body temperature), and (4) pressure injury prevention interventions. The types and number of comorbidities were determined. For a variable to be marked yes, it had to be documented on any of the 4 days. For example, if "chemically sedated (IV drip)" was marked no on the HAPI date, yes on 1 day before, yes on 2 days before, and yes on 3 days before, then it was counted as yes (Figure 4).

Subject ID: \_\_\_\_\_ **Pressure Ulcer Prevention Inventory** HAPU Acquisition Date: \_\_\_\_/\_\_\_\_/\_\_\_\_

HAPU Location: 1= Sacrum/Coccyx 2= Ischium 3= Trochanter 4= Heel 5= Occipital 6= Ear 7= Other \_\_\_\_\_

HAPU Laterality: 1= Right 2= Left 3= Midline \_\_\_\_\_

HAPU Stage: 1= sDTI 2= 2 3= 3 4= 4 5= Unstageable 6= Indeterminate \_\_\_\_\_

Audit Date: \_\_\_\_\_ Auditor: \_\_\_\_\_

**Review medical record 3 days prior to the documented development of the HAPU**  
Assign the appropriate score for each item:  
1= NO, not appropriate 2= YES, appropriate

1. Clinical Condition Evaluation	SCORE
History and physical completed upon admission	
Braden pressure ulcer assessment upon admission	
Braden pressure ulcer assessment per policy (daily or every shift)	
Skin assessment (nursing) completed upon admission	

**Review medical record 3 days prior to the documented development of the HAPU**  
Assign the appropriate score for each item:  
1= NO, not appropriate 2= YES, appropriate

**2. Defined and Implemented Intervention(s) Consistent With Patient's Needs**

	HAPU DAY 0 Date _____	1 day prior to HAPU Date _____	2 days prior to HAPU Date _____	3 days prior to HAPU Date _____
2.1 Sensory Perception Interventions Appropriate?				
2.2 Moisture Interventions Appropriate?				
2.3 Activity Interventions Appropriate?				
2.4 Mobility Interventions Appropriate?				
2.5 Nutritional Interventions Appropriate?				
2.6 Friction and Shear Interventions Appropriate?				
<b>3. Monitored/Evaluated Impact of Interventions</b>				
Skin assessment completed every shift				
<b>4. Revised Interventions as Appropriate</b>				

**Summary/Conclusion**

5. Pressure Ulcer Avoidable \_\_\_\_\_

**Figure 2 Pressure Ulcer Prevention Inventory.**  
Reproduced with permission from Indiana University Health. ©2018 Indiana University Health  
Abbreviations: HAPU, hospital-acquired pressure ulcer; sDTI, suspected deep tissue injury. (The Pressure Ulcer Prevention Inventory was developed before the National Pressure Ulcer Advisory Panel changed the staging terminology in 2016.)

ill en- ite  
gator- ronic d to 14 for HAPI ond data  
Imis- f the infor- res, unt in body tion idities res, it exam- ed no 2 days un-  
line.org

# BRADEN INVENTORY WORKSHEET

HAPU Acquisition Date: \_\_\_\_\_

Days Prior to HAPU	Date	2.0 Braden Scale														Total Daily Braden Score	
		Sensory Perception		Moisture		Activity		Mobility		Nutrition		Friction and Shear					
		DAY	NIGHT	DAY	NIGHT	DAY	NIGHT	DAY	NIGHT	DAY	NIGHT	DAY	NIGHT	DAY	NIGHT		
-1PUDAY																	
-2PUDAY																	
-3PUDAY																	

Review documentation. Based on the lowest score for each Braden category, check all types of interventions found in documentation.

2.1 Sensory Perception Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Elevate Heels Off Surface			
Reposition Frequently			
Pressure Redistribution			
Bony Prominence Protection			
<b>Total</b>			

2.2 Moisture Management Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Breathable Pads			
Diapers			
Barrier Creams			
Scheduled Voiding			
Urinary Device			
IO Catheter			
Fecal Device			
Extra Linen Change			
<b>Total</b>			

2.3 Activity Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Elevate Heels Off Surface			
Reposition Frequently			
Pressure Redistribution			
Bony Prominence Protection			
Mobility Promotion (PT, OT, ROM)			
<b>Total</b>			

2.4 Mobility Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Elevate Heels Off Surface			
Reposition Frequently			
Pressure Redistribution			
Mobility Promotion (PT, OT, ROM)			
Rotoprone, Rotorest			
Exception Documented			
<b>Total</b>			

2.5 Nutrition Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Enteral Nutrition			
Parenteral Nutrition			
PO Diet			
Supplements			
Contraindications Documented			
Dietary Consult			
<b>Total</b>			

2.6 Friction and Shear Interventions	-1PU DAY	-2PU DAY	-3PU DAY
Elevate Heels Off Surface			
Reposition Frequently			
Pressure Redistribution			
Mobility Promotion (PT, OT, ROM)			
Drawsheet/Lift Sheet			
HOB $\leq$ 30 degrees			
Bony Prominence Protection			
Other			
<b>Total</b>			

**Figure 3 Braden Inventory Worksheet.**  
 Reproduced with permission from Indiana University Health. ©2012 Indiana University Health  
 Abbreviations: HAPU, hospital-acquired pressure ulcer/injury; HOB, head of bed; IO, in and out; OT, occupational therapy; PO, by mouth; PT, physical therapy; ROM, range of motion.

**BANNER BAR**

1. Subject ID: \_\_\_\_\_

2. Age: \_\_\_\_\_

3. Gender: 1=Male 2=Female \_\_\_\_\_

4. Hospital: 1 2 3 4 5 6 7 \_\_\_\_\_

5. LOS prior to HAPU (in days): \_\_\_\_\_

6. Mortality this admission: 1=NO 2=YES \_\_\_\_\_

**FACE SHEET**

7. Race: 1=African American 2=White 3=Asian 4=Native American  
5=Pacific Islander/Hawaiian 6=Other \_\_\_\_\_

**CLINICAL NOTES**

8. Admitting diagnosis: \_\_\_\_\_

9. Patient admitted from: 1=Home 2=OSH 3=LTAC 4=Nursing Home 5=Other \_\_\_\_\_

10. Any hospital admission within 30 days prior to HAPU: 1=NO 2=YES 3=UNKNOWN \_\_\_\_\_

11. Smoker: 1=Never 2=Past 3=Current \_\_\_\_\_

12. Comorbidities:

1=MI	6=Chronic pulmonary disease	11=Hemiplegia/paraplegia
2=CHF	7=Rheumatic disease	12=Renal disease
3=PVD	8=PUD	13=Malignancy
4=Cerebrovascular disease	9=Liver disease	14=AIDS/HIV
5=Dementia	10=DM	15=NONE

Total # of comorbidities: \_\_\_\_\_

13. History of pressure ulcer: 1=NO 2=YES 3=UNKNOWN \_\_\_\_\_

14. Previous pressure ulcer location:

1=Sacrum/coccyx	4=Heel	7=Other
2=Ischium	5=Occipital	8=NA
3=Trochanter	6=Ear	

15. Previous PU laterality: 1=Left 2=Right 3=Midline 4=NA \_\_\_\_\_

16. Previous Pressure Ulcer Stage:

0=Unknown	3=Stage 3	6=Indeterminate
1=sDTI	4=Stage 4	7=NA
2=Stage 2	5=Unstageable	

17. Previous PU device related? 1=NO 2=YES 3=NA \_\_\_\_\_

**FORMS**

18. Unintentional weight loss of 10 or more pounds in 1 month? 1=NO 2=YES 3=UNKNOWN \_\_\_\_\_

**RESULTS REVIEW**

19. Height (cm): \_\_\_\_\_

20. Admission weight (kg): \_\_\_\_\_

21. Pressure ulcer location:

1=Sacrum/coccyx	3=Trochanter	5=Occipital	7=Other
2=Ischium	4=Heel	6=Ear	

22. Pressure ulcer laterality: 1=Left 2=Right 3=Midline \_\_\_\_\_

23. Pressure ulcer stage:

1=sDTI	3=Stage 3	5=Unstageable
2=Stage 2	4=Stage 4	6=Indeterminate

24. Pressure ulcer device related? 1=NO 2=YES \_\_\_\_\_

Com

Figure 4 Data collected from electronic medical record.  
 Reproduced with permission from Indiana University Health. ©2018 Indiana University Health.

	PU Date	-1 Day	-2 Days	-3 Days
<b>RESULTS REVIEW</b>				
25. Daily weight:				
26. BMI:				
27. Mechanical ventilation: 1=NO 2=YES				
28. MAP <60: 1=NO 2=YES				
29. SBP <90: 1=NO 2=YES				
30. FiO2 (highest):				
31. O2 saturation (lowest):				
32. Temperature (highest):				
33. Temperature (lowest):				
34. Glucose (highest):				
35. Glucose (lowest):				
36. Hemoglobin (lowest):				
37. Blood pH (lowest):				
38. ScVO2/SVO2:				
39. Any stage I PU present? 1=NO 2=YES				
<b>EXPLORER MENU</b>				
40. Unit/location: 1=CC 2=PCU 3=M/S 4=ED 5=OR				
<b>MAR SUMMARY</b>				
41. Chemically sedated (IV drip): 1=NO 2=YES				
42. Paralytic agent >2 hrs: 1=NO 2=YES				
43. Vasopressors (concurrently): 0=NONE 1=1 2=2 3=3 or more				
44. Steroids: 1=NO 2=YES				
<b>CLINICAL NOTES</b>				
45. Perioperative/procedure time >4 hrs: 1=NO 2=YES				
46. Any type of dialysis: 1=NO 2=YES				
<b>I/O FLOW SHEET</b>				
47. Nutrition: 1=NPO 2=Tube feed 3=TPN 4=Oral 5=Multiple				
48. Bed type: 1=Standard for unit 2=Specialty				
49. Specialty bed type: 1=Bariatric 2=Rotoprone 3=Rotorest 4=Clinitron				
50. Waffle mattress: 1=NO 2=YES				
51. Moisture: 0=None 1=Urinary incontinence 2=Fecal incontinence 3=Both 4=Other				
52. Bowel management system? 1=NO 2=YES				
53. Lowest daily Braden score:				
54. Capillary refill >3 seconds? 1=NO 2=YES				
55. APACHE score within 24 hrs of ICU admit:				
56. APACHE score 3 days prior to PU:				

Figure 4 Continued

Abbreviations: APACHE, Acute Physiology and Chronic Health Evaluation; BMI, body mass index; CC, critical care unit; CHF, congestive heart failure; DM, diabetes mellitus; ED, emergency department; FiO2, fraction of inspired oxygen; HAPU, hospital-acquired pressure injury/ulcer; ICU, intensive care unit; ID, identification; I/O, Inspection/Observation; IV, intravenous; LOS, length of stay; LTAC, long-term acute care hospital; MAP, mean arterial pressure; MAR, medication administration record; MI, myocardial infarction; M/S, medical/surgical unit; NA, not applicable; NPO, nothing by mouth; O2, oxygen; OR, operating room; OSH, outside hospital; PCU, progressive care unit; PU, pressure ulcer; PUD, peptic ulcer disease; PVD, peripheral vascular disease; SBP, systolic blood pressure; ScVO2, central venous oxygen saturation; sDTI, stage deep tissue injury; SvO2, mixed venous oxygen saturation; TPN, total parenteral nutrition.



**Table 1**  
Participants' demographic and clinical data

Variable	No. (%) of participants <sup>a</sup>			P
	Total (N=165)	Avoidable (n=98)	Unavoidable (n=67)	
Age, mean (SD), y	59.9 (16.4)	61.5 (16.1)	57.7 (16.6)	.14
Acute Physiology and Chronic Health Evaluation (APACHE) score within 24 hours of ICU admission, mean (SD)	23.8 (8.7)	23.6 (8.2)	24.0 (9.4)	.72
Total No. of comorbid conditions, mean (SD)	1.5 (1.2)	1.7 (1.3)	1.3 (1.1)	.08
Days in hospital before HAPI was documented, mean (SD)	15.1 (13.7)	13.4 (12.7)	17.6 (14.8)	.06
Female sex	67 (41)	43 (44)	24 (36)	.30
Race				.43
White	132 (83)	81 (87)	51 (77)	
Nonwhite	26 (16)	12 (13)	14 (21)	
Multiracial	1 (1)	0 (0)	1 (2)	
Ethnicity				.43
Hispanic or Latinx	1 (1)	0 (0)	1 (3)	
Not Hispanic or Latinx	90 (99)	52 (100)	38 (97)	
Smoker				.05
Never	68 (45)	35 (39)	33 (55)	
Current/past	82 (55)	55 (61)	27 (45)	
Congestive heart failure				.01
No	142 (87)	80 (82)	62 (95)	
Yes	21 (13)	18 (18)	3 (5)	
Unintentional weight loss of ≥10 lb (4.5 kg) in 1 month				.91
No	121 (86)	71 (86)	50 (86)	
Yes	20 (14)	12 (14)	8 (14)	
History of pressure injury				.10
No	114 (86)	68 (91)	46 (81)	
Yes	18 (14)	7 (9)	11 (19)	
Pressure injury stage				.07
Deep tissue pressure injury	102 (63)	67 (70)	35 (52)	
2	34 (21)	17 (18)	17 (25)	
3	1 (1)	0 (0)	1 (1)	
4	1 (1)	1 (1)	0 (0)	
Unstageable	25 (15)	11 (11)	14 (21)	
Medical device–related pressure injury				.27
No	105 (64)	59 (60)	46 (69)	
Yes	60 (36)	39 (40)	21 (31)	
Pressure injury location				.78
Sacrum/coccyx	70 (42)	41 (42)	29 (43)	
Ischium	4 (2)	1 (1)	3 (4)	
Heel	23 (14)	14 (14)	9 (13)	
Occipital	3 (2)	2 (2)	1 (1)	
Ear	13 (8)	7 (7)	6 (9)	
Other	52 (32)	33 (34)	19 (28)	
Mortality				.24
Yes	45 (27)	30 (31)	15 (22)	
Mechanical ventilation				.49
No	35 (21)	19 (19)	16 (24)	
Documented at least once in the 3 days before HAPI and day when HAPI was first documented	130 (79)	79 (81)	51 (76)	
Chemically sedated (intravenous infusion)				.003
No	73 (44)	34 (35)	39 (58)	
Documented at least once in the 3 days before and the day when HAPI was first documented	92 (56)	64 (65)	28 (42)	
Systolic blood pressure <90 mm Hg				.046
No	59 (36)	29 (30)	30 (45)	
Documented at least once in the 3 days before and the day when HAPI was first documented	106 (64)	69 (70)	37 (55)	

Continued

Continued

Variable	No. (%) of participants <sup>a</sup>			P
	Total (N=165)	Avoidable (n=98)	Unavoidable (n=67)	
Mean arterial pressure < 60 mm Hg				
No				
Documented at least once in the 3 days before and the day when HAPI was first documented	65 (39)	33 (34)	32 (48)	.07
Vasopressors (concurrently)	100 (61)	65 (66)	35 (52)	
No				
Documented at least once in the 3 days before and the day when HAPI was first documented	97 (59)	50 (51)	47 (70)	.01
Incontinence	68 (41)	48 (49)	20 (30)	
No				
Documented at least once in the 3 days before and the day when HAPI was first documented	74 (45)	43 (44)	31 (46)	.76
Bowel management system	91 (55)	55 (56)	36 (54)	
No				
Documented at least once in the 3 days before and the day when HAPI was first documented	131 (79)	83 (85)	48 (72)	.04
	34 (21)	15 (15)	19 (28)	

Abbreviations: HAPI, hospital-acquired pressure injury; ICU, intensive care unit.  
<sup>a</sup> Values are mean (SD) in the first 4 rows; all remaining entries are No. (%) of participants. Some percentages do not total 100 because of rounding.

**Table 2**  
 Univariate models of factors associated with unavoidable hospital-acquired pressure injury (HAPI)

Variable	Mean (SD)		Odds ratio <sup>a</sup> (95% CI)	P
	Avoidable	Unavoidable		
Days in hospital before HAPI occurred	13.4 (12.7)	17.6 (14.8)	1.02 (1.00-1.05)	.07
Total No. of comorbidities	1.7 (1.3)	1.3 (1.1)	0.79 (0.60-1.03)	.08
Daily Braden Inventory Worksheet (data reflect the mean of day and night Braden Scale scores totaled, then the mean across 3 days)				
Mobility	2.0 (0.7)	2.2 (0.8)	1.35 (0.88-2.07)	.18
Activity	1.3 (0.5)	1.4 (0.7)	1.56 (0.91-2.67)	.11
Sensory perception	2.7 (0.8)	2.8 (0.9)	1.14 (0.78-1.68)	.49
Moisture	3.2 (0.5)	3.2 (0.5)	0.90 (0.49-1.66)	.74
Nutrition	2.5 (0.5)	2.7 (0.4)	2.82 (1.27-6.24)	.01
Friction and shear	1.9 (0.5)	1.9 (0.5)	0.79 (0.43-1.44)	.44
Braden Scale total score	13.6 (2.5)	14.2 (2.7)	1.09 (0.97-1.23)	.16
No. of pressure ulcer preventive interventions				
Mobility	2.6 (0.7)	3.1 (0.6)	4.02 (2.17-7.43)	<.001
Activity	3.4 (0.7)	4.0 (0.7)	4.12 (2.24-7.58)	<.001
Sensory perception	3.4 (0.7)	3.8 (0.5)	2.96 (1.55-5.64)	.001
Moisture	2.8 (0.6)	2.9 (0.7)	1.34 (0.80-2.22)	.26
Nutrition	1.9 (0.7)	2.3 (0.5)	3.09 (1.75-5.47)	<.001
Friction and shear	4.7 (1.1)	5.3 (1.1)	1.70 (1.22-2.36)	.002

<sup>a</sup> Odds of being an unavoidable HAPI with continuous variable in first column controlled for.

of having an unavoidable HAPI, and participants with a history of pressure injury were 5 times more likely to have an unavoidable HAPI (Table 3).

### Discussion

An important finding of this study was the identification of 41% (n=67) of the HAPIs as being

unavoidable. Using a valid and reliable instrument (PUI) provided an objective measure to identify unavoidable HAPIs. No other similar studies using such a tool were found in the literature.

In addition, this study is unique in being the first of its kind with the aim of objectively quantifying pressure injury prevention interventions in use

**Table 3**  
Multivariate models of clinical factors associated with unavoidable hospital-acquired pressure injury (HAPI)

Risk factor	Odds ratio <sup>a</sup> (95% CI)	P
Days in hospital before HAPI	1.04 (1.002-1.08)	.04
Total No. of comorbidities	0.96 (0.65-1.43)	.85
Daily Braden Scale total score per unit increase	1.12 (0.92-1.36)	.27
History of pressure injury	5.27 (1.20-23.15)	.03
Chemically sedated (intravenous infusion)	0.49 (0.20-1.25)	.14
Congestive heart failure	0.028 (0.002-0.36)	.006
Systolic blood pressure < 90 mm Hg	0.82 (0.31-2.22)	.70
Vasopressors (concurrently)	0.45 (0.16-1.24)	.12
Bowel management system	1.49 (0.47-4.71)	.50
Smoker	0.92 (0.38-2.19)	.84

<sup>a</sup> Odds of being an unavoidable HAPI with clinical risk factor controlled for.

before pressure injury development. The only other study we found that examined appropriate pressure injury preventive care was that of Beekman and colleagues,<sup>15</sup> a randomized controlled trial of 464 nursing home residents in 4 nursing homes in Belgium.

**This study provides an objective, though retrospective, means to identify unavoidable pressure injuries.**

In that study, the authors examined adherence to guideline-based pressure injury preventive care recommendations. Pressure injury preventive protocol was tailored to the resident and was described as skin observation, use of support surface, repositioning, and heel elevation.

Preventive interventions were defined as either fully adequate, meaning that all were performed, or not. The results of the study were limited, as the researchers found that fully adequate preventive care was provided to the intervention group only when patients were seated; no improvements to preventive care were found while patients were in bed.<sup>15</sup> Preventive interventions were not described in as much detail as in the present study. The main factor distinguishing avoidable from unavoidable HAPIs in the present study was the number of interventions documented. The unavoidable HAPI group had more interventions documented than the avoidable HAPI group; this finding makes sense given that the documented nursing care was deemed appropriate to the patient condition in the unavoidable HAPI group.

Another interesting finding of the present study was the high proportion (36%) of medical device-related HAPIs. This finding is most likely due to the nature of the study population and the large number of devices used in critical care areas.

The finding is consistent with the results of Black and colleagues,<sup>16</sup> who reported a HAPI rate of 5.4% (113 of 2079), with 34.5% (39 of 113) being related to medical devices.

In this study, unavoidable HAPIs were defined as those that developed in spite of consistent documentation of evidence-based preventive interventions. Thus, the question arises: Why did a HAPI develop in these individuals? The literature includes discussion of the complexity of pressure injury etiology and the potential for acute skin failure in critically ill patients<sup>4,6,17,20</sup>; however, with our current level of evidence, the distinction between acute skin failure and unavoidable pressure injury remains obscure.

The NPUAP hosted a multidisciplinary conference in 2014 to explore the issue of pressure injury unavoidability using an organ system framework.<sup>5</sup> Participants achieved consensus that unavoidable pressure injuries do indeed occur and that risk factors such as multiorgan dysfunction syndrome, shock or sepsis, hemodynamic instability and impaired tissue oxygenation, cardiac dysfunction, CHF, and skin failure are associated with pressure injury development and increase the likelihood of unavoidable pressure injury development.<sup>5</sup> These factors have also been described as part of the concept of acute skin failure.<sup>18-21</sup>

Langemo and Brown conducted a systematic review and defined skin failure as "an event in which skin and underlying tissues die due to hypoperfusion concurrent with severe dysfunction or failure of other organ systems."<sup>18(p200)</sup> They reported that the term *skin failure* appeared in the literature as early as 1993 and was described as the damage that occurs in skin and underlying tissue at the end of life and in the intensive care setting.<sup>18</sup> Langemo and Brown stated that the distinguishing factor between skin failure and a pressure injury is the coexistence of a significant disease process or organ failure. Their definition of acute skin failure is consistent with the overall characteristics of the patients in our study.

In a study of 552 critical care patients, Delmo and colleagues<sup>19</sup> developed a predictive risk model of development of acute skin failure versus pressure injury in patients admitted to critical care units. Specifically, they identified 5 predictors of acute skin failure: peripheral artery disease, mechanical ventilation lasting more than 72 hours, respiratory failure, liver failure, and severe sepsis or septic shock.<sup>19</sup> In a retrospective correlational study of 347 critical care patients, Cox<sup>20</sup> identified norepinephrine as a predictive risk factor for pressure injuries. Our study results are consistent with these findings, as al

9% (n = 130) of the participants with HAPIs in our study were receiving mechanical ventilation, 56% (n = 92) were chemically sedated, 64% (n = 106) had SBP below 90 mm Hg, 61% (n = 100) had mean arterial pressure below 60 mm Hg, and 41% (n = 68) were receiving 1 or more vasopressors.

Advances in medicine have enabled critically ill patients to survive situations that in the past led to death. Now, increasing numbers of patients with multiorgan failure are surviving but remain highly susceptible to adverse events involving skin integrity, specifically skin failure. In a prospective descriptive study of 29 critical care patients with acute skin failure, Curry and colleagues<sup>21</sup> reported that 5 patients had 2 organ failures at the time that skin failure was noted, 15 patients had either 3 organ failures or 2 organ failures plus sepsis, and 9 patients experienced 4 or more organ system failures and/or sepsis. All patients in that study had a diagnosis of nonskin organ failure and low albumin levels. Although the study of Curry et al<sup>21</sup> provides additional information, it lacks an objective definition of acute skin failure and does not differentiate between pressure injuries and acute skin failure. The present study provides an objective, although retrospective, means to identify pressure injuries that are unavoidable. The results may suggest a process more aligned with acute skin failure than with pressure injury.

The present study's findings regarding risk factors and avoidable HAPIs are difficult to explain, but they may be due to nurses' perceptions of patients' hemodynamic instability and thus their inability to implement preventive interventions, primarily repositioning. Patients with CHE, chemical sedation, SBP less than 90 mm Hg, or vasopressor use were more likely to lack sufficient preventive measures. These findings support current discussions in the literature on the challenges of repositioning critical care patients. Brindle and colleagues<sup>22</sup> reported on the development of consensus recommendations by a group of experts related to safe repositioning of patients. They noted that the critical care unit's culture and clinicians' perceptions about hemodynamic instability may lead to staff members' not repositioning patients. Krapfl and colleagues,<sup>23</sup> in a review of the literature, noted that the complexity and instability of the patient's condition often limit repositioning by the nurse. However, best practice continues to suggest that slow, gradual turning allows sufficient time for stabilization of blood pressure and oxygen saturation and should be considered.<sup>11</sup>

#### Limitations

Several limitations of this study are evident. The study had a retrospective design and relied on the accuracy of documentation, which did not allow confirmation of findings through observation. All participants in the study had pressure injuries, and no case-control group was used for comparison, thus limiting the ability to compare risk factors across groups. The study was conducted at a large health care system with high patient acuity, and the findings may not be generalizable to other populations and settings. Data retrieval to complete the PUIPI relied on the accuracy of the documentation in the EMR. The complexity of the EMR was evident during data collection, even though the EMR software is commonly used and provided by an international EMR software company. Another limitation was the inclusion of progressive care patients, which may have diluted the acuity of the sample. Finally, the PUIPI is based on the Braden Scale subscales and was not originally intended for use with medical device-related pressure injuries, thus limiting its application to this type of HAPI.

#### Implications for Nursing

The findings of this study provide important information and new knowledge for critical care nurses and other health care providers and highlight the importance of pressure injury prevention documentation. Attention to standard nursing care (preventive interventions) and accuracy of documentation is essential in the complex critical care setting. In addition, this study objectively identified the occurrence of unavoidable pressure injuries, suggesting a possible etiology of acute skin failure rather than a lack of preventive nursing care. Further research—particularly rigorous, controlled studies—is needed to investigate the occurrence of unavoidable HAPIs and acute skin failure in critically ill patients.

#### ACKNOWLEDGMENTS

This study was performed at Indiana University Health, Indianapolis, Indiana.

#### FINANCIAL DISCLOSURES

This study was funded by an American Association of Critical-Care Nurses Research Impact Grant.

#### SEE ALSO

For more about pressure injuries, visit the *Critical Care Nurse* website, [www.ccnonline.org](http://www.ccnonline.org), and read the article by McGee et al, "Pressure Injuries at Intensive Care Unit Admission as a Prognostic Indicator of Patient Outcomes" (June 2019).

REFERENCES

- Cuddigan J. Critical Care. In: Phipps R, National Pressure Ulcer Advisory Panel, eds. *Pressure Ulcers: Prevention and Treatment*. Philadelphia: Elsevier, 2014:1-10.
- National Pressure Ulcer Advisory Panel. *National Pressure Ulcer Prevention and Treatment Guidelines*. Washington, DC: Centers for Medicare and Medicaid Services, 2014:247-265.
- Surveys for Long-term Care. *Centers for Medicare and Medicaid Services, Guidance to States*. www.cms.gov/Medicare/Provider-EnrollmentandCertification/05-17-Long-term-Care-Info/downloads/SCLetter05-17.pdf. Accessed May 27, 2019.
- Black J, Edberg L, Baharestani M, et al. Risk factors for avoidable or unavoidable pressure ulcers: results of the National Pressure Ulcer Advisory Panel Consensus Conference. *Ostomy Wound Ostomy and Continence Nurs*. 2016;43(9):242-247.
- Wound, Ostomy and Continence Nurses Society. *WOCN Society Position Paper: Avoidable Versus Unavoidable Pressure Ulcers*. Mt Laurel, NJ: Wound, Ostomy and Continence Nurses Society; 2016. [www.wocn.org/resources/publications/Avoidable and Unavoidable\\_P.pdf](http://www.wocn.org/resources/publications/Avoidable%20and%20Unavoidable_P.pdf). Accessed May 27, 2019.
- Langemo D, Baharestani M, Postauer M, Goldberger M, Unruh M, Langemo D. Risk factors for avoidable and nonavoidable pressure ulcers in hospitalized cardiac surgery patients: a systematic review. *J Wound Ostomy Continence Nurs*. 2016;43(9):242-247.
- Pittman J, Beeson T, Dillon J, et al. Unavoidable pressure ulcers: development and testing of the Indiana University Health Pressure Ulcer Assessment Tool. *J Wound Ostomy Continence Nurs*. 2016;43(1):32-38.
- Bradley B, Bergstrom N. A conceptual schema for the study of the etiology of pressure ulcers. *Rehabil Nurs*. 1987;2(1):8-12.
- Coleman S, Gorecki C, Nelson EA, et al. *Prevention and Treatment of Pressure Ulcers*. In: Healer E, ed. *Prevention and Treatment of Pressure Ulcers: Clinical Practice Guidelines*. Western Australia: Western Australia; 2014:2-61.
- Hessler E, ed. National Pressure Ulcer Advisory Panel. *European Pressure Ulcer Advisory Panel, Pan Pacific Pressure Ulcer Alliance. Prevention and Treatment of Pressure Ulcers: Clinical Practice Guidelines*. Osborne Park, Western Australia, Australia: Cambridge Media; 2014.
- Edberg L, Black J, Goldberg M, McNichol L, Moore L, Slip green M. Revised National Pressure Ulcer Prevention and Treatment Guidelines. *J Wound Ostomy Continence Nurs*. 2016;43(6):358-397.
- National Database of Nursing Quality Indicators. *Guidelines for Data Collection*. www.ndnqi.org. Accessed May 27, 2019.
- Bradley B, Bergstrom N, Braden L, et al. *Pressure Ulcer Risk Assessment*. www.ndnqi.org. Accessed May 27, 2019.
- Sore Risk. <https://www.hhs.gov/od/officeofpublicaffairs/press-releases/2015/05/15/sore-risk/>.
- Verhaeghe S. A multi-faceted tailored strategy to implement an electronic clinical decision support system and pressure ulcer prevention in a long-term care facility. *J Wound Ostomy Continence Nurs*. 2015;42(1):38-48.
- Black J, Cuddigan J, Walke M, Diller LA, Lander M, Kelso A, et al. Medical device related pressure ulcers in hospitalized controlled trial. *Int J Nurs Stud*. 2015;50(4):475-486.
- Schmitt S, Andries M, Ashmore P, Brunette G, Judga K, Bonham P. WOCN Society Position Paper: avoidable versus unavoidable pressure ulcers. *J Wound Ostomy Continence Nurs*. 2017;44(5):458-468.
- Langemo D, Brown G. Skin falls too: acute, chronic and end stage skin failure. *Adv Skin Wound Care*. 2006;19(4):209-214.
- Dalmore B, Cox J, Robinson M, et al. Risk factors for avoidable skin failure in a long-term care patient. *Adv Skin Wound Care*. 2015;28(1):14-23.
- Cox J. Predictors of pressure ulcers in adult critical care patients. *Am J Crit Care*. 2016;21(5):363-368.
- Healer E, et al. *Prevention and Treatment of Pressure Ulcers: A Prospective, Descriptive Study of Characteristics Associated with Skin Failure in Critically Ill Adults*. *Ostomy Wound Care*. 2017;59(5):36-43.
- Bridle C, et al. *Prevention and Treatment of Pressure Ulcers in Critically Ill Patients with Hemodynamic Instability*. *Am J Crit Care*. 2015;20(3):254-267.
- Wound Ostomy Continence Nurs. 2015;40(3):254-267.
- Krapfl L, Langin J, Pike C, Pezalla P. Does medical malpractice litigation affect pressure ulcer prevention? *J Wound Ostomy Continence Nurs*. 2017;44(4):319-322.

To purchase electronic or print reprints, contact American Association of Critical-Care Nurses, 101 Columbia, Avejeff, CA 92656. Phone: (800) 899-1712 or (949) 361-5532; fax: (949) 362-2049; email: reprints@aacn.org

CE 1.0 Hour Category A

**Notice to CE enrollees:** This article has been designated for CE contact hour(s). The evaluation demonstrates your knowledge of the following objectives:

1. Identify 4 components of an unavoidable pressure injury.
2. Describe the purpose of this study.
3. Describe 3 risk factors that were associated with pressure injury development in this study.

To complete the evaluation for CE contact hour(s) for this article #A1928052, visit [www.aacnline.org](http://www.aacnline.org) and click the "CE Articles" button. No CE evaluation fee for AACN members. This expires on September 1, 2020.

The American Association of Critical-Care Nurses is an accredited provider of continuing nursing education. AACN has been approved as a pro-gram for continuing education in nursing by the State Boards of Registered Nursing of California (#01036) and Louisiana (#LSBN12).