



Content Focus: **Interactions** Correct: 8 / 22 Incorrect: 14 / 22 Time Spent: 00:24:25

Incorrect Answers:

Q: What are potential interactions for methotrexate?

A:

Concurrent use of methotrexate and digoxin can reduce digoxin level.

Concurrent use of methotrexate and NSAIDs, salicylates, and sulfonamides can cause toxicity.

Alcohol use can increase risk of

36%

Final Score

Congratulations on completing your Flash Cards.

How would you like to continue?



START OVER



RESHUFFLE



REPLAY
INCORRECT



PRINT RESULTS



Content Focus: **Contraindications** ▼

Correct: 10 / 22

Incorrect: 12 / 22

Time Spent: 00:40:22

Incorrect Answers:

Q: What contraindications are there for the use of carmustine ?

A:

Pregnancy/lactation risk

History of impaired lung function

Recent infection or exposure to herpes or chickenpox

Severe bone marrow

45%

Final Score

Congratulations on completing your Flash Cards.

How would you like to continue?



START OVER



RESHUFFLE



REPLAY
INCORRECT



PRINT RESULTS



Content Focus: **Interventions** Correct: 6 / 22 Incorrect: 16 / 22 Time Spent: 00:56:06

Incorrect Answers:

Q: What interventions should be planned for a client taking cytarabine?

A:
Monitor client's weight and report gastrointestinal effects.

Premedicate with an antiemetic prior to administration of cytarabine.

Monitor CBC, Hgb,

27%

Final Score

Congratulations on completing your Flash Cards.

How would you like to continue?



START OVER



RESHUFFLE



REPLAY
INCORRECT



PRINT RESULTS