

3. Relative Risk Ratio - a ratio of the incidence rate in the exposed group and the incidence rate in the nonexposed group.

Example: Relative Risk = $\frac{\text{Incidence rate in the exposed group}}{\text{Incidence rate in the nonexposed group}}$

**A relative risk of 1.0 indicates that the risk is equal for both groups, and conversely, a relative risk greater than 1.0 indicates that the risk is greater in the exposed group. A relative risk less than 1.0 indicates that the risk is less in the exposed group.

Rates to Determine Validity and Reliability

1. Sensitivity: the ability of a test to identify correctly people who have the health problem under study

Example: Sensitivity = $\frac{\text{true positive results}}{\text{(true positive results + false negatives)}}$

2. Specificity: the ability of a test to correctly identify people who do not have the health problem

Example: Specificity = $\frac{\text{true negative results}}{\text{(true negative results + false positives)}}$

Calculate the following rates using the information provided. Use the scaling factor of 100,000 for all of the problems.

1. The total death in County Z last year was 6,092. The population of County Z last year was 524,263. What was the crude death rate?

$\frac{6092}{524,263} \times 100,000 = 1162$ 11.62 per 1,000

2. There were 4,953 deaths from neoplasms in City B during the past year. The year end population was 3,495,678. What was the specific cancer death rate for last year?

~~4953~~ $\frac{4953}{3,495,678} \times 100,000 = 141.689$ 141.6 per 100,000
142

3. The population of the US in 2000 was 281,421,906. The number of deaths from heart disease in the US in 2000 was 710,760. The total number of deaths in the US in 2000 was 2,403,351.

a Calculate the percentage (%) of heart disease deaths for the US in 2000.

29% $\frac{710,760}{2,403,351} = 0.296 \times 100$ or 29.6%

b Calculate the rate of heart disease deaths in the US in 2000 for the US.

$\frac{710,760}{281,421,906} \times 100,000 = 252.56$ 252.56 per 100,000
25

4. In Illinois in 2000, the population was 12,419,293. The number of Salmonella cases in 2000 was 1,502 in Illinois. Calculate the incidence rate for Salmonella for Illinois in 2000.

$$12.04$$

5. There were 45,238 neonatal deaths out of 5,672,000 live births in City F. Calculate the neonatal mortality (death) rate.

$$\frac{45,238}{5,672,000} \times 1000 = \boxed{7.97 \text{ per } 1,000}$$

6. The population in Sangamon county in 2000 was 188,951

- a The number of live births in Sangamon County in 2000 was 2,646. Figure the Live Birth rate for Sangamon country for 2000.

$$14\% \quad \frac{2646}{188951} \times 1000 = \boxed{14\%}$$

- b The number of infant deaths in Sangamon County in 2000 was 18. Figure the infant mortality rate for Sangamon country for 2000.

$$70\% \quad \frac{18}{2646} \times 1000 = \boxed{6.8\%}$$

7. A city has a population of 250,000. Of these, 10,000 have disease X, which is incurable. There are 1,000 new cases and 400 deaths each year from this disease. There are 2,500 deaths per year from all causes. What is the prevalence rate based on a multiplier 100,000.

$$\boxed{4,000 \text{ per } 100,000}$$

$$\frac{10,000}{250,000} = 0.04 \times 100,000$$

	Quintile of CRP Level				
	1	2	3	4	5
	0.49 mg/dL	>0.49-1.08 mg/dL	>1.08-2.09 mg/dL	>2.09-4.19 mg/dL	>4.19 mg/dL
Relative Risk	1.0	1.8	2.3	3.2	4.5
Number of women	6000	6000	6000	6000	6000

8. Based on the relative risk data above, one can conclude:
- There is no risk of heart attack/stroke for women with CRP levels in the first quintile.
 - Decreasing CRP level appears to increase the risk of heart attack/stroke.
 - Increasing CRP level appears to increase the risk of heart attack/stroke.
 - There appears to be no association between CRP levels and heart attack/stroke.

9. In 2020, the population of Illinois is 12.63 million. Total cases of COVID-19 is 900,370. Using 100,000 as a multiplier, what is the period prevalence rate?

$$\frac{900,370}{12.63 \text{ mil}} \times 100,000 = 7128.8 \text{ per } 100,000$$

10. In 2020, the population of Illinois is 12.63 million. The total deaths from COVID-19 is 128,000. Using 100,000 as a multiplier, what is the cause-specific mortality rate?

$$\frac{128,000}{12,630,000} \times 100,000 = 1013.46 \text{ per } 100,000$$

EPIDEMIOLOGY EXERCISES
 INFANT MORTALITY, CHICAGO COMMUNITY AREAS

Anderson

COMMUNITY AREA	# of LIVE BIRTHS	DEATHS UNDER ONE YEAR	
		#	RATE PER 1000 BIRTHS
			12.3
Argyle Park	1,064	13	5.2
West Ridge	966	5	19.4
Uptown	1,580	26	17.1
Lincoln Square	760	13	11.5
South Center	610	7	24.9
East Garfield Park	763	19	25.4
New West Park	1,338	34	27.1
Oakland	295	8	19.9
Grand Boulevard	1,209	24	23.1
Washington Park	735	17	20.7
Englewood	1,303	27	33.5
Washington Heights	507	17	
CHICAGO	55,216	914	16.6
UNITED STATES	--	--	12.5

A. Fill in the blank columns in Table 1 using the formula to calculate infant mortality rate. A hand or desk calculator speeds calculations but is not essential. ✓

B. Compare the infant mortality rate you calculated for Lincoln Square with that of East Garfield Park. Are they the same or different?

LS 17.1
 EGP 24.9

C. What general trends, if any, are apparent from these data?

↑ death rates in communities with multiracial minorities.

D. Compare the infant mortality rate of the City of Chicago with that of the United States. Are the rates the same or different? What factors may account, between Chicago and the United States, affect these rates?

Chi 16.6
 US 12.5

Resources to the vulnerable population may be a large factor.