



Individual Performance Profile

DOWNLOAD REPORT

Score Explanation

ADJUSTED INDIVIDUAL TOTAL SCORE 83.3% TIME SPENT 55:26	Individual Name: conor deering Student Number: cdeering Institution: Lakeview CON Program Type: BSN	Focused Review Progress View missed topics and launch study materials below. Last accessed: 12/11/2021 Time spent: 01:32:59
	Test Completed Date: 12/9/2021 # of Points: 60 Attempt: 1	

Proficiency Level	Mean		Percentile Rank	
Level 3	National 66.6%	Program 67.2%	National 93	Program 92

Individual Performance in the Major Content Areas Show all topics to review OFF

Content Area	Topics to Review	Total # Points	MEAN		PERCENTILE RANK		Individual Score	
			National	Program	National	Program		
Management of Care	0	2	n/a	n/a	n/a	n/a	100.0%	
+ Safety and Infection Control	2	3	n/a	n/a	n/a	n/a	33.3%	<div style="background-color: #92d050; padding: 5px; text-align: center;"> FOCUSED REVIEW > </div> Last Accessed: 12/11/2021
+ Health Promotion and Maintenance	2	13	65.3%	65.7%	93	92	84.6%	<div style="background-color: #92d050; padding: 5px; text-align: center;"> FOCUSED REVIEW > </div> Last Accessed:

● **Accident/Error/Injury/Prevention - (2)**

- Assessment and Management of Newborn Complications: Caring for a Newborn Who Has Abstinence Syndrome (Active Learning Template - Basic Concept, RM MN RN 11.0 Chp 27 Assessment and Management of Newborn Complications)
 - Seizures, disturbed sleep patterns, stillbirth, and SIDS incidences are increased.
 - Infants suffering withdrawal can have poor feeding, regurgitation, and diarrhea.
 - Increased crying, irritability, and increased muscle tone occur with abstinence syndrome.
- Nursing Care and Discharge Teaching: Evaluating Understanding of Car Seat Safety (Active Learning Template - Basic Concept, RM MN RN 11.0 Chp 26 Nursing Care and Discharge Teaching)
 - Use a rear-facing car seat that is officially approved for use.
 - It is preferable to place the car seat in the middle portion of the back seat.
 - A car seat should be used until age 2 or if the child reaches the maximum height and weight for the seat.

● **Ante/Intra/Postpartum and Newborn Care - (2)**

- Labor and Delivery Processes: Caring for a Client Who Is in the Second Stage of Labor (Active Learning Template - Basic Concept, RM MN RN 11.0 Chp 11 Labor and Delivery Processes)
 - Involuntary powers and the mother consciously pushing helps bring out the fetus.
 - Frequent position changes help reduce pain as well as aid in fetal descent.
 - It is important that the nurse focus on the mother's anxiety level and provide comfort.
- Prenatal Care: Expected Changes During the Second Trimester (Active Learning Template - Basic Concept, RM MN RN 11.0 Chp 4 Prenatal Care)
 - Fundal height measurement starts in the second trimester.
 - Fundal assessment measurements should match fundal height within a deviation of 2cm.
 - Fetal movement can be felt within the second trimester.
- **Nutrition and Oral Hydration - (1)**
 - Sources of Nutrition: Teaching a Client About High-Calcium Food (Active Learning Template - Basic Concept, RM Nutrition 7.0 Chp 1 Sources of Nutrition)
 - Major sources of calcium are broccoli, kale, and dairy products.
 - Excess calcium can cause constipation, kidney stones, and depressed deep-tendon reflexes.
 - Vitamin D aids in calcium absorption and is necessary for facilitating it.
- **Medication Administration - (2)**
 - Early Onset of Labor: Evaluating Understanding of Teaching About Terbutaline (Active Learning Template - Medication, RM MN RN 11.0 Chp 10 Early Onset of Labor)
 - Terbutaline is contraindicated in patients with severe gestational hypertension.
 - Monitor patients for chest discomfort, vomiting, or hypotension after terbutaline administration.
 - Terbutaline 0.24mg can be administered every 4 hours for 24 hours.
 - Therapeutic Procedures to Assist with Labor and Delivery: Labor Induction Using Oxytocin (Active Learning Template - Medication, RM MN RN 11.0 Chp 15 Therapeutic Procedures to Assist with Labor and Delivery)
 - Labor can be induced with IV oxytocin.
 - The nurse should confirm the fetus is engaged at a minimum station of 0 before oxytocin administration.
 - Assess blood pressure, pulse, and respirations every 30 minutes with oxytocin administration.
- **Potential for Complications of Diagnostic Tests/Treatments/Procedures - (1)**
 - Therapeutic Procedures to Assist with Labor and Delivery: Newborn Complications Following a Forceps-Assisted Birth (Active Learning Template - Therapeutic Procedure, RM MN RN 11.0 Chp 15 Therapeutic Procedures to Assist with Labor and Delivery)
 - Fetal heart rate needs to be monitored before, during and after forceps delivery.
 - Compression of the cord by the forceps will decrease the fetal heart rate.
 - Assess the newborn for facial palsy after applying forceps.
- **Alterations in Body Systems - (2)**
 - Assessment and Management of Newborn Complications: Caring for a Newborn Who Has Necrotizing Enterocolitis (Active Learning Template - System Disorder, RM MN RN 11.0 Chp 27 Assessment and Management of Newborn Complications)
 - Necrotizing enterocolitis is an inflammatory process that causes ischemia of the bowel.
 - Short-gut syndrome can be secondary to necrotizing enterocolitis.
 - Necrotising enterocolitis can result in sepsis if not treated.

- Infections: Planning Care for a Client Who Has HIV (Active Learning Template - System Disorder, RM MN RN 11.0 Chp 8 Infections)
 - Risk factors of HIV are IV drug use, multiple sexual partners, and a multi-STI history.
 - Early discovery of HIV decreases the chance of transmission to the newborn.
 - Amniocentesis procedure should be avoided due to exposing the baby to maternal blood.