

N321 CARE PLAN

N321 Care Plan # 1

Lakeview College of Nursing

Name

Shivani Patel

N321 CARE PLAN

Demographics (3 points)

Date of Admission 3/1/2022	Client Initials M.Q.	Age 66	Gender Female
Race/Ethnicity Asian	Occupation Not employed	Marital Status Single	Allergies Plasma Human
Code Status Full code	Height 5'1"(154.9)	Weight 137 lbs	

Medical History (5 Points)

Past Medical History: Allergic rhinitis, anxiety, borderline diabetes, CVA, depression, GERD, HTN, hyperthyroidism, diffuse large B cell lymphoma, abdominal fistula, pancytopenia

Past Surgical History: C-section, cholecystectomy, endoscopic ultrasound, IR venous access, Lasik, port placement, stereotactic core biopsy, upper gastrointestinal endoscopy

Family History: Father/Mother- heart attack, Maternal grandmother- hypertension

Social History (tobacco/alcohol/drugs including frequency, quantity and duration of use):

History of smoking cigarettes

Quit date: 1/1/2001

Quantity and duration of use: not on file

Assistive Devices: None

Living Situation: Lives alone at home

Education Level: College graduate

Admission Assessment

Chief Complaint (2 points): The patient came into the emergency department with complaints of abdominal pain.

N321 CARE PLAN

History of Present Illness – OLD CARTS (10 points): The patient was admitted to the emergency department with abdominal pain, and with a diagnosis of a mesenteric tumor in the abdomen. The pain started a week ago in the middle of the abdomen. The patient has been experiencing pain in her abdomen for about a week. The pain radiates throughout the middle of her abdomen. The pain is intermittent, and she describes it as “shooting and stabbing” pain. When the patient moves around too much, she starts to exhibit an increase in pain. The patient takes acetaminophen (Tylenol) to relieve pain. Resting also helps to reduce the pain. The patient has previously been in the ED in February regarding abdominal pain, but she was diagnosed with an abdominal fistula. At this moment, on a scale of 1-10, the patient states they have been at a pain level of 0.

Primary Diagnosis

Primary Diagnosis on Admission (2 points): Mesenteric tumor (perforation of mesenteric abdominal pass)

Secondary Diagnosis (if applicable): Intra-abdominal abscess

Pathophysiology of the Disease, APA format (20 points):

The mesentery is a set of tissues located in the abdomen, and it attaches the intestine to the wall of the abdomen and attaches it into place. Mesenteric tumors are rare and consist of a group of lesions. Masses can arise from any mesenteric components like the peritoneum, lymphatic tissue, fat, and connective tissue. The tumors can be classified as solid or cystic, benign or malignant (Dufay et al., 2019). A mesenteric tumor larger than 2cm can signify a pN2 disease. Lymphoma is the most common solid mesenteric tumor. Some signs and symptoms of mesenteric tumors are nausea, vomiting, diarrhea, bloating, and constipation.

N321 CARE PLAN

The patient can experience a fast heart rate and respirations due to nausea. Potassium levels are usually lower with a mesenteric tumor. While the diagnosis is formed from a CT scan, the most common symptoms leading to the diagnosis of a mesenteric tumor are abdominal pain or mass. A complete blood cell count is necessary to diagnose a mesenteric mass. The treatment of mesenteric tumors is surgery which consists of local excision. Most cystic mesenteric lesions can be easily excised. It is important to drain the abdominal abscess caused by the mesenteric tumor. The patient was diagnosed with a mesenteric tumor considering she was experiencing abdominal pain. A CT scan of the patient's abdomen was performed and it showed a large mesentery necrotic mass with internal abscess.

Pathophysiology References (2) (APA):

Sembawa, H. A. (2020). Primary mesenteric gastrointestinal stromal tumor presented with acute small bowel obstruction. *National Health Institutes of Health*, 8(1), 1-10.

<https://doi.org/10.1093/jscr/rjaa225>

Dufay, C., Abdelli, A., Le Pennec, V., & Chiche, L. (2019). Mesenteric tumors: Diagnosis and treatment. *Journal of Visceral Surgery*, 149(4), e239-e251.

<https://doi.org/10.1016/j.jviscsurg.2012.05.005>

Laboratory Data (15 points)

N321 CARE PLAN

CBC Highlight All Abnormal Labs—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason for Abnormal Value
RBC	3.50-5.20	2.64	2.86	Low levels of RBCs can be due to anemia (Pagana et al., 2019).
Hgb	11.0-16.0	7.6	8.5	The patient has low iron (Pagana et al., 2019).
Hct	34.0-47.0%	23.5%	26.9%	Low levels of hematocrit can be due to nutritional problems like low iron (Pagana et al., 2019).
Platelets	140-400	85	87.6	Low platelet levels can indicate thrombocytopenia (Pagana et al., 2019).
WBC	4.00-11.00	10.49	6.91	
Neutrophils	47.0%-73.0%	N/A	N/A	
Lymphocytes	18.0%-42.0%	1.7%	8.8%	Low levels indicate the blood doesn't have enough white blood cells. It can also indicate infections (Pagana et al., 2019).
Monocytes	4.0%-12.0%	2.6%	2.3%	Low levels of monocytes can result from chemotherapy (Pagana et al., 2019).
Eosinophils	0.0-5.0%	0.0	0.0	
Bands	Less than or equal to 10%	5.2%	N/A	

Chemistry Highlight All Abnormal Labs—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason For Abnormal
Na-	136-145	134	135	Low levels of sodium can be caused by nausea and muscle weakness (Pagana et al., 2019).
K+	3.5-5.1	3.2	4.7	Low potassium may be a result of diarrhea (Pagana et al., 2019).
Cl-	98-107	102	109	High levels can indicate dehydration (Pagana et al., 2019).
CO2	220-290	250	190	It can indicate an electrolyte imbalance, or that there is a problem removing carbon dioxide

N321 CARE PLAN

				through your lungs (Pagana et al., 2019).
Glucose	74-100	109	168	The cause of high blood sugar can be caused by dehydration and a lack of exercise (Pagana et al., 2019).
BUN	10-20	10	22	High levels of BUN can be due to dehydration (Pagana et al., 2019).
Creatinine	0.55-1.02	0.69	0.59	
Albumin	3.4-4.8	2.6	N/A	Low levels of albumin can be caused by malnutrition (Pagana et al., 2019).
Calcium	8.9-10.6	8.1	N/A	Low levels of calcium can be due to a disorder that interferes with the body's ability to absorb calcium (Pagana et al., 2019).
Mag	1.6-2.6	1.3	N/A	It is caused by decreased absorption of magnesium in the gut or increased excretion of magnesium in the urine (Pagana et al., 2019).
Phosphate	2.5-4.5	N/A	N/A	
Bilirubin	0.2-0.5	N/A	N/A	
Alk Phos	40-150	147	N/A	
AST	5-34	16	N/A	
ALT	0-55	19	N/A	
Amylase	40-140	N/A	N/A	
Lipase	8-78	13	N/A	
Lactic Acid	0.50-2.20	0.57	N/A	

Other Tests **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

N321 CARE PLAN

Lab Test	Normal Range	Value on Admission	Today's Value	Reason for Abnormal
INR	2.0-3.0	N/A	N/A	
PT	11-13.5	N/A	N/A	
PTT	25-35	N/A	N/A	
D-Dimer	less than 0.5	N/A	N/A	
BNP	less than 100	N/A	N/A	
HDL	45-70	N/A	N/A	
LDL	less than 100	N/A	N/A	
Cholesterol	less than 200	N/A	N/A	
Triglycerides	less than 150	N/A	N/A	
Hgb A1c	less than 5.7%	N/A	N/A	
TSH	0.5-5.0	N/A	N/A	

Urinalysis **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab Test	Normal Range	Value on Admission	Today's Value	Reason for Abnormal
Color & Clarity	Yellow/clear	Straw/clear	N/A	Straw- colored urine can indicate dehydration (Pagana et al., 2019).
pH	5.0-7.0	7.0	N/A	
Specific Gravity	1.003-1.035	1.035	N/A	
Glucose	Negative	Negative	N/A	
Protein	Negative	Negative	N/A	
Ketones	Negative	Negative	N/A	
WBC	0-25	5	N/A	

N321 CARE PLAN

RBC	0-20	4	N/A	
Leukoesterase	Negative	Negative	N/A	

Cultures **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Test	Normal Range	Value on Admission	Today's Value	Explanation of Findings
Urine Culture	Negative	N/A	N/A	
Blood Culture	No growth	No growth	N/A	
Sputum Culture	Negative	N/A	N/A	
Stool Culture	Negative	N/A	N/A	

Lab Correlations Reference (1) (APA):

Pagana, K. D., Pagana, T. J., & Pagana T. N. (2019). *Mosby's diagnostic and laboratory desk reference* (14th ed.). Elsevier.

Diagnostic Imaging

All Other Diagnostic Tests (5 points):

The patient had a CT of the abdomen done and a CT of the pelvis with contrast. A CT of the abdomen and pelvis helps to detect diseases of the small bowel, colon, and other internal organs. It is performed to assess tumors, lesions, obstructions, and pain regarding the abdomen.

Diagnostic Test Correlation (5 points):

N321 CARE PLAN

The CT scan revealed a central mesentery large necrotic mass with internal abscess. There is a persistent fistulous track with an adjacent small bowel loop. There is a slight gas leaking into the adjacent mesenteric fat. There is no acute bowel obstruction, but the large and small bowel loops are slightly prominent showing fluid and gas-filled content. There was a small bowel perforation from the lymphoma mass.

Diagnostic Test Reference (1) (APA):

Dufay, C., Abdelli, A., Le Pennec, V., & Chiche, L. (2019). Mesenteric tumors: diagnosis and treatment. *Journal of Visceral Surgery*, 149(4), e239-e251.

<https://doi.org/10.1016/j.jvisc Surg.2012.05.005>

Current Medications (10 points, 1 point per completed med)
10 different medications must be completed

Hospital Medications (5 required)

Brand/ Generic	Fluconazole (Diflucan)		Magnesium L- lactate (Mag-Tab SR)	Sertraline (Zoloft)	
Dose	400 mg		84 mg	50 mg	
Frequency	100 mL/hr daily		BID	Daily	

N321 CARE PLAN

Route	IV		Oral	Oral	
Classification	Pharmacological class: Azole antifungal Therapeutic class: Antifungal		Pharmacological class: Mineral Therapeutic class: Electrolyte replacement	Pharmacologic al class: Selective serotonin reuptake inhibitor (SSRI) Therapeutic class: Antianxiety, antidepressant, antiobsessant, antipanic, antiposttraumatic stress, antipremenstrual dysphoric	
Mechanism of Action	Damages fungal cells by interfering with a cytochrome P-450 enzyme needed to convert lanosterol to ergosterol, an essential part of the fungal cell membrane.		Assists all enzymes involved in phosphate transfer reactions that use adenosine triphosphate (ATP). Magnesium is required for the normal function of the ATP-dependent sodium-potassium pump in muscle membranes.	Inhibits reuptake of the neurotransmitter serotonin available in nerve synapses. An elevated serotonin level may result in elevated mood and reduced depression. This action may also relieve symptoms of other psychiatric conditions attributed to serotonin deficiency and premenstrual dysmorphic disorder	
Reason Client Taking	Treat esophageal candidiasis		To correct magnesium deficiency	It is used to treat depression	

N321 CARE PLAN

Contraindications (2)	-Coadministration of drugs known to prolong QT interval and which are metabolized via the enzyme CYP3A4 or concurrent therapy with terfenadine -Hypersensitivity to fluconazole or its components		-Hypersensitivity to magnesium salts -Hypersensitivity to any component to any magnesium-containing preparations	-Concurrent use of disulfiram or pimozide -Hypersensitivity to sertraline or its components	
Side Effects/Adverse Reactions (2)	-Prolonged QT interval -Adrenal insufficiency		-Arrhythmias -Hypermagnesemia	-Drowsiness -Ventricular tachycardia	
Nursing Considerations (2)	-Assess for rash every 8 hours during, and notify prescriber if rash occurs -Monitor coagulation test results and assess the patient for bleeding if the patient is receiving an oral anticoagulant		-Be aware that magnesium sulfate is the elemental form of magnesium. Oral preparations aren't all equivalent. -Be aware that the drug isn't metabolized. Drug remaining in the GI tract produces watery stool within 30 minutes to 3 hours.	-Monitor patient for hypo-osmolality of serum and urine and for hyponatremia, which may indicate sertraline-induced syndrome of inappropriate ADH secretion -Be aware that effective antidepressant therapy can promote the development of mania in predisposed people. If mania develops, notify the prescriber immediately	

N321 CARE PLAN

				and expect to withhold sertraline.	
--	--	--	--	------------------------------------	--

Home Medications (5 required)

Brand/Generic	Acetaminophen (Tylenol)	Albuterol sulfate (Accuneb)	Omeprazole (Prilosec)	Pantoprazole (Protonix)	
Dose	325 mg	90 mcg	20 mg	10 mg	
Frequency	PRN	PRN	Daily	PRN	
Route	Oral	Inhalation	Oral	Oral	
Classification	Pharmacologic class: Nonsalicylate, paraaminophenol derivative Therapeutic class: Antipyretic, nonopioid analgesic	- Pharmacologic class: Adrenergic - Therapeutic class: Bronchodilator	- Pharmacologic class: Proton pump inhibitor - Therapeutic class: Antiulcer	- Pharmacologic class: Proton pump inhibitor Therapeutic class: Antiulcer	

N321 CARE PLAN

Mechanism of Action	Inhibits the enzyme cyclooxygenase, blocking prostaglandin production and interfering with pain impulse generation in the peripheral nervous system. Acetaminophen also acts directly on temperature-regulating center in the hypothalamus by inhibiting the synthesis of prostaglandin in E2	Albuterol attaches to beta 2 receptors on bronchial cell membranes, which stimulates the intracellular enzyme adenylate cyclase to convert adenosine triphosphate (ATP) to cyclic adenosine monophosphate (cAMP). This reaction decreases intracellular calcium levels. It also increases intracellular levels of cAMP, as shown. Together, these effects bronchial smooth muscle cells and inhibit histamine release	Omeprazole interferes with gastric acid secretion by inhibiting the hydrogen potassium adenosine triphosphatase enzyme system, or proton pump, in gastric parietal cells.	Interferes with gastric acid secretion by inhibiting the hydrogen-potassium-adenosine triphosphatase ($H^+-K^+-ATPase$) enzyme system, or proton pump, in gastric parietal cells. Normally, the proton pump uses energy from hydrolysis of ATPase to drive H^+ and chloride (Cl^-) out of parietal cells and into the stomach lumen in exchange for potassium (K^+), which leaves the stomach lumen and enters parietal cells. After this exchange,	
----------------------------	--	--	--	--	--

N321 CARE PLAN

				H⁺ and Cl⁻ combine in the stomach to form hydrochloric acid (HCl). Pantoprazole irreversibly inhibits the final step in gastric acid production by blocking the exchange of intracellular H⁺ and extracellular K⁺, thus preventing H⁺ from entering the stomach and additional HCl from forming.	
Reason Client Taking	To relieve mild to moderate pain	To prevent exercise-induced bronchospasm	It is used to treat GERD	It is used to treat GERD	
Contraindications (2)	- Hypersensitivity to acetaminophen or its components - Severe hepatic impairment	- Hypersensitivity to albuterol or its components	- Concurrent therapy with rilpivirine-containing products - Hypersensitivity to	- Concurrent therapy with rilpivirine-containing products - Hypersensitivity to	

N321 CARE PLAN

			omeprazole, substituted benzimidazoles, or their components	pantoprazole, substituted benzimidazoles, or their components	
Side Effects/Adverse Reactions (2)	- Hypotension - Anaphylaxis	-Anxiety -Pharyngitis	-Chest pain - Hypertension	- Hyperglycemia -Bronchitis	
Nursing Considerations (2)	-Use acetaminophen cautiously in patients with hepatic impairment or active hepatic disease, alcoholism, chronic malnutrition, severe hypovolemia, or severe renal impairment -Monitor renal function in patient on long-term therapy. Keep in mind that blood or albumin in urine may indicate nephritis; decreased urine	-Administer pressurized inhalations of albuterol during the second half of inspiration, when airways are open wider and aerosol distribution is more effective -Be aware that drug tolerance can develop with prolonged use.	-Give omeprazole before meals, preferably in the morning for once-daily dosing. If needed, also give an antacid, as prescribed -Know that because drugs can interfere with absorption of vitamin-B12, monitor patient for macrocytic anemia.	-Ensure the continuity of gastric acid suppression during transition from oral to I.V. pantoprazole (or vice versa) because even a brief interruption of effective suppression can lead to serious complications. -Don't give pantoprazole within 4 weeks of testing for <i>Helicobacter pylori</i> because antibiotics, bismuth preparations, and	

N321 CARE PLAN

	output may indicate nephritis; decreased urine output may indicate renal failure; and dark brown urine may indicate the presence of the metabolite phenacetin.			proton pump inhibitors suppress H. pylori and may lead to false-negative results. Drug also may cause false-positive results in urine screening tests for tetrahydrocannabinol. Consult guidelines for pantoprazole use before testing.	
--	--	--	--	---	--

Medications Reference (1) (APA):

Jones & Bartlett Learning, LLC. (2021). 2021 *Nurse's drug handbook* (twentieth).

Assessment

Physical Exam (18 points) – **HIGHLIGHT ALL PERTINENT ABNORMAL FINDINGS**

GENERAL: Alertness: Orientation: Distress: Overall appearance:	The patient is alert and oriented The patient does not seem visibly distressed Pt well dressed in clean gown Pt's skin, hair, nails clean and well maintained
INTEGUMENTARY: Skin color: Character:	Skin color: White Character: Skin is warm and dry upon palpation

N321 CARE PLAN

Temperature: Turgor: Rashes: Bruises: Wounds: Braden Score: Drains present: Y ☐ N ☐ Type:	Temperature: Taken orally and was 97.8 F Turgor: Skin has normal turgor Pt has no visible bruising Normal quantity, distribution, and texture of hair Braden score: 20 The patient has a JP drain
HEENT: Head/Neck: Ears: Eyes: Nose: Teeth:	Head/Neck: Head and neck are symmetrical. Normocephalic and atraumatic Ears: Left/right external ear normal Eyes: No visible drainage from eyes, the bilateral sclera is white, the bilateral cornea is clear, bilateral conjunctiva is pink. Bilateral lids are moist and pink without any discharge Extraocular movements: extraocular movements intact Conjunctiva/sclera: conjunctivae normal Pupils: pupils are equal, round, reactive to light Nose: Septum is midline and no visible bleeding from nose Teeth: Did not notice plaque or tartar. Teeth are white and somewhat aligned with gums. The mucous membrane is moist
CARDIOVASCULAR: Heart sounds: S1, S2, S3, S4, murmur etc. Cardiac rhythm (if applicable): Peripheral Pulses: Capillary refill: Neck Vein Distention: Y ☐ N ☐ Edema Y ☐ N ☐ Location of Edema:	Normal heart rate and rhythm. Clear S1 and S2 without any murmurs Peripheral pulse: 3+ Capillary refill: 2 seconds No edema or neck vein distention
RESPIRATORY: Accessory muscle use: Y ☐ N ☐ Breath Sounds: Location, character	No accessory muscle use Breath sounds: normal breath sounds Effort: pulmonary effort is normal. No respiratory distress
GASTROINTESTINAL: Diet at home: Current Diet Height: Weight: Auscultation Bowel sounds:	Diet at home: regular Current diet: full liquid Height: 5'11" Weight: 137 lbs Bowel sounds: hyperactive Last BM: 3/10/22

N321 CARE PLAN

Last BM: Palpation: Pain, Mass etc.: Inspection: Distention: Incisions: Scars: Drains: Wounds: Ostomy: Y ☐ N ☐ Nasogastric: Y ☐ N ☐ Size: Feeding tubes/PEG tube Y ☐ N ☐ Type:	Upon palpation there is pain and an abdominal mass present. The abdomen is soft Tenderness: There is abdominal tenderness. There is no guarding or rebound Inspection: none Distention: none Incisions: none Scars: none Drains: JP drain Wounds: none No ostomy, nasogastric, or feeding tubes Swelling od abdomen
GENITOURINARY: Color: Character: Quantity of urine: Pain with urination: Y ☐ N ☐ Dialysis: Y ☐ N ☐ Inspection of genitals: Catheter: Y ☐ N ☐ Type: Size:	Clear yellow No pain with urination No dialysis Genitals appear to be normal No catheter
MUSCULOSKELETAL: Neurovascular status: ROM: Supportive devices: Strength: ADL Assistance: Y ☐ N ☐ Fall Risk: Y ☐ N ☐ Fall Score: Activity/Mobility Status: Independent (up ad lib) ☐ Needs assistance with equipment ☐ Needs support to stand and walk ☐	Neurovascular status: normal No swelling. Normal range of motion Cervical back- normal range of motion Strength: Patient noticeably strong Supportive devices: none ADL assistance: none Fall risk: yes Fall risk score: 9 Mobility status: patient able to move freely without equipment. Pt independent. Does not need assistance when standing or walking.
NEUROLOGICAL: MAEW: Y ☐ N ☐ PERLA: Y ☐ N ☐ Strength Equal: Y ☐ N ☐ if no - Legs ☐ Arms ☐ Both ☐ Orientation: Mental Status: Speech: Sensory:	MAEW: no PERLA: no Strength is equal Orientation: pt oriented Mental status: pt is alert and oriented to person, place, and time Speech: normal Sensory: normal sensation Pt awake and alert

N321 CARE PLAN

LOC:	LOC: pt awake and alert
PSYCHOSOCIAL/CULTURAL: Coping method(s): Developmental level: Religion & what it means to pt.: Personal/Family Data (Think about home environment, family structure, and available family support):	Coping method: resting, smartphone, television Developmental level: developmental status appropriate for age Patient is calm and cooperative. The patient is also accepting and participates in care. Behavior is appropriate to the situation. The patient does not state that they are religious. The patient's personal family data is undetermined.

Vital Signs, 2 sets (5 points) – **HIGHLIGHT ALL ABNORMAL VITAL SIGNS**

Time	Pulse	B/P	Resp Rate	Temp	Oxygen
7:52	70	143/70	18	97.8 F	96%
11:41	62	146/78	18	97.7 F	96%

Pain Assessment, 2 sets (2 points)

Time	Scale	Location	Severity	Characteristics	Interventions
7:52	1-10	N/A	N/A	N/A	N/A
11:41	1-10	N/A	N/A	N/A	N/A

IV Assessment (2 Points)

IV Assessment	Fluid Type/Rate or Saline Lock
Size of IV: Location of IV: Date on IV: Patency of IV:	Size of IV: 20 Location of IV: Right lower arm Date on IV: 3/1/22 Patency of IV: open

N321 CARE PLAN

Signs of erythema, drainage, etc.: IV dressing assessment:	Signs of erythema, drainage, etc.: no signs IV dressing assessment: dry and intact
---	---

Intake and Output (2 points)

Intake (in mL)	Output (in mL)
800- fluids	800 mL
200- Fluconazole	
2000- clinimix-e-parenteral nutrition mixture	

Nursing Care**Summary of Care (2 points)**

Overview of care: Patient is diagnosed with a mesenteric tumor. Labs are being done daily to monitor how effective treatment is.

Procedures/testing done: CT scan of abdomen/pelvis

Complaints/Issues: No complaints/issues

Vital signs (stable/unstable): Vital signs are stable

Tolerating diet, activity, etc.: Patient tolerating diet and activity

Physician notifications: None

Future plans for client: Undetermined

Discharge Planning (2 points)

Discharge location: Patients home

Home health needs (if applicable): The patient has a contract with a home care agency. The patient needs a home TPN

Equipment needs (if applicable): Not applicable

N321 CARE PLAN

Follow-up plan: No surgical intervention planned. The patient will need to follow-up with outpatient

Education needs: The patient will be educated on how to care for a JP drain, and how often the patient needs to empty it

Nursing Diagnosis (15 points)

Must be NANDA approved nursing diagnosis and listed in order of priority

Nursing Diagnosis	Rationale	Interventions (2 per dx)	Outcome Goal (1 per dx)	Evaluation
● Include full nursing diagnosis with “related to” and “as evidenced by” components ● Listed in order by priority – highest priority to lowest priority pertinent to this client	● Explain why the nursing diagnosis was chosen			● How did the client/family respond to the nurse’s actions? ● Client response, status of goals and outcomes, modifications to plan.
1. Risk for falls related to a hemoglobin of 7.6 as evidenced by a mesenteric tumor	After admission, the patient’s lab indicated a hemoglobin level of 7.6 which indicates anemia	1. It is important to be aware of the factors that may cause injury from a fall 2. It is important to constantly assess the patient's fall risk. Before the	1. Increase the amount of iron intake. Continue to monitor hemoglobin levels on a daily basis.	The client was cooperative when teaching how to increase hemoglobin levels. The patient agreed to eat foods with more iron.

N321 CARE PLAN

		patient gets up make sure to remove objects from the ground		
2. Risk for fluid volume deficit related to inadequate fluid intake as evidenced by low levels of sodium .	After admission, the patient's lab indicated a sodium level of 134	1. Assess the client's skin turgor and mucous membranes for signs of dehydration. 2. Assess the volume and frequency of vomiting.	1. Raise the levels of sodium and continue to monitor intakes of fluid and vomiting frequency.	The patient was willing to understand their abnormal labs and continue to monitor their labs on a daily basis. Patients will help achieve their goal by being cooperative and attempting to consume more liquids.
3. Risk for diarrhea related to an abscess in the mesentery as evidenced by a mesenteric tumor	The patient was admitted to the emergency department with abdominal pain which indicated there is a likelihood of getting diarrhea	1. Assess type and frequency of bowel movement 2. Assess the patient's fluid intake	1. The patient was given IV fluids in a timely manner. Give patient foods with more fiber	Patient was very understanding of the interventions and agreed to consume foods high in fiber and make sure they drink more fluids.

Other References (APA):

N321 CARE PLAN

Phelps, L. L. (2020). *Sparks & Taylor's nursing diagnosis reference manual*. Wolters Kluwer.

Concept Map (20 Points):

Objective data:

BP: 143/70
Pulse: 70
Oxygen: 96%
Pulse level: N/A
Temp: 97.8 F
Resp: 18
Mesenteric tumor found on CT scan

Patient information:

M.Q.
66 year-old
Asian
Not employed
Single
Allergies: Plasma Human
Full Code
5'1"
137 lbs

Risk for fall:

Risk for falls related to patient hemoglobin of 7.0 as evidenced by a mesenteric tumor
Risk for fluid volume deficit related to inadequate fluid intake as evidenced by low levels of sodium
Risk for diarrhea related to an assess in the mesentery as evidenced by a mesenteric tumor

Risk for fluid volume deficit:

1. Assess the client's skin turgor and mucous membranes for signs of dehydration.

Nursing outcomes/goals: the volume and frequency of vomiting and vomiting frequency.
1. Increase the amount of fluid intake. Continue to monitor hemoglobin levels on a daily basis.
2. Raise the levels of sodium and continue to monitor intake of fluid and vomiting frequency.
3. The patient was given IV fluids in a timely manner. Give patient foods with more fiber

Risk for diarrhea:

1. Assess type and frequency of bowel movement
2. Assess the patient's fluid intake

N321 CARE PLAN

N321 CARE PLAN