

N321 Care Plan #1

Lakeview College of Nursing

Jamal Drea

## N321 CARE PLAN

**Demographics (3 points)**

<b>Date of Admission</b> 5/15/22	<b>Client Initials</b> J.M.	<b>Age</b> 84	<b>Gender</b> Male
<b>Race/Ethnicity</b> White or Caucasian	<b>Occupation</b> Not employed	<b>Marital Status</b> Widowed	<b>Allergies</b> Morphine
<b>Code Status</b> FULL	<b>Height</b> 5' 11.5"	<b>Weight</b> 170	

**Medical History (5 Points)**

**Past Medical History:** Atrial fibrillation, acute renal failure (2009), rectal cancer, stenosis of rectum and anus, colostomy (2013), HTN, hearing loss (2013), hyperlipidemia, thrombocytopenia, cataract

**Past Surgical History:** polypectomy, retinal detachment surgery (right), cataract removal with implant, colonoscopy stoma, capsulotomy laser (right - 2019)

**Family History:** N/A

**Social History (tobacco/alcohol/drugs including frequency, quantity and duration of use):**  
Former tobacco smoker of 35 years (stopped 30 years ago), used to drink 6 cans of beer a week, no history of drug use

**Assistive Devices:** Hearing aid, walker, reading glasses

**Living Situation:** Lives alone

**Education Level:** Adult

**Admission Assessment**

**Chief Complaint (2 points):** Confusion

**History of Present Illness – OLD CARTS (10 points):**

**Onset:** Confusion occurred 1 week ago, but patient was found negative for stroke

**Location:** Brain

N321 CARE PLAN

**Duration: Intermittent**

**Characteristic Symptoms: N/A during time of assessment**

**Aggravating Factors: N/A**

**Associated Manifestations: Impaired memory, weakness and unequal strength in extremities**

**Relieving Factors: N/A**

**Treatments: No previous treatment for confusion**

### **Primary Diagnosis**

**Primary Diagnosis on Admission (2 points): Acute metabolic encephalopathy**

**Secondary Diagnosis (if applicable):**

**Pathophysiology of the Disease, APA format (20 points):**

**An encephalopathy is a condition of the brain in which function is impaired as a result of several possible factors that can cause a patient to be confused, experience muscle weakness, lose memory, or have other symptoms that are neuropathic in origin. These factors include and could also be a combination of abnormal blood flow, abnormal energy metabolism, abnormal neurotransmission, and cellular maintenance processes that may contribute to cerebral metabolic insufficiency (Williams, 2013). Since metabolism has a significant influence on energy and cellular processes, problems with liver function could be the cause of the disorder that leads patients to be in an altered mental state. Hepatic encephalopathy happens due to liver failure and is commonly linked to cirrhosis that is associated with alcohol abuse and inflammation. Ammonia is a nitrogenous waste and toxin that can build up if the liver is unable to remove it from the body, which could affect the**

**N321 CARE PLAN**

health of the brain and lead to symptoms like confusion, disorientation, and even coma (Capriotti & Frizzell, 2020). Another example would be pulmonary encephalopathy in which the lungs function insufficiently as a result of infection or congestive heart failure. In this case, carbon dioxide is not exchanged and arterial pressure rises to levels of hypercapnia that exerts onto cerebrospinal fluid and changes level of consciousness with impaired neuronal metabolism (Angel & Young, 2011). An imbalance in electrolyte levels is an additional instance that could cause metabolic encephalopathy. In a situation where a patient has a significantly low level of sodium, or hyponatremia, it could alter the osmotic pressure that is responsible for the movement of water through the intracellular space and this may bring about cerebral edema that impairs brain function with symptoms like headaches, respiratory failure, and seizures (Angel & Young, 2011). There are multiple ways that can make a patient suffer this condition but they are similarly rooted in metabolism and homeostasis. It is important to be mindful of the vitals and lab diagnostics that could indicate an issue that would help prevent damage by this condition.

**Pathophysiology References (2) (APA):**

Angel, M. J., & Young, G. B. (2011). Metabolic encephalopathies. *Neurologic clinics*, 29(4), 837–882. <https://doi.org/10.1016/j.ncl.2011.08.002>

Capriotti, T. & Frizzell, J.P. (2020). *Pathophysiology: Introductory concepts and clinical perspectives*. (2nd ed.). F.A. Davis Company.

Williams S. T. (2013). Pathophysiology of encephalopathy and delirium. *Journal of clinical neurophysiology : official publication of the American Electroencephalographic Society*, 30(5), 435–437. <https://doi.org/10.1097/WNP.0b013e3182a73e04>

**Laboratory Data (15 points)**

## N321 CARE PLAN

**CBC Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason for Abnormal Value
RBC	4.7-6.1	3.65	3.68	Hemolytic medications that the patient takes could lower RBC. The low value for red blood count could also be attributed to overhydration as the patient was put on a liquid intake restriction of 1 L by the provider (Pagana, 2018). A urinalysis shows hematuria so blood loss could be a factor.
Hgb	14-18	10.8	11.4	A lower than expected HgB is related to a reduced RBC count (Pagana, 2018).
Hct	42-52%	30.9	32.6	Hematocrit reflects the HgB and RBC count which could explain a less than normal range (Pagana, 2018).
Platelets	150-400	109	131	A urinalysis of the patient shows an infection, which could be the reason for a reduced platelet count. The patient may have a thrombocytopenia disorder since this is shown in their medical history (Pagana, 2018).
WBC	5-10	4.9	4.58	WBC count is slightly reduced and could be caused by impaired bone marrow function since RBC is also lower than the expected value (Pagana, 2018).
Neutrophils	55-70%	52	51	
Lymphocytes	20-40%	26.9	31	
Monocytes	2-8%	16.5	13.1	
Eosinophils	1-4%	3.1	0.7	
Bands	0.5-1%	N/A	N/A	

## N321 CARE PLAN

Chemistry **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason For Abnormal
Na-	136-145	125	129	Hyponatremia could be caused by renal insufficiency. The patient has a history of renal complications that could be the reason for this low level of sodium (Pagana, 2018). Patient is currently treated with sodium tablets.
K+	3.5-5.0	4.1	4.7	
Cl-	98-106	90	94	Hypochloremia could be caused by diuretic therapy and the patient has been prescribed furosemide (Pagana, 2018).
CO2	23-30	22	25	Slightly low CO2 levels could be due to the patient's renal insufficiency (Pagana, 2018).
Glucose	82-115	100	90	
BUN	10-20	12	28	
Creatinine	0.6-1.2	0.94	1.18	
Albumin	3.5-5.0	3.5	N/A	
Calcium	9.0-10.5	8.3	8.5	Low calcium could be the result of renal insufficiency but the patient is also of elderly age and may suffer from malabsorption of vitamin C. It is not likely that it is by dietary causes since the patient enjoys milk and ice cream during meals (Pagana, 2018).
Mag	1.3-2.1	1.3	1.8	
Phosphate	3.0-4.5	3.5	N/A	
Bilirubin	0.3-1.0	1.2	N/A	Bilirubin is elevated and this could be caused by salicylate

## N321 CARE PLAN

				medications. The patient takes aspirin as a home medication (Pagana, 2018).
Alk Phos	30-120	78	N/A	
AST	0-35	16	N/A	
ALT	4-36	11	N/A	
Amylase	6.6-35.2	N/A	N/A	
Lipase	0-160	N/A	N/A	
Lactic Acid	5-20	N/A	N/A	

Other Tests **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab Test	Normal Range	Value on Admission	Today's Value	Reason for Abnormal
INR	0.8-1.1	N/A	1.0	
PT	11-12.5	N/A	12.3	
PTT	30-40	N/A	27.3	PTT is slightly low which can be supported by overactive blood clotting proteins (Pagana, 2018).
D-Dimer	< 250	N/A	865	Anemia shown by RBC could have increased the value of D-Dimer (Pagana, 2018).
BNP	< 100	N/A	147	BNP levels are usually higher in older adults (Pagana, 2018).
HDL	> 45	N/A	50	
LDL	<130	N/A	87	
Cholesterol	< 200	N/A	149	
Triglycerides	40-160	N/A	61	

## N321 CARE PLAN

Hgb A1c	4-5.9 in nondiabetic	N/A	5.1	
TSH	2-10	N/A	N/A	

Urinalysis **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab Test	Normal Range	Value on Admission	Today's Value	Reason for Abnormal
Color & Clarity	Amber-Light Yellow and clear	Light yellow and clear	N/A	
pH	4.6-8.0	6	N/A	
Specific Gravity	1.005-1.030	1.011	N/A	
Glucose	Negative	Negative	N/A	
Protein	0-8	Negative	N/A	
Ketones	Negative	15	N/A	The patient has no record of diabetes in their health history so a diet that lacked carbohydrates before admission could have resulted in ketones in the urine sample (Pagana, 2018).
WBC	0-4	Negative	N/A	
RBC	<2	Moderate	N/A	Overaggressive anticoagulant therapy or bleeding disorders may have caused RBCs in the urine (Pagana, 2018).
Leukoesterase	Negative	Negative	N/A	

Cultures **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Test	Normal Range	Value on Admission	Today's Value	Explanation of Findings
Urine Culture	Negative < 100,000	>100,000 gram	N/A	Klebsiella oxytoca was found in the urine culture.

## N321 CARE PLAN

		<b>negative bacilli</b>		
<b>Blood Culture</b>	<b>Negative</b>	<b>Negative</b>	<b>N/A</b>	
<b>Sputum Culture</b>	<b>Negative</b>	<b>Negative</b>	<b>N/A</b>	
<b>Stool Culture</b>	<b>Negative</b>	<b>Negative</b>	<b>N/A</b>	

**Lab Correlations Reference (1) (APA):**

**Pagana, K.D., Pagana, T.J., & Pagana, T.N. (2018). *Mosby's Diagnostic and Laboratory Test Reference* (14th ed.). Mosby.**

**Diagnostic Imaging**

**All Other Diagnostic Tests (5 points): CT Chest w/o contrast shows pleural thickening, stomach lining thickening, coronary artery disease, trace pericardial effusion, mild mediastinal lymphadenopathy, subpleural nodularity in right upper lobe.**

**Diagnostic Test Correlation (5 points): CT of chest is a noninvasive x-ray procedure for diagnosing and evaluating pathologic conditions such as tumors, nodules, hematomas, parenchymal lesions, cysts, abscesses, pleural effusion, and enlarged lymph nodes (Pagana, 2018).**

**Diagnostic Test Reference (1) (APA):**

**Pagana, K.D., Pagana, T.J., & Pagana, T.N. (2018). *Mosby's Diagnostic and Laboratory Test Reference* (14th ed.). Mosby.**

**Current Medications (10 points, 1 point per completed med)  
\*10 different medications must be completed\***

## N321 CARE PLAN

**Home Medications (5 required)**

<b>Brand/Generic</b>	<b>Lipitor (Atorvastatin)</b>	<b>Flomax (Tamsulosin)</b>	<b>Lopressor (Metoprolol tartrate)</b>	<b>Prilosec (Omeprazole magnesium)</b>	<b>Spiriva respiant (Tiotropium bromide)</b>
<b>Dose</b>	<b>20 mg</b>	<b>0.4 mg</b>	<b>75 mg</b>	<b>40 mg</b>	<b>2.5 mcg</b>
<b>Frequency</b>	<b>Daily with evening meal</b>	<b>Daily</b>	<b>BID</b>	<b>Daily</b>	<b>Daily</b>
<b>Route</b>	<b>oral</b>	<b>oral</b>	<b>oral</b>	<b>oral</b>	<b>inhalation</b>
<b>Classification</b>	Pharmacologic class: HMG-CoA reductase inhibitor  Therapeutic class: Antihyperlipidemic	Pharmacologic class: Alpha adrenergic antagonist  Therapeutic class: Benign prostatic hyperplasia (BPH) agent	Pharmacologic class: Beta1-adrenergic blocker  Therapeutic class: Antianginal, antihypertensive	Pharmacologic class: Proton pump inhibitor  Therapeutic class: Antiulcer	Pharmacologic class: Anticholinergic  Therapeutic class: Bronchodilator
<b>Mechanism of Action</b>	Reduces plasma cholesterol and lipoprotein levels by inhibiting HMG-CoA reductase and cholesterol synthesis in the liver and by increasing the number of LDL receptors on liver cells to enhance LDL uptake and breakdown.	Blocks alpha1-adrenergic receptors in the prostate. This action inhibits smooth muscle contraction in the bladder neck and prostate, prostatic capsule, and prostatic urethra, which improves the rate of urine flow and reduces symptoms of BPH.	Inhibits stimulation of beta1-receptor sites, located mainly in the heart, resulting in decreased cardiac excitability, cardiac output, and myocardial oxygen demand. These effects help relieve angina, minimize cardiac tissue damage from a myocardial infarction, and help relieve symptoms of heart failure. Metoprolol	Omeprazole interferes with gastric acid secretion by inhibiting the hydrogen potassium adenosine triphosphatase (H <sup>+</sup> K <sup>+</sup> -ATPase) enzyme system, or proton pump, in gastric parietal cells. Normally, the proton pump uses energy from hydrolysis of adenosine triphosphate to drive hydrogen (H <sup>+</sup> ) and chloride	Prevents acetylcholine from attaching to muscarinic receptors on membranes of smooth-muscle cells. By blocking acetylcholine's effects in the bronchi and bronchioles, tiotropium relaxes smooth muscles and causes bronchodilation.

## N321 CARE PLAN

			also helps reduce blood pressure by decreasing renal release of renin.	(Cl <sup>-</sup> ) out of parietal cells and into the stomach lumen in exchange for potassium (K <sup>+</sup> ), which leaves the stomach lumen and enters parietal cells. After this exchange, H <sup>+</sup> and Cl <sup>-</sup> combine in the stomach to form hydrochloric acid (HCl), as shown below left. Omeprazole irreversibly blocks the exchange of intracellular H <sup>+</sup> and extracellular K <sup>+</sup> , as shown below right. By preventing H <sup>+</sup> from entering the stomach lumen, omeprazole keeps additional HCl from forming.	
<b>Reason Client Taking</b>	To control lipid levels and disorders such as hypercholesterolemia and dyslipidemia. The client has a past medical history with hyperlipidemia.	To treat benign prostatic hyperplasia	To treat acute MI early and treat hypertension	To provide active treatment of active benign gastric ulcer or maintain GERD	To provide maintenance treatment of asthma
<b>Contraindications (2)</b>	Active hepatic disease and persistent rise in serum	Sensitivity to tamsulosin and quinazolines	Hypersensitivity to beta blockers and heart blocks	Concurrent therapy with rilpivirine-containing products;	Hypersensitivity to atropine and its derivatives

## N321 CARE PLAN

	transaminase level		greater than 1sr degree	hypersensitivity to omeprazole or substituted benzimidazoles	such as ipratropium and tiotropium
<b>Side Effects/Adverse Reactions (2)</b>	Arrhythmias and hypoglycemia	Arrhythmia and respiratory impairment	Cardiac arrest and leukopenia	Hypoglycemia and hepatic dysfunction or failure	Atrial fibrillation and intestinal obstruction
<b>Nursing Considerations (2)</b>		Flomax should be given 30 minutes after the same meal every day. If the patient takes the medication on an empty stomach then their blood pressure should be monitored due to risk for orthostatic hypotension.	Use cautiously in patients with angina or hypertension who have congestive heart failure because beta blockers such as metoprolol can further depress myocardial contractility, worsening heart failure. Before starting therapy for heart failure, expect to give an ACE inhibitor, digoxin, and a diuretic to stabilize the patient.	Give omeprazole before meals, preferably in the morning for once-daily dosing. If needed, also give an antacid, as prescribed. Monitor the patient for macrocytic anemia because the drug can interfere with the absorption of vitamin K.	This drug should not be used for acute bronchospasms, it is strictly for maintenance. You should monitor the patient's pulmonary function to evaluate effectiveness of treatment.

## N321 CARE PLAN

## Hospital Medications (5 required)

<b>Brand/Generic</b>	<b>Plavix (Clopidogrel bisulfate)</b>	<b>Lovenox (Enoxaparin)</b>	<b>Lasix (Furosemide)</b>	<b>Seroquel (Quetiapine fumarate)</b>	<b>Zofran (Ondansetron)</b>
<b>Dose</b>	<b>75 mg</b>	<b>40 mg</b>	<b>20 mg</b>	<b>12.5 mg</b>	<b>4 mg</b>
<b>Frequency</b>	<b>Daily</b>	<b>Daily</b>	<b>BID</b>	<b>Daily at bedtime</b>	<b>Daily PRN</b>
<b>Route</b>	<b>Oral</b>	<b>Subcutaneous</b>	<b>Oral</b>	<b>Oral</b>	<b>IV</b>
<b>Classification</b>	Pharmacologic class: P2Y <sub>12</sub> platelet inhibitor  Therapeutic class: Platelet aggregation inhibitor	Pharmacologic class: Low-molecular-weight heparin  Therapeutic class: Anticoagulant	Pharmacologic class: Loop diuretic  Therapeutic class: Antihypertensive, diuretic	Pharmacologic class: Dibenzothiazepine derivative  Therapeutic class: Antipsychotic	Pharmacologic class: Selective serotonin (5-HT <sub>3</sub> ) receptor antagonist  Therapeutic class: Antiemetic
<b>Mechanism of Action</b>	Binds to adenosine diphosphate (ADP) receptors on the surface of activated platelets. This action blocks ADP, which deactivates nearby glycoprotein IIb/IIIa receptors and prevents fibrinogen from attaching to receptors. Without fibrinogen, platelets can't aggregate and form thrombi.	Potentiates the action of antithrombin III, a coagulation inhibitor. By binding with antithrombin III, enoxaparin rapidly binds with and inactivates clotting factors (primarily factor Xa and thrombin). Without thrombin, fibrinogen can't convert to fibrin and clots can't form.	Inhibits sodium and water reabsorption in the loop of Henle and increases urine formation. As the body's plasma volume decreases, aldosterone production increases, which promotes sodium reabsorption and the loss of potassium and hydrogen ions. Furosemide also increases the excretion of calcium, magnesium, bicarbonate, ammonium, and phosphate. By	May produce antipsychotic effects by interfering with dopamine binding to dopamine type 2 (D <sub>2</sub> )-receptor sites in the brain and by antagonizing serotonin 5-HT <sub>2</sub> , dopamine type 1 (D <sub>1</sub> ), histamine H <sub>1</sub> , and adrenergic alpha <sub>1</sub> and alpha <sub>2</sub> receptors.	Blocks serotonin receptors centrally in the chemoreceptor or trigger zone and peripherally at vagal nerve terminals in the intestine. This action reduces nausea and vomiting by preventing serotonin release in the small intestine and by blocking signals to the CNS.

## N321 CARE PLAN

			reducing intracellular and extracellular fluid volume, the drug reduces blood pressure and decreases cardiac output. Over time, cardiac output returns to normal.		
<b>Reason Client Taking</b>	To reduce thrombotic events, such as MI and stroke, in patients with atherosclerosis documented (in this case by arterial disease as evidenced by CT of chest)	To prevent DVT	To reduce edema and hypertension	To treat disorders with behavior	To prevent nausea and vomiting
<b>Contraindications (2)</b>	Active pathological bleeding and hypersensitivity to clopidogrel	Active major bleeding and history of immune-mediated heparin-induced thrombocytopenia	Anuria and hypersensitivity to furosemide	Hypersensitivity to quetiapine or components of quetiapine	Use of apomorphine or hypersensitivity to ondansetron
<b>Side Effects/Adverse Reactions (2)</b>	Hypotension and bronchospasm	CHF and Hematemesis	Arrhythmia and thromboembolism	Cardiomyopathy or thrombocytopenia	Hypotension and arrhythmias or chest pain
<b>Nursing Considerations (2)</b>	Use with caution for patients with hepatic or renal disease and those at risk for bleeding from trauma or surgery. Obtain blood cell count, as ordered, whenever	Use with caution for patients with increased risk of hemorrhage. This medication should not be used in patients with prosthetic heart valves.	Use furosemide cautiously in patients with advanced hepatic cirrhosis, especially those who also have a history of electrolyte imbalance or hepatic encephalopathy	Know that quetiapine shouldn't be used for elderly patients with dementia-related psychosis because the drug increases the risk of death in these patients. Know that this drug	Electrolyte imbalances should be corrected before administration of ondansetron. Monitor the patient for arrhythmia or CHF.

N321 CARE PLAN

	signs and symptoms suggest a hematologic problem.		y. Obtain the patient's weight before and during administration to evaluate fluid loss.	should not be administered to patients who have a history of cardiac arrhythmias. Monitor for dehydration.	
--	---	--	---	--	--

Medications Reference (1) (APA):

Jones & Bartlett Learning. (2020). 2021 Nurse's Drug Handbook (19 th ed.). Jones & Bartlett Learning

Assessment

Physical Exam (18 points) – **HIGHLIGHT ALL PERTINENT ABNORMAL FINDINGS**

<b>GENERAL:</b> <b>Alertness:</b> <b>Orientation:</b> <b>Distress:</b> <b>Overall appearance:</b>	<b>Alert x 3</b> <b>Oriented x 3</b> <b>No acute distress</b> <b>Calm and normal appearance</b>
<b>INTEGUMENTARY:</b> <b>Skin color:</b> <b>Character:</b> <b>Temperature:</b> <b>Turgor:</b> <b>Rashes:</b> <b>Bruises:</b> <b>Wounds:</b> <b>Braden Score:</b> <b>Drains present: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Type:</b>	<b>Normal</b> <b>Normal, dry</b> <b>Cold on palpation</b> <b>Normal, no tenting</b> <b>None</b> <b>None</b> <b>Wound present on left elbow</b> <b>Braden Score: 20</b> <b>No observable drainage present</b>
<b>HEENT:</b> <b>Head/Neck:</b> <b>Ears:</b> <b>Eyes:</b> <b>Nose:</b> <b>Teeth:</b>	<b>Normocephalic, atraumatic/no deviation</b> <b>External ears normal</b> <b>Sclera and conjunctiva normal, PERRL, EOMs intact</b> <b>Nose appears normal with no congestion or rhinorrhea</b> <b>Has upper dentures in good condition</b>

## N321 CARE PLAN

	<b>Mucosa of the mouth and oropharynx is pink and moist</b>
<b>CARDIOVASCULAR:</b> <b>Heart sounds:</b> <b>S1, S2, S3, S4, murmur etc.</b> <b>Cardiac rhythm (if applicable):</b> <b>Peripheral Pulses:</b> <b>Capillary refill:</b> <b>Neck Vein Distention: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Edema Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Location of Edema:</b>	<b>Normal S1 and S2 heart sounds</b> <b>Normal heart rate and rhythm</b> <b>Peripheral pulses 2+</b> <b>Capillary refill &lt;3 seconds</b> <b>No neck vein distention</b> <b>No edema present</b>
<b>RESPIRATORY:</b> <b>Accessory muscle use: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Breath Sounds: Location, character</b>	<b>Pulmonary effort is normal, no accessory muscle use</b> <b>No respiratory distress</b> <b>Breathing sounds normal bilaterally, no wheezing or rales</b>
<b>GASTROINTESTINAL:</b> <b>Diet at home:</b> <b>Current Diet</b> <b>Height:</b> <b>Weight:</b> <b>Auscultation Bowel sounds:</b> <b>Last BM:</b> <b>Palpation: Pain, Mass etc.:</b> <b>Inspection:</b> <b>Distention:</b> <b>Incisions:</b> <b>Scars:</b> <b>Drains:</b> <b>Wounds:</b> <b>Ostomy: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Nasogastric: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Size:</b> <b>Feeding tubes/PEG tube Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Type:</b>	<b>Normal diet, no restrictions</b> <b>Current diet is soft bite and thin liquid</b> <b>5'11.5"</b> <b>170 lbs (23.1 BMI)</b>  <b>Last BM: 1115</b> <b>Bowel sounds are normoactive in all quadrants</b> <b>No distension or reported pain in abdomen</b> <b>No tenderness or guarding</b> <b>No incisions around abdominal region</b> <b>No scars around abdominal region</b> <b>No drainage around abdominal region</b> <b>No wounds around abdominal region</b> <b>Patient has a colostomy</b> <b>No nasogastric or feeding tubes</b>
<b>GENITOURINARY:</b> <b>Color:</b> <b>Character:</b> <b>Quantity of urine:</b> <b>Pain with urination: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Dialysis: Y <input type="checkbox"/> N <input type="checkbox"/></b> <b>Inspection of genitals:</b> <b>Catheter: Y <input type="checkbox"/> N <input type="checkbox"/></b>	<b>Light yellow, clear urine</b> <b>450 mL output</b> <b>No reported pain with urination</b> <b>No dysuria, urgency, frequency; hematuria present according to urinalysis</b> <b>No dialysis</b> <b>No catheter</b>

## N321 CARE PLAN

Type: Size:	
<b>MUSCULOSKELETAL:</b> Neurovascular status: ROM: Supportive devices: Strength: ADL Assistance: Y <input type="checkbox"/> N <input type="checkbox"/> Fall Risk: Y <input type="checkbox"/> N <input type="checkbox"/> Fall Score: Activity/Mobility Status: Independent (up ad lib) <input type="checkbox"/> Needs assistance with equipment <input type="checkbox"/> Needs support to stand and walk <input type="checkbox"/>	Alert and responsive Normal ROM Walker Strength unequal in legs during neuro assessment, plantar flexion and dorsiflexion especially weak in left leg ADL assistance required for mobility and food preparation Fall Risk: 15 Requires support with standing and walking
<b>NEUROLOGICAL:</b> MAEW: Y <input type="checkbox"/> N <input type="checkbox"/> PERLA: Y <input type="checkbox"/> N <input type="checkbox"/> Strength Equal: Y <input type="checkbox"/> N <input type="checkbox"/> if no - Legs <input type="checkbox"/> Arms <input type="checkbox"/> Both <input type="checkbox"/> Orientation: Mental Status: Speech: Sensory: LOC:	MAEW ✓ PERLA ✓ Strength unequal in legs (left noticeably weaker) Oriented x 3 Normal during assessment Normal Normal Alert and responsive
<b>PSYCHOSOCIAL/CULTURAL:</b> Coping method(s): Developmental level: Religion & what it means to pt.: Personal/Family Data (Think about home environment, family structure, and available family support):	Watches TV and sleeps throughout the day Developmental level: Adult N/A Is widowed but has daughter that calls in to receive updates on condition of the patient

## Vital Signs, 2 sets (5 points) – HIGHLIGHT ALL ABNORMAL VITAL SIGNS

Time	Pulse	B/P	Resp Rate	Temp	Oxygen
0810	76	112/50	12	98.3	95
1100	72	110/52	12	97.7	92

## N321 CARE PLAN

**Pain Assessment, 2 sets (2 points)**

<b>Time</b>	<b>Scale</b>	<b>Location</b>	<b>Severity</b>	<b>Characteristics</b>	<b>Interventions</b>
<b>0810</b>	<b>0</b>	<b>N/A</b>	<b>N/A</b>	<b>N/A</b>	<b>N/A</b>
<b>1100</b>	<b>0</b>	<b>N/A</b>	<b>N/A</b>	<b>N/A</b>	<b>N/A</b>

**IV Assessment (2 Points)**

<b>IV Assessment</b>	<b>Fluid Type/Rate or Saline Lock</b>
<b>Size of IV:</b> <b>Location of IV:</b> <b>Date on IV:</b> <b>Patency of IV:</b> <b>Signs of erythema, drainage, etc.:</b> <b>IV dressing assessment:</b>	20 G Left antecubital 5/15/22 Open and functional, no swelling No signs of erythema or drainage IV is secure

**Intake and Output (2 points)**

<b>Intake (in mL)</b>	<b>Output (in mL)</b>
<b>980</b>	<b>450</b>

**Nursing Care****Summary of Care (2 points)**

**Overview of care: Plan is for long term care**

**Procedures/testing done: CBC, CMP, CT of chest**

**Complaints/Issues: CC is confusion, needs assistance with mobility**

**Vital signs (stable/unstable): Stable**

## N321 CARE PLAN

**Tolerating diet, activity, etc.: Patient tolerates diet, activity is limited**

**Physician notifications: N/A**

**Future plans for client: Physical therapy in long term care facility**

**Discharge Planning (2 points)**

**Discharge location: Extended care facility**

**Home health needs (if applicable): Arrangements for reduced fall risk**

**Equipment needs (if applicable): Walker for assistance with mobility**

**Follow up plan: Follow up on diagnostic tests of chest due to presence of nodule in lungs**

**Education needs: Mobility, fall risk, medications, diagnosis**

**Nursing Diagnosis (15 points)**

**\*Must be NANDA approved nursing diagnosis and listed in order of priority\***

<p><b>Nursing Diagnosis</b></p> <ul style="list-style-type: none"> <li>● Include full nursing diagnosis with “related to” and “as evidenced by” components</li> <li>● Listed in order by priority – highest priority to lowest priority pertinent to this client</li> </ul>	<p><b>Rationale</b></p> <ul style="list-style-type: none"> <li>● Explain why the nursing diagnosis was chosen</li> </ul>	<p><b>Interventions (2 per dx)</b></p>	<p><b>Outcome Goal (1 per dx)</b></p>	<p><b>Evaluation</b></p> <ul style="list-style-type: none"> <li>● How did the client/family respond to the nurse’s actions?</li> <li>● Client response, status of goals and outcomes, modifications to plan.</li> </ul>
<p><b>1. Risk of fall related to acute</b></p>	<p><b>The patient is at risk for fall as a result of their condition,</b></p>	<p><b>1. Remove anything from the environment that will increase the</b></p>	<p><b>1. The patient will assist in identifying and applying physical changes in the</b></p>	<p><b>The patient will optimize activities of daily living</b></p>

## N321 CARE PLAN

<p>metabolic encephalopathy as evidenced by need for assisted mobility and unequal strength in extremities.</p>	<p>which has left them with weakness and limited energy for physical activity. The high fall risk score indicates that it is important that safety measures are put in place to protect the client from injury.</p>	<p>risk of falls. Patient's immediate environment must be reviewed frequently to prevent unnecessary falls.</p> <p>2. Identify and educate on the factors that may cause or contribute to injury from a fall in order to enhance the patient, family, and caregiver awareness of the risks.</p>	<p>environment to ensure increased safety.</p>	<p>within limitations. The patient and their family will understand the factors that could contribute to fall risk.</p>
<p>2. Risk of acute confusion related to acute metabolic encephalopathy as evidenced by intermittent disorientation and memory loss.</p>	<p>The patient is at risk for acute confusion, which has been shown by intermittent instances of confusion and disorientation that affect the patient's cognitive ability and puts them at risk of injury.</p>	<p>1. Monitor neurological status every 4 hours to determine if the client is alert and oriented or experiencing a decline in brain function.</p> <p>2. Use appropriate safety measures to protect the patient from injury.</p>	<p>1. Patient's neurological status will remain stable and they will participate in ADLs.</p>	<p>The patient will understand safety measures that need to be put in place to protect them and family will know when to report change in behavior.</p>
<p>3. Risk for electrolyte imbalance related to acute metabolic encephalopathy as evidenced by hyponatremia and hypocalcemia.</p>	<p>Metabolic encephalopathies can be caused by electrolyte imbalances so it is important to educate the patient about how it relates to their condition so that they could avoid further effects.</p>	<p>1. Assess the patient's fluid status and monitor for physical signs of electrolyte imbalance (cardiac, neuro, musculoskeletal).</p> <p>2. Educate the patient and their family members on the risk of electrolyte imbalances along with their medical condition and</p>	<p>1. The patient will maintain normal values for their electrolyte levels.</p>	<p>The patient will be able to verbalize risks and symptoms that require interventions. The patient and family will understand the goal of the teachings.</p>

## N321 CARE PLAN

		symptoms to watch for in the future.		
--	--	--------------------------------------	--	--

**Other References (APA):**

**Phelps, L.L. (2020). *Sparks and Taylor's Nursing Diagnosis Reference Manual* (11 th ed.).**

**Wolters Kluwer.**

N321 CARE PLAN

Concept Map (20 Points):







