

N311 Care Plan # 3

Lakeview College of Nursing

Jayda Davis

Demographics (5 points)

Date of Admission 9/8/2022	Client Initials B. R	Age 54	Gender Male
Race/Ethnicity Caucasian	Occupation Factory worker	Marital Status Single	Allergies None
Code Status Full Code	Height 69 inches	Weight 89.4 kg	

Medical History (5 Points)

Past Medical History: Unspecified protein-calorie malnutrition, gout, GERD, right foot drop, hypokalemia, hypertension, alcohol dependence, nutritional anemia, hyponatremia, hypo-osmolarity, depression, and history of falling.

Past Surgical History: Patient-reported no surgical history

Family History: Patient-reported no family history

Social History (tobacco/alcohol/drugs including frequency, quantity and duration of use):

Cigarettes- 1 whole pack every day for 30 years. Nicotine patch- 1 patch every day for the past 6 months. Beer, 7-8 bottles every day for the past 25 years. Vodka- 2 pints every other day for the past 25 years. No drug uses.

Admission Assessment

Chief Complaint (2 points): Generalized weakness

History of Present Illness – OLD CARTS (10 points): On September 8th, a 54-year-old male was admitted to Mattoon rehabilitation and Health center for rehabilitation care due to weakness and recurrent falls. The patient reported experiencing weakness in his legs for about a month now. The patient reports that he has had an increased amount of falls due to his weakness. Walking, standing up, and transferring have become more difficult due to being weak in both

legs. The patient reports that laying down and sitting make him feel better. The patient is in physical therapy to help him gain strength in his lower legs.

Primary Diagnosis

Primary Diagnosis on Admission (3 points): Alcoholic cirrhosis of the liver with ascites.

Secondary Diagnosis (if applicable): Failure to thrive

Pathophysiology of the Disease, APA format (20 points):

Alcoholic cirrhosis of the liver with ascites is a disease that is caused due to excessive ingestion of alcohol. Alcoholic cirrhosis develops over a long period and is permanent (Capriotti, 2022, p. 771). It can cause severe functional impairment and has a mortality rate of nearly 60% after four years of being diagnosed (Capriotti, 2022, p. 771). Ascites are the presence of excessive fluid in the peritoneal cavity (Perri, 2017). Ascites in cirrhosis cause hypertension and renal sodium retention (Perri, 2017). Symptoms of cirrhosis and ascites are abdominal distension, nausea, lower-extremity edema, jaundice, and reduced mobility (Perri, 2017).

In cirrhosis and ascites, alcohol is a potent hepatocyte toxin (Capriotti, 2022, p. 772). When hepatocytes are constantly damaged due to alcohol, they start to regenerate slower (Capriotti, 2022, p. 772). Fatty liver develops when an individual consumes more than 60 grams of alcohol daily (Capriotti, 2022, p. 772). Constantly ingesting ethanol will inhibit the oxidation of fatty acids in the liver (Capriotti, 2022, p. 772). Fat will accumulate around the hepatocytes and disrupt the organelles' integrity (Capriotti, 2022, p. 772). Disrupting the mitochondrion in hepatocytes leads to free radical release and inflammation (Capriotti, 2022, p.772). Ascites occur because the disruption in pressure forces between intravascular and extravascular fluid space allows fluid from the extravascular to accumulate in the peritoneal cavity (Perri, 2017).

The diagnosis of cirrhosis and ascites requires a blood draw to look at the person's levels involved with the liver. Elevated liver enzymes are accompanied by hypercholesteremia, hypertriglyceridemia, and sometimes hyperbilirubinemia (Capriotti, 2022, p. 772). Specifically, in alcoholic cirrhosis, AST and ALT liver enzymes are elevated (Capriotti, 2022, p. 772). An ultrasound of the liver can also be used to detect fatty infiltration (Capriotti, 2022, p. 772). A liver biopsy also can confirm alcoholic liver disease (Capriotti, 2022, p. 772). Treatment of alcoholic cirrhosis and ascites includes cessation of alcohol use (Capriotti, 2022, p. 772). Patients should be on a high-protein, low-sodium, low-fat diet to help with the liver (Capriotti, 2022, p. 772). Vitamins and minerals such as folate and thiamine are also recommended (Capriotti, 2022, p. 772).

The patient was diagnosed with alcoholic cirrhosis with ascites due to the extreme amount of use of alcohol consumption. The patient drank up to 8 beers a day plus 2 pints of vodka. The patient was diagnosed with this from a blood draw as his AST and ALT liver enzymes were elevated. The patient also had yellowing of the skin and a very distended abdomen due to the ascites. The patient has been feeling very weak due to his diagnosis of ascites and cirrhosis. The patient must go to physical therapy to strengthen his muscles and get his abdomen drained once a week due to the excessive amount of fluid caused by the ascites.

Pathophysiology References (2) (APA):

Capriotti, T. M. (2020). *Davis advantage for pathophysiology: Introductory concepts and clinical perspectives* (2nd ed.). F. A. Davis Company.

<https://fadavisreader.vitalsource.com/books/9781719641470>

Perri, G.-A. (2017). *Ascites in patients with cirrhosis*. Canadian family physician. Retrieved from <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3860926/>

Laboratory Data (20 points)

If laboratory data is unavailable, values will be assigned by the clinical instructor

CBC Highlight All Abnormal Labs—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason for Abnormal Value
RBC (x10 ⁶ /μL)	4.28-5.56	N/A	4.28	N/A
Hgb (g/dL)	13.0-17.0	N/A	13.0	N/A
Hct (%)	38.1-48.9	N/A	42.0	N/A
Platelets (K/μL)	149-393	N/A	150	N/A
WBC (K/μL)	4.0-11.7	N/A	7.0	N/A
Neutrophils (%)	45.3-79.0	N/A	61.3	N/A
Lymphocytes (%)	11.8-45.9	N/A	14.8	N/A
Monocytes (%)	4.4-12.0	N/A	4.4	N/A
Eosinophils (%)	0.0-6.3	N/A	0.0	N/A
Bands (%)	1-5	N/A	N/A	N/A

Chemistry Highlight All Abnormal Labs—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab	Normal Range	Admission Value	Today's Value	Reason For Abnormal
Na- (mmol/L)	136-145	N/A	125	Sodium is low because in cirrhosis it causes an impairment in the renal capacity. It causes the retention of water and that becomes disproportionate to the retention of sodium. This causes a reduction in sodium concentration and hypo-osmolality (Capriotti, 2022).
K+ (mmol/L)	3.5-5.1	N/A	3.5	N/A
Cl- (mmol/L)	98-107	N/A	100	N/A

CO2 (mmol/L)	21-31	N/A	26	N/A
Glucose (mg/dL)	74-109	N/A	98	N/A
BUN (mg/dL)	7-25	N/A	7	N/A
Creatinine (mg/dL)	0.60-1.20	N/A	0.90	N/A
Albumin (g/dL)	3.5-5.2	N/A	2.6	N/A
Calcium (mg/dL)	8.6-10.3	N/A	9.0	N/A
Mag (mg/dL)	1.8-3.0	N/A	2.0	N/A
Phosphate (units/L)	1.7-2.6	N/A	2.6	N/A
Bilirubin (mg/dL)	0.3-1.0	N/A	6	The bilirubin level is high because the portal of blood flow is distorted and accompanied by a decreased hepatic clearance of bilirubin. Also, can be caused due to the inflammation of the liver (Capriotti, 2022).
Alk Phos (units/L)	34-104	N/A	104	N/A
AST (units/mL)	5-40	N/A	100	The AST is high because when liver cells are damaged, AST will leak out into the bloodstream and cause the levels to become high. It also is caused by the depletion of vitamin B6 which is known in chronic alcoholic users (Capriotti, 2022).
ALT (units/mL)	5-35	N/A	98	The ALT is high because of overuse of alcohol which causes the liver to be damaged. When the liver is damaged, it causes ALT to leak into the bloodstream. This then elevates the ALT levels (Capriotti, 2022).

Urinalysis **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Lab Test	Normal Range	Value on Admission	Today's Value	Reason for Abnormal
Color & Clarity	Light yellow and clear	N/A	N/A	N/A
pH	5.0-8.0	N/A	N/A	N/A
Specific Gravity	1.005-1.034	N/A	N/A	N/A
Glucose	Negative	N/A	N/A	N/A
Protein	Negative	N/A	N/A	N/A
Ketones	Negative	N/A	N/A	N/A
WBC	0-5	N/A	N/A	N/A
RBC	0-5	N/A	N/A	N/A
Leukoesterase	Negative	N/A	N/A	N/A

Cultures **Highlight All Abnormal Labs**—Explanations must be in complete sentences and contain in-text citations in APA format.

Test	Normal Range	Value on Admission	Today's Value	Explanation of Findings
Urine Culture	Negative	N/A	N/A	N/A
Blood Culture	Negative	N/A	N/A	N/A
Sputum Culture	Negative	N/A	N/A	N/A
Stool Culture	Negative	N/A	N/A	N/A

Lab Correlations Reference (1) (APA):

Capriotti, T. M. (2020). *Davis advantage for pathophysiology: Introductory concepts and*

clinical perspectives (2nd ed.). F. A. Davis Company.

<https://fadavisreader.vitalsource.com/books/9781719641470>

Kee, J. L. F. (2017). *Pearson Handbook of Laboratory & Diagnostic tests with nursing implications* (8th ed.). Pearson Publication.

Sarah Bush Lincoln Hospital. (2022). *Lab Values*. Sarah Bush Lincoln Hospital.

Diagnostic Imaging

All Other Diagnostic Tests (10 points): Ultrasound of the liver- The patient had a liver ultrasound to confirm the diagnosis of alcoholic cirrhosis. An ultrasound is used to determine the size of the liver (Capriotti, 2022). The ultrasound can show nodular liver surfaces and hypoechoic nodules in liver parenchyma (Capriotti, 2022). The ultrasound allows the investigation of the hepatic tissue through ultrasonic waves (Capriotti, 2022). This ultrasound is essential to the client because it can show how badly the liver has been affected by the overuse of alcohol. It will show abnormalities of the liver and can help with the confirmation of alcoholic cirrhosis.

Diagnostic Imaging Reference (1) (APA):

Capriotti, T. M. (2020). *Davis advantage for pathophysiology: Introductory concepts and clinical perspectives* (2nd ed.). F. A. Davis Company.

<https://fadavisreader.vitalsource.com/books/9781719641470>

Current Medications (10 points, 2 points per completed med)

5 different medications must be completed**Medications (5 required)**

Brand/Generic	Celexa/ Citalopram hydrobromide	Lopressor /Metoprolol Tartrate	Aldactone/ Spironolacto ne	Zyloprim/ Allopurin ol	Protonix /Pantoprazol e sodium
Dose	40 mg	25 mg	25 mg	100 mg	40 mg
Frequency	QD	QD	QD	QD	BID
Route	PO	PO	PO	PO	PO
Classification	Selective serotonin reuptake inhibitor & Antidepressant	Beta- adrenergic blocker & Antihypertensi ve	Potassium- sparing diuretic & Diuretic	Xanthine oxidase inhibitor & Antigout	Proton pump inhibitor & Antiulcer
Mechanism of Action	Blocks serotonin reuptake by adrenergic nerves. The blocked reuptake increases the serotonin levels at the nerves synapse which can reduce depression. (Jones, 2022)	Reduces blood pressure by decreasing the release of renin. Also inhibits the stimulation of beta-receptors sites in the heart. (Jones, 2022)	Spironolacto ne competes with aldosterone and prevents sodium and water reabsorption. This causes their excretion through the distal convoluted tubules. This can reduce blood volume and blood pressure. (Jones, 2022)	Allopurin ol inhibits xanthine oxidase and causes it to block the metabolis m of xanthine to uric acid. This helps relieve the symptoms of gout. (Jones, 2020)	Pantoprazol e inhibits the last step in gastric acid production and blocks the exchange of intracellular hydrogens and extracellular potassium. Prevents additional hydrochloric acid from forming. (Jones, 2022)
Reason Client Taking	To manage depression	To manage hypertension	To treat cirrhosis	To treat gout	To manage gouty arthritis
Contraindicati	Hypersensitivi	Hypotension	Hyperkalemi	Renal	Concurrent

ons (2)	ty to citalopram Coadministrati on with pimozide. (Jones, 2022)	Bradycardia (Jones, 2022)	a Addison’s disease (Jones, 2022)	impairme nt Hepatic disease (Jones, 2022)	therapy with rilpivirine- containing products Hyponatrem ia (Jones, 2022)
Side Effects/Adverse Reactions (2)	Anxiety Impaired concentration (Jones, 2022)	Confusion Dizziness (Jones, 2022)	Hypotension Hyponatremi a (Jones, 2022)	Jaundice Ecchymos is (Jones, 2022)	Hepatotoxici ty Hepatic failure (Jones, 2022)

Medications Reference (1) (APA):

Jones & Bartlett Learning. (2022). *2022 nurse’s drug handbook* (21st ed.). Jones & Bartlett Learning.

Assessment

Physical Exam (18 points) – **HIGHLIGHT ALL PERTINENT ABNORMAL FINDINGS**

<p>GENERAL: Alertness: Orientation: Distress: Overall appearance:</p>	<p>ANO x4. Patient was alert and oriented. Patient was in no acute distress. Patient was well groomed.</p>
<p>INTEGUMENTARY: Skin color: yellow Character: Temperature: Turgor: Rashes: Bruises: Wounds: scratches on left forearm Braden Score: 20 Drains present: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Type:</p>	<p>Patient’s skin color was yellow. Skin was cool, dry, and thin. Patient had scratches on left forearm. Skin turgor was loose and less than 3 seconds. Braden score is 20</p>
<p>HEENT: Head/Neck: Ears: Eyes: Nose: Teeth:</p>	<p>Patients head and neck are symmetrical. Thyroid is non palpable. Trachea is midline with no deviation. Bilateral carotid pulses are palpable and 2+. Bilateral sclera white, bilateral conjunctiva is pink, and bilateral cornea is clear. Bilateral lids are pink and moist without any lesions or discharge. PERRLA is bilaterally and EOM’s intact bilaterally. The nose is midline, and the septum is midline. Turbinate’s are moist and pink bilaterally with no visible drainage or polyps. Bilateral frontal sinuses are nontender to palpation. Tongue and buccal mucosa were pink, and moist, with no lesions. The patient had all 32 teeth present.</p>
<p>CARDIOVASCULAR: Heart sounds: S1, S2, S3, S4, murmur etc. Cardiac rhythm (if applicable): Peripheral Pulses: 2+ in all pulses Capillary refill: <3 Neck Vein Distention: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Edema Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Location of Edema: N/A</p>	<p>S1 and S2 heart sounds were clear and audible without murmurs or gallops. Cardiac rhythm is steady and regular. Carotid and radial pulses were palpable and are 2+. Capillary refill was <3 seconds in fingers bilaterally. No jugular vein distention was seen.</p>

<p>RESPIRATORY: Accessory muscle use: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Breath Sounds: Location, character</p>	<p>No abnormal lung sounds during auscultation. Lung sounds were clear anterior/posterior bilaterally. No accessory muscles were used for respiration. No wheezes, crackles, or rhonchi were noted.</p>
<p>GASTROINTESTINAL: Diet at home: N/A Current Diet: Regular diet Height: 69 inches Weight: 89.4 kg Auscultation Bowel sounds: present in all quadrants Last BM: 2 days ago 10/18/22 Palpation: Pain, Mass etc.: Inspection: Round abdomen Distention: Yes Incisions: N/A Scars: N/A Drains: N/A Wounds: N/A Ostomy: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Nasogastric: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Size: Feeding tubes/PEG tube Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Type:</p>	<p>The patient is on a regular diet. Patient's height is 69 inches and weighs 89.4 kg. Last bowel movement was on 10/18/22. The abdomen was very hard and distended due to ascites. The abdomen was also very round. Able to hear bowel sounds in all quadrants. The patient has no ostomy, nasogastric, or feeding tubes.</p>
<p>GENITOURINARY: Color: Dark yellow urine Character: Quantity of urine: Pain with urination: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Dialysis: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Inspection of genitals: Catheter: Y <input type="checkbox"/> N <input checked="" type="checkbox"/> Type: Size:</p>	<p>The patient reports a weak urine stream and does not urinate often. The patient reports having dark yellow urine.</p>
<p>MUSCULOSKELETAL: Neurovascular status: Unable to assess ROM: Unable to assess Supportive devices: Wheelchair and walker Strength: Unable to assess ADL Assistance: Y <input checked="" type="checkbox"/> N <input type="checkbox"/> Fall Risk: Y <input checked="" type="checkbox"/> N <input type="checkbox"/> Fall Score: 65 Activity/Mobility Status: 1 assist Independent (up ad lib) <input type="checkbox"/></p>	<p>Patient uses a walker to walk and transfer. When the patient is feeling weaker, he uses the wheelchair. Patient needs assistance in dressing, showering, and help with walking. The patient is a fall risk. The fall score is 65.</p>

Needs assistance with equipment <input checked="" type="checkbox"/> Needs support to stand and walk <input checked="" type="checkbox"/>	
NEUROLOGICAL: MAEW: Y <input type="checkbox"/> N <input type="checkbox"/> PERLA: Y <input type="checkbox"/> N <input type="checkbox"/> Strength Equal: Y <input type="checkbox"/> N <input type="checkbox"/> if no - Legs <input type="checkbox"/> Arms <input type="checkbox"/> Both <input type="checkbox"/> Orientation: Mental Status: Speech: Sensory: LOC:	Unable to assess.
PSYCHOSOCIAL/CULTURAL: Coping method(s): Developmental level: Religion & what it means to pt.: Personal/Family Data (Think about home environment, family structure, and available family support):	The patient uses deep breathing exercises to cope. The patient has a high school diploma. Patient is a Christian and utilizes prayer. The patient's parents are deceased. The patient has support from siblings. The patient also has no children.

Vital Signs, 1 set (5 points) – HIGHLIGHT ALL ABNORMAL VITAL SIGNS

Time	Pulse	B/P	Resp Rate	Temp	Oxygen
0912	76 bpm	102/88 LA	18 resp/min	36.7 °C Oral	100% RA

Pain Assessment, 1 set (5 points)

Time	Scale	Location	Severity	Characteristics	Interventions
0838	Numeric pain scale	N/A	0	N/A	N/A

Intake and Output (2 points)

Intake (in mL)	Output (in mL)
100% breakfast	Voided x1

75% lunch	
200 mL of orange juice	
200 mL of water	

Nursing Diagnosis (15 points)
Must be NANDA approved nursing diagnosis

Nursing Diagnosis	Rationale	Interventions (2 per dx)	Outcome Goal (1 per dx)	Evaluation
<ul style="list-style-type: none"> • Include full nursing diagnosis with “related to” and “as evidenced by” components • Listed in order by priority – highest priority to lowest priority pertinent to this client 	<ul style="list-style-type: none"> • Explain why the nursing diagnosis was chosen 			<ul style="list-style-type: none"> • How did the client/family respond to the nurse’s actions? <ul style="list-style-type: none"> • Client response, status of goals and outcomes, modifications to plan.
1.Risk for impaired liver function related to cirrhosis as evidenced by the yellowing of the skin.	This diagnosis was given because the patient has yellowing of the skin.	1.Monitor lab tests. 2.Monitor clinical manifestations of hepatic inflammation	1.Patient will no longer have yellowing of the skin.	The patient is aware that the nurse will be monitoring labs for changes in liver functions. The patient will notify the nurse if there are any changes within himself that can indicate hepatic inflammation.
2.Ineffective	This diagnosis	1. Teach	1. Patient will	The patient is

<p>coping related to the failure to thrive as evidenced by social isolation.</p>	<p>was chosen because the patient is withdrawn from life and is socially isolated.</p>	<p>patient coping skills. 2. Get the patient out of the room and involved with fun activities.</p>	<p>demonstrate the ability to use newly learned coping skills.</p>	<p>compliant with learning new coping skills. The patient showed interest in wanting to learn different coping skills. The patient was a little hesitant with getting out of the room to try new activities but is willing to give it a shot.</p>
--	--	---	--	---

Other References (APA):

Phelps, L. L. (2020). *Sparks and Taylor's nursing diagnosis reference manual* (11th ed.). Wolters Kluwer.

Concept Map (20 Points):



