

5. There were 45,238 neonatal deaths out of 5,672,000 live births in City F. Calculate the neonatal mortality (death) rate. Use 1,000 as the multiplier. Round to the nearest whole number.

$$\frac{45238}{5672000} = 0.00797567$$

7.97 \rightarrow 8 cases per 1000

6. The population in Sangamon county in 2000 was 188,951

- a The number of live births in Sangamon County in 2000 was 2,646. Figure the Live Birth rate for Sangamon country for 2000. Use 1,000 as the multiplier. Round to the nearest whole number.

$$\frac{2646}{188951} = 0.01400363$$

14 births

- b The number of infant deaths in Sangamon County in 2000 was 18. Figure the infant mortality rate for Sangamon country for 2000. Use 1,000 as the multiplier. Round to the nearest whole number.

$$\frac{18}{188951}$$

0.095 \rightarrow 0 deaths

7. A city has a population of 250,000. Of these, 10,000 have disease X, which is incurable. There are 1,000 new cases and 400 deaths each year from this disease. There are 2,500 deaths per year from all causes. What is the prevalence rate based on a multiplier 100,000. Round to the nearest whole number.

$$\frac{\text{\# of cases}}{\text{\# of people population}} = \frac{10000}{250000} = 0.04$$

4000 cases per 100000

Quintile of CRP Level					
	1	2	3	4	5
	0.49 mg/dL	>0.49-1.08 mg/dL	>1.08-2.09 mg/dL	>2.09-4.19 mg/dL	>4.19 mg/dL
Relative Risk	1.0	1.8	2.3	3.2	4.5
Number of women	6000	6000	6000	6000	6000

8. Based on the relative risk data above, one can conclude:
- There is no risk of heart attack/stroke for women with CRP levels in the first quintile.
 - Decreasing CRP level appears to increase the risk of heart attack/stroke.
 - Increasing CRP level appears to increase the risk of heart attack/stroke.
 - There appears to be no association between CRP levels and heart attack/stroke.

9. In 2020, the population of Illinois is 12.63 million. Total cases of COVID-19 is 900,370. Using 100,000 as a multiplier, what is the period prevalence rate? Round to the nearest whole number.

$$\frac{900370}{12.63 \text{ million}} = 0.0071$$

7129 cases per 100000

10. In 2020, the population of Illinois is 12.63 million. The total deaths from COVID-19 is 128,000. Using 100,000 as a multiplier, what is the cause-specific mortality rate? Round to the nearest whole number.

$$\frac{128000}{12630000} = 0.0101346$$

1013.46

1013 cases per 100000

Calculate the following rates using the information provided.

1. The total death in County Z last year was 6,092. The population of County Z last year was 524,263. What was the crude death rate? Use 100,000 as the multiplier. Round to the nearest whole number.

$$\frac{6092}{524263} = 0.01162012$$

1162 cases per 100000 12 per 1000

2. There were 4,953 deaths from neoplasms in City B during the past year. The year-end population was 3,495,678. What was the specific cancer death rate for last year? Use 100,000 as the multiplier. Round to the nearest whole number.

$$\frac{4953}{3495678} = 0.00141689$$

rate 142 per 100000

3. The population of the US in 2000 was 99,421,906. The number of deaths from heart disease in the US in 2000 was 710,760. The total number of deaths in the US in 2000 was 2,403,351.

- a Calculate the percentage (%) of heart disease deaths for the US in 2000. Round to the nearest whole number.

population = 99421906

$$\frac{710760}{2403351} = 0.2957$$

$$\frac{710760}{2403351} = 0.2957 \rightarrow 29.57 = \boxed{30\%}$$

0.2957

29.57

30%

- b Calculate the rate of heart disease deaths in the US in 2000 for the US. Use 100,000 as the multiplier. Round to the nearest whole number.

$$\frac{710760}{99421906} = 0.00714893$$

714.89 → 715 cases per 100000

4. In Illinois in 2000, the population was 12,419,293. The number of Salmonella cases in 2000 was 1,502 in Illinois. Calculate the incidence rate for Salmonella for Illinois in 2000. Use 100,000 as the multiplier. Round to the nearest whole number.

$$\frac{1502}{12419293} = 0.00012094$$

12 cases per 100000