

N434 Evidence-Based Practice

Grammar Practice with Draft Coach

Summer 2024

Instructions: Working in groups of 2-3, students will practice using Grammar feature in Draft Coach. Students will modify the following statements to ensure correct clarity, mechanics, grammar, and punctuation is used.

NOTE: Some examples may have zero (0) errors. If that is the case, put "N/A" as the correction.

Restraints are in use when determining whether the less restrictive interventions are ineffective.
Correction: N/A
Studies haev arisen on wheather restraints are moer beneficial than potential harm.
Correction: Studies have arisen on whether restraints are more beneficial than potential harm.
Restraints can have a negative effect on patients' physical and mental health, a reduction of usage has been recommended for all healthcare settings
Correction: Restraints can have a negative effect on patients' physical and mental health, and a reduction of usage has been recommended for all healthcare settings.
According to Dong et al. (2020), women that test positive for covid-19 may be able to pass antibodies to infants through breast milk and while in gestation.
Correction: According to Dong et al. (2020), women who test positive for Covid-19 may be able to pass antibodies to infants through breast milk and during pregnancy.
Approximately 100 million Americans experience chronic pain daily
Correction: Approximately 100 million Americans experience chronic pain daily.
More than 50% of health care visits in children were due to reported pain.
Correction: More than 50% of healthcare visits in children were due to reported pain.
Regarding nursing management of the disease, nurses can promote optimal nursing care for this disease by achieving airway clearance, improving breathing patterns, encouraging effective coping, and promoting self-care (Hinkle & Cheever, 2018).

Correction: N/A – 2% similarity at another university
In the world hypertension is the primary cause of cardiovascular disease and early death.
Correction: In the world, hypertension is the primary cause of cardiovascular disease and early death. – 4% similarity at another university
I think it is so crazy that global mean blood pressure has remained constant or reduced somewhat during the last four decades due to widespread usage of antihypertensive drugs.
Correction: It is crazy that global mean blood pressure has remained constant or reduced somewhat during the last four decades due to the widespread usage of antihypertensive drugs. – 7% similarity to an article that isn't cited
Approximately 45 percent of individuals had symptoms of GAD for at least two years before being diagnosed with the illness
Correction: Approximately 45 percent of individuals had symptoms of GAD for at least two years before being diagnosed with the illness. – define GAD
This decrease in bone density, can make a patient, more likely to fracture teh bones.
Correction: This decrease in bone density can make a patient more likely to fracture the bones.
The last step, consists of providing resources to each patient; to help them be successful through the transition
Correction: The last step consists of providing resources to each patient to help them be successful through the transition.
I have found that the most frequent comorbidities associated with ADHD are learning disorders such as reading disorders and dyscalculia
Correction: The most frequent comorbidities associated of ADHD are learning disorders, such as reading disorders and dyscalculia.
To expand upon this, according to Joaquin (2019), another factor that contributes to lower nurse-to-patient ratios is spending. Hospitals would have to hire many nurses, increasing the annual cost up to millions.
Correction: To expand upon this, according to Joaquin (2019), another factor that contributes to lower nurse-to-patient ratios is spending. Hospitals would have to hire many nurses, increasing the annual cost to millions.

Although lifestyle changes such as a low salt diet and more exercise are good changes when dealing with hypertension, it shows that when a patient has both hypertension and diabetes, the most common way to control it is through medication.

Correction:

N/A