



Expert Chart Darian Oduya

This expert chart is intended to assist in evaluating student performance in documentation for this activity. Only the tabs and tables of the chart that warrant entries are included.

Darian Oduya, Male	DOB: 7 years old	Attending: John Mack, MD
MRN: 10483948	Height: 119 cm	Code Status: Full Code
Allergies: none	Weight: 25.9 kg	Comments: none

Allergies & Home Medications

Allergies	
No known allergies	Reviewed

Vital Signs

Measurements	
Height/Length	119 cm
Weight	25.9 kg

Copyright ©2023 Assessment Technologies Institute, L.L.C. All rights reserved. This resource is copyright protected material of ATI and is provided for use solely under license by ATI. Copying, distributing, or posting this material on the internet or social media is strictly prohibited. Posting on institution LMS requires ATI's prior written permission.

Orders/Prescriptions

Start Date	Order	Details
17.0 hr before start	<p>Cephalexin</p> <p><i>Entered by Provider</i> Dose/Frequency: 50 mg/kg/day Every 6 hr Route: Oral</p> <p>Start Date: 17.0 hr before start Dispensed: ORAL; SUSPENSION 250 mg/5 mL</p>	<p>5.0 hr before start - Given 6.5 mL</p> <p>11.0 hr before start - Given 6.5 mL</p> <p>17.0 hr before start - Given 6.5 mL</p> <p>Child weight: 25.9 kg = 6.5 mL per dose</p>
17.0 hr before start	<p>Dextrose 5% and 0.9% Sodium Chloride (D5NS)</p> <p><i>Entered by Provider</i> Dose/Frequency: 1000 mL Rate: 67 mL/hr Route: IV</p> <p>Start Date: 17.0 hr before start End Date: Dispensed: INJECTABLE, IV 1000 mL</p>	<p>2.0 hr before start - Given</p> <p>17.0 hr before start - Started</p>
17.0 hr before start	<p>Ibuprofen</p> <p><i>Written order</i> Dose/Frequency: 200 mg Every 6 hr PRN Route: Oral</p> <p>Start Date: 17.0 hr before start End Date: Dispensed: ORAL; SUSPENSION 100 mg/5 mL</p>	<p>6.0 hr before start - Given 200 mg</p> <p>12.0 hr before start - Given 200 mg</p> <p>As needed for pain</p>
17.0 hr before start	<p>Multivitamin</p> <p><i>Entered by Provider</i> Dose/Frequency: 1 tablet daily Route: Oral</p> <p>Start Date: 17.0 hr before start End Date: Dispensed: ORAL; CHEWABLE TABLET</p>	

MAR

Medication Administration for 0 Days after start

Medication	-17 hr	-12 hr	-11 hr	-5 hr	-2 hr
Cephalexin <i>Entered by Provider</i> Dose/Frequency: 50 mg/kg/day Every 6 hr Route: Oral Start Date: 17.0 hr before start Dispensed: ORAL; SUSPENSION 250 mg/5 mL	Given KW		Given KW	Given KW	
Dextrose 5% and 0.9% Sodium Chloride (D5NS) <i>Entered by Provider</i> Dose/Frequency: 1000 mL Rate: 67 mL/hr Route: IV Start Date: 17.0 hr before start End Date: Dispensed: INJECTABLE, IV 1000 mL	Started KW				Given KW
Ibuprofen <i>Written order</i> Dose/Frequency: 200 mg Every 6 hr PRN Route: Oral Start Date: 17.0 hr before start End Date: Dispensed: ORAL; SUSPENSION 100 mg/5 mL		Given KW		Given KW	
Multivitamin <i>Entered by Provider</i> Dose/Frequency: 1 tablet daily Route: Oral Start Date: 17.0 hr before start End Date: Dispensed: ORAL; CHEWABLE TABLET					

Flowsheet

Interventions (Lines, Drains, etc)

IVs/Lines	
<i>LINES</i>	2.0 min after start
Type	Peripheral IV
Action	Assessment
Location	Forearm
Orientation	Right
Site Assessment	Clean, dry, intact

Assessment

Head, Face, Anterior Fontanel, Neck	
Head, Face	Head round and symmetrical, Normocephalic

Respiratory	
Resp. Effort/Pattern	Bilaterally even and unlabored
Breath Sound, Comments	Clear in all lobes bilaterally, anterior and posterior
Oxygen rate and comments	Symmetrical movements of anterior and posterior chest with inspiration and expiration. Rate is 22/min.

Cardiac	
Cardiac Rhythm/Sounds	Regular rhythm, S1 and S2 present
Cardiac Comments	Apical pulse 115/min

Peripheral Vascular	
RUE	Capillary refill less than 3 secs
LUE	Capillary refill less than 3 secs
RLE	Capillary refill less than 3 secs, Pulse is palpable and strong. No edema or pain.
LLE	Capillary refill less than 3 secs, Pulse is palpable and strong
Periph. Vasc. Comments	Swelling noted in left lower leg

Integumentary	
Skin Color	Appropriate for ethnicity, Even distribution
Skin Temperature/Condition	Intact, Warm
Skin Comments	Skin intact with even tone and consistent with genetic background, except for cellulitis on left lower calf, which is erythematous. Skin warm on right lower leg, but not on left lower leg.

Musculoskeletal	
RLE	Full range of mobility
LLE	Full range of mobility
Musculoskeletal Comments	Upper extremities symmetrical in appearance

Gastrointestinal	
Abdomen	Soft, Nontender, Rounded
Bowel Sounds	Active
GI Comment	Symmetrical appearance, umbilicus inverted and at midline

Pain Assessment	
Pain Location	Left leg
Pain Rating - Faces	4
Pain Comments	Unable to describe pain. Appears in pain and tired. States he is "so tired" and his dad reports that the child did not sleep well. Left leg elevated