

## PHARM MED MATH ASSIGNMENT

### Proficiency test 1 – Riley Arend

1. Intravenous (IV)
2. Do not use hs → use at bedtime
3. As needed
4. Don't use OU - use in both eyes
5. Keep the vein open
6. In the rectum
7. Sublingual
8. No known drug allergies
9. Do not use tiw → use three times weekly
10. Milliliter, mL
11. Every 4 hours
12. Do not use cc → use milliliter
13. Do not use SC → use subcutaneous
14. Do not use AU → use each ear
15. No known allergies
16. After meals
17. Do not use qd → use everyday
18. Immediately
19. Controlled release
20. 3 times a day
21. Do not use OS → write out in the left eye
22. Kilogram (kg)
23. To keep open
24. Every hour
25. Do not use OD → write out in the right eye
26. Milliequivalent
27. Before meals
28. Four times a day
29. Microgram (mcg)
30. Intramuscular
31. 1500
32. 1715
33. 0700
34. 1045
35. 0930

### Proficiency Test 2:

1. Pentobarbital one hundred milligrams at hour of sleep, as needed, by mouth.
2. Propranolol hydrochloride forty milligrams by mouth two times a day

- (Example times: 9am and 9pm)
3. Ampicillin one-gram intravenous piggyback every six hours  
(Example times: 6am, noon, 6pm, midnight)
  4. Acetaminophen three hundred twenty-five milligrams, two tablets by mouth immediately.  
(Give two tablets, each tablet is 325mg)
  5. Pilocarpine two drops in both eyes every three hours. Do not use OU; write in both eyes”  
(Example times: 3am, 6am, 9am)
  6. Scopolamine eight-tenths of a milligram subcutaneously immediately.
  7. Nystatin four hundred thousand units’ swish and swallow four times a day.  
(Example times: 10am, 2pm, 6pm, 10pm)
  8. Potassium thirty milliequivalents by mouth twice a day.  
(Example times: 10am and 6pm)
  9. Heparin six thousand units subcutaneous every twelve hours.  
(Example times – 6am and 6pm)
  10. Morphine two to four milligrams intravenous push, as needed, every two to three hours for pain.  
(Example times: 6am, 9am, noon, 3pm, 6pm, 9pm, midnight)
  11. Colace one hundred milligrams by mouth three times a day.  
(Example times: 10am, 2pm, 6pm)
  12. Ativan one milligram intravenous push times one dose now.  
(Example: Administer one dose now.)
  13. Ten milliequivalents potassium chloride in one hundred cubic centimeters of normal saline over one hour, times one dose.  
(Should be one hundred milliliters instead of cubic centimeters)
  14. Tylenol #3 two tablets by mouth every four hours as needed for pain.
  15. Lopressor twenty-five milligrams by mouth twice a day.  
(Example times – 9am, 9pm)