

## N322 Drug Card Template

<b>Brand/Generic</b>	<b>Lovenox/ Enoxaparin</b>
<b>Typical Prescribed Dose</b>	<b>30 mg</b>
<b>Typical Frequency</b>	<b>Q 12hr</b>
<b>Route</b> (most common)	<b>Subcutaneously</b>
<b>Classification</b>	<b>Anticoagulant</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Lovenox is an anticoagulant that speeds up the activity of antithrombin III.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>This drug is used to prevent blood clots, DVT, and pulmonary embolisms.</b>
<b>Contraindications</b> (3 most important)	<b>Three major contraindication are active major bleeding, GI ulcerations, severe kidney or liver disease</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>Central nervous system adverse reactions include confusion, epidural or spinal hematoma and fever.</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Use extreme caution in patients with history oh heparin induced thrombocytopenia and patients with increased risk of hemorrhage.</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	

<b>Client Teaching needs</b> (1 or 2 most important)	<b>Notify prescriber immediately of any adverse events, Emphasize importance of complying with follow up physicals with physician.</b>
---	--

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Heparin</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	<b>IV or Subcutaneously</b>
<b>Classification</b>	<b>Anticoagulant</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>It binds with antithrombins III. Heparin will inactivate thrombin which will prevent existing clot extension and block fibrin.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>To prevent and treat peripheral arterial embolism, thromboembolic complications, pulmonary embolism and venous thrombosis.</b>
<b>Contraindications</b> (3 most important)	<b>Contraindications include breastfeeding women, infants, and neonates.</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>Common adverse reactions include chills, chest pain and fever.</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Always read heparin label carefully to help reduce the risk of miscalculation that could result in medication errors.</b>

<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Explain to the patient that heparin can not be taken orally. Inform the patient about increased risk of bleeding and encourage them to avoid injuries and use soft bristle toothbrushes and an electrical razor</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Coumadin/ Warfin</b>
<b>Typical Prescribed Dose</b>	<b>2-5 mg/d for 2-3 d</b>
<b>Typical Frequency</b>	<b>Daily infused</b>
<b>Route</b> (most common)	<b>PO</b>
<b>Classification</b>	<b>Anticoagulant</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Prevents liver for synthesizing vitamin K, which interferes with clotting and warren prevents coagulation.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Typically prescribed for DVT, PE and embolization associated with atrial fibrillation and prosthetic heart valves.</b>
<b>Contraindications</b> (3 most important)	<b>Three most important contraindications are GI ulcerations, blood disorders associated with bleeding, severe kidney or liver disease.</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Hemorrhage 2. Nausea 3. Vomiting</b>

<b>Nursing Considerations</b> (1 or 2)	<b>Women of childbearing age should have a negative pregnancy test before beginning warfarin therapy. Monitor INR and assess for therapeutic effects as prescribed.</b>
<b>Patient Assessment</b> (how you are monitoring the medication's effect)	<b>Monitor INR and PT results.</b>
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Instruct patient to take drug as prescribed at the same time each evening.</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Xarelto/ Rivaroxaban</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	<b>PO</b>
<b>Classification</b>	<b>Anticoagulant</b>
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Prophylaxis of DVT, treatment and secondary prevention of DVT and stroke prevention in atrial fibrillation.</b>
<b>Contraindications</b> (3 most important)	<b>1. Artificial heart valve surgery 2. Severe liver disease 3. Increased risk of bleeding due to clotting disorder</b>

<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<ol style="list-style-type: none"> <li>1. More likely to bruise more easy</li> <li>2. It may take longer for bleeding to stop</li> <li>3. Increase risk for bleeding, which can be serious</li> </ol>
<b>Nursing Considerations</b> (1 or 2)	It is used to reduce the risk of systemic embolism and stroke and should not be given to patient with moderate or severe hepatic impairment.
<b>Patient Assessment</b> (how you are monitoring the medications effect)	Monitor creatinine
<b>Client Teaching needs</b> (1 or 2 most important)	Emphasize the importance of taking medication as prescribed.

### N322 Drug Card Template

<b>Brand/Generic</b>	Epogen/ Epoetin Alfa
<b>Typical Prescribed Dose</b>	Cancer chemotherapy adult: 150-300 units/kg Chronic kidney disease adults: 50-100 units/kg
<b>Typical Frequency</b>	Chronic kidney disease- Adults-three times daily Cancer Chemotherapy- Adults- three times weekly 1 mon to 12 y, 600 units/ kg IV, sub
<b>Route</b> (most common)	Intravenous or Subcutaneous
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	

<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<ol style="list-style-type: none"> <li>1. Headache</li> <li>2. Body aches</li> <li>3. Diarrhea</li> </ol>
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	-statin family
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	

<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

N322 Drug Card Template

<b>Brand/Generic</b>	<b>Aspirin</b>
<b>Typical Prescribed Dose</b>	<b>325 to 650 mg</b>
<b>Typical Frequency</b>	<b>Q4hr</b>

<b>Route</b> (most common)	<b>PO</b>
<b>Classification</b>	<b>Anti-inflammatory, antiplatelet, antipyretic, nonopioid analgesic</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Blocks the enzyme needed for prostaglandin synthesis. When the inhibition of prostaglandins causes inflammatory symptoms to subside, This we cause pain to be relived because of pain transmission from periphery to the spinal cord. In the hypothalamus aspirin causes peripheral vasodilation, heat loss and diaphoresis.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Prescribed to relieve mild pain or fever</b>
<b>Contraindications</b> (3 most important)	<b>Allergy to tartrazine dye, bleeding problems and asthma</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Confusion 2. CNS depression 3. Hearing loss</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Ask client about tinnitus</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Advise parents not to give aspirin to a child with chickenpox or flu symptoms because of risk of Reye's syndrome</b>

<b>Brand/Generic</b>	<b>Zetia/ ezetimibe</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	<b>antihypercholesterolemic</b>
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Prescribed to treat heterozygous familial and nonfamilial hypercholesterolemia or homozygous sitosterolemia</b>
<b>Contraindications</b> (3 most important)	<b>1. Active hepatic dysfunction 2. hypersensitivity to ezetimibe or its components</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Depression 2. Dizziness 3. Fatigue</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Monitor liver enzymes before and during ezetimibe therapy</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	<b>Monitor liver enzymes</b>

<b>Client Teaching needs</b> (1 or 2 most important)	<b>Instruct patient to follow a low-cholesterol diet. Advice client to report unexplained muscle pain, tenderness.</b>
---	--

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Welchol</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	<b>PO</b>
<b>Classification</b>	<b>Bile acid sequestrant</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>In the intestine Welchol binds with bile acids which will prevent their absorption and will be excreted in feces.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Use in adjunction to diet and exercise to improve glycemic</b>
<b>Contraindications</b> (3 most important)	<b>1. Blood dyscrasia 2. Hypersensitivity to colchicine or its components 3. Serious cardiac conditions</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Peripheral neuropathy 2. Abdominal pain 3. Anorexia</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Stop cochlincine if patient develops evidence to toxicity</b>

<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Aq1</b>	<b>1. Patient should take medication with meals and drink plenty of fluids</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Fibrates (-fibrozil)</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	

<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Remicade/infliximab</b>
<b>Typical Prescribed Dose</b>	<b>3mg</b>
<b>Typical Frequency</b>	<b>Infused over 2 hrs</b>
<b>Route</b> (most common)	<b>IV infusion</b>
<b>Classification</b>	<b>Anti inflammatory</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Prevents TNF-alpha from binding with it receptors.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>To control moderate to sever crohns disease and long term</b>
<b>Contraindications</b> (3 most important)	<b>1. Breast feeding 2. Hypersensitivity to infliximab 3. Monitor for severe heart failure.</b>

<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<ol style="list-style-type: none"> <li>1. Chills</li> <li>2. Fatigue</li> <li>3. Fever</li> </ol>
<b>Nursing Considerations</b> (1 or 2)	<b>Infliximab therapy shouldn't be started in patients with active infections.</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	<b>Monito liver function</b>
<b>Client Teaching needs</b> (1 or 2 most important)	<b>May take up between 1 to 2 weeks to see effect</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	<b>Adalimumab/ Humira</b>
<b>Typical Prescribed Dose</b>	<b>40mg</b>
<b>Typical Frequency</b>	<b>Every other week</b>
<b>Route</b> (most common)	<b>Subcutaneous</b>
<b>Classification</b>	<b>TNF blocker</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Prevents TNF-alpha from binding with it receptors, which TNF-alpha can't produce proinlfammatory cytokines and endothelial permeability.</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>Patient with rheumatoid arthritis</b>

<b>Contraindications</b> (3 most important)	<ol style="list-style-type: none"> <li>1. Active infection</li> <li>2. Breastfeeding</li> <li>3. Hypersensitivity to adalimumab or its components</li> </ol>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<ol style="list-style-type: none"> <li>1. Confusion</li> <li>2. CVA</li> <li>3. Headache</li> </ol>
<b>Nursing Considerations</b> (1 or 2)	Use cautiously in patients with recurrent infections
<b>Patient Assessment</b> (how you are monitoring the medications effect)	Lab work
<b>Client Teaching needs</b> (1 or 2 most important)	Explain to patient first injection must be taken with physician present

### N322 Drug Card Template

<b>Brand/Generic</b>	Enebrel entanercept
<b>Typical Prescribed Dose</b>	50 mg
<b>Typical Frequency</b>	Once a week
<b>Route</b> (most common)	Subcutaneous
<b>Classification</b>	Disease-modifying antirheumatoid drug
<b>Mechanism of Action</b> (brief description use your own words)	Reduces joint inflammation by binding with TNF.

<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>To reduce signs and symptoms of psoriatic pr rheumatoid arthritis</b>
<b>Contraindications</b> (3 most important)	<b>1. Hypersensitivity to entanjrcept 2. Sepsis 3. Hypersensitivity to hamster protein</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Asthenia 2. Chills 3. Demyelination</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Screen patients for latent tuberculosis with a skin test before starting therapy</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Teach patient of proper injection technique</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	• Methotrexate robin
<b>Typical Prescribed Dose</b>	<b>1, 500 mg</b>
<b>Typical Frequency</b>	<b>4 times daily for 2-3 days</b>
<b>Route</b> (most common)	<b>Tablets</b>

<b>Classification</b>	<b>Skeletal muscle relaxant</b>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Depresses CNS which leads to sedation</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>To relieve discomfort caused by acute painful musculoskeletal conditions.</b>
<b>Contraindications</b> (3 most important)	<b>1. Renal disease 2. Hypersensitivity to methocarbamol</b>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<b>1. Dizziness 2. Fever 3. Drowsiness</b>
<b>Nursing Considerations</b> (1 or 2)	<b>Monitor IV site for phlebitis</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	<b>Make sure muscle spasms are relieved</b>
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Take drug as prescribed</b>

### N322 Drug Card Template

<b>Brand/Generic</b>	<ul style="list-style-type: none"> <li>Sandimmune (cyclosporine)</li> </ul>
<b>Typical Prescribed Dose</b>	<ul style="list-style-type: none"> <li>12 to 15 mg/kg</li> </ul>

<b>Typical Frequency</b>	<b>Daily</b>
<b>Route</b> (most common)	<b>Oral, IV</b>
<b>Classification</b>	<ul style="list-style-type: none"> <li>• Antipsoriatic, anti rheumatic, immunosuppressant</li> </ul>
<b>Mechanism of Action</b> (brief description use your own words)	<b>Cause suppression of the immune system by inhibiting the proliferation of T lymphocytes</b>
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	<b>To prevent or treat organ rejection</b>
<b>Contraindications</b> (3 most important)	<ol style="list-style-type: none"> <li>1. <b>Abnormal renal function</b></li> <li>2. <b>Neoplastic diseases</b></li> <li>3. <b>Uncontrolled hypertension</b></li> </ol>
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	<ol style="list-style-type: none"> <li>1. <b>Altered level of consciousness</b></li> <li>2. <b>Confusion</b></li> <li>3. <b>Headache</b></li> </ol>
<b>Nursing Considerations</b> (1 or 2)	<b>Be aware that rapid IV infusion may cause acute nephrotoxicity</b>
<b>Patient Assessment</b> (how you are monitoring the medications effect)	<b>Organ not being rejected</b>
<b>Client Teaching needs</b> (1 or 2 most important)	<b>Instruct patient to take medication at the same time each day</b>

N322 Drug Card Template

<b>Brand/Generic</b>	<b>Profrac tacrolimus</b>
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	

<b>Client Teaching needs</b> (1 or 2 most important)	
---	--

N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	

<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	

<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	

<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	

<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	

<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	

<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	

<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	

<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

N322 Drug Card Template

<b>Brand/Generic</b>	
----------------------	--

<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

## N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	

<b>Client Teaching needs</b> (1 or 2 most important)	
---	--

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	

<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	

<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	

<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	
<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	

<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	

### N322 Drug Card Template

<b>Brand/Generic</b>	
<b>Typical Prescribed Dose</b>	
<b>Typical Frequency</b>	
<b>Route</b> (most common)	
<b>Classification</b>	
<b>Mechanism of Action</b> (brief description use your own words)	

<b>Reasons Prescribed</b> (conditions in which this drug is usually prescribed)	
<b>Contraindications</b> (3 most important)	
<b>Common Side Effects/Adverse Reactions</b> (3 most common)	
<b>Nursing Considerations</b> (1 or 2)	
<b>Patient Assessment</b> (how you are monitoring the medications effect)	
<b>Client Teaching needs</b> (1 or 2 most important)	