

Coombs test

From Wikipedia, the free encyclopedia
[Jump to navigation](#)[Jump to search](#)



This article **needs additional citations for verification**. Please help [improve this article](#) by [adding citations to reliable sources](#). Unsourced material may be challenged and removed.

Find sources: ["Coombs test"](#) – [news](#) · [newspapers](#) · [books](#) · [scholar](#) · [JSTOR](#) (October 2015) ([Learn how and when to remove this template message](#))

Coombs test	
<i>Medical diagnostics</i>	
MeSH	D003298
MedlinePlus	003344

A **Coombs test** (also known as **antiglobulin test** or **AGT**) is either of two [clinical blood tests](#) used in [immunohematology](#) and [immunology](#). The two Coombs tests are the **direct Coombs test (DCT)**, also known as **direct antiglobulin test** or **DAT**, and the **indirect Coombs test** (also known as **indirect antiglobulin test** or **IAT**). The direct Coombs test detects antibodies that are stuck to the surface of the red blood cells.^[1] Since these antibodies sometimes destroy red blood cells, a person can be anemic and this test can help clarify the condition. The indirect Coombs detects antibodies that are floating freely in the blood.^[1] These antibodies could act against certain red blood cells and the test can be done to diagnose reactions to a [blood transfusion](#).^[1]


The direct Coombs test is used to test for [autoimmune hemolytic anemia](#)—that is, a condition where the immune system breaks down [red blood cells](#), leading to [anemia](#). The direct Coombs test is used to detect antibodies or [complement proteins](#) attached to the surface of red blood cells. To perform the test, a blood sample is taken and the red blood cells are washed (removing the patient's own plasma and unbound antibodies from the red blood cells) and then incubated with anti-human globulin ("Coombs reagent"). If the red cells then [agglutinate](#), the direct Coombs test is positive, a visual indication that antibodies or complement proteins are bound to the surface of red blood cells and may be causing destruction of those cells.

The indirect Coombs test is used in prenatal testing of pregnant women and in testing prior to a blood transfusion. The test detects antibodies against foreign red blood cells. In this case, [serum](#) is extracted from a blood sample taken from the patient. The serum is incubated with foreign red blood cells of known [antigenicity](#). Finally, anti-human globulin is added. If agglutination occurs, the indirect Coombs test is positive.^[2]



- 1Mechanism
- 2Direct Coombs test
 - o 2.1Examples of diseases that give a positive direct Coombs test
 - 2.1.1Examples of alloimmune hemolysis
 - 2.1.2Examples of autoimmune hemolysis/immunohemolytic hemolysis
 - 2.1.3Drug-induced immune-mediated hemolysis
 - o 2.2Laboratory
- 3Indirect Coombs test
 - o 3.1Examples of clinical uses of the indirect Coombs test
 - 3.1.1Blood transfusion preparation
 - 3.1.2Antenatal antibody screening
 - o 3.2Laboratory method
 - 3.2.1First stage
 - 3.2.2Second stage
 - 3.2.3Titrations
- 4Coombs reagent
- 5Enhancement media
- 6History
- 7In popular culture
- 8References
- 9External links

Mechanism^[edit]



Schematic showing the **direct and indirect Coombs tests**.

The two Coombs tests are based on anti-human [antibodies](#) binding to human antibodies, commonly [IgG](#) or [IgM](#). These anti-human antibodies are produced by [plasma cells](#) of non-human animals after immunizing them with [human plasma](#). Additionally, these anti-human antibodies will also bind to human antibodies that may be fixed onto [antigens](#) on the surface of [red blood cells](#) (RBCs). In the appropriate test tube conditions, this can lead to [agglutination](#) of RBCs and

allowing for visualisation of the resulting clumps of RBCs. If clumping is seen, the Coombs test is positive; if not, the Coombs test is negative.

Common clinical uses of the Coombs test include the preparation of blood for [transfusion](#) in [cross-matching](#), atypical antibodies in the [blood plasma](#) of [pregnant](#) women as part of [antenatal care](#), and detection of antibodies for the diagnosis of immune-mediated [haemolytic anemias](#).

Coombs tests are performed using RBCs or serum (direct or indirect, respectively) from venous whole blood samples which are taken from patients by [venipuncture](#). The venous blood is taken to a laboratory (or blood bank), where trained scientific technical staff do the Coombs tests. The clinical significance of the result is assessed by the [physician](#) who requested the Coombs test, perhaps with assistance from a laboratory-based [hematologist](#).

Direct Coombs test^[edit]

The direct Coombs test (also known as the **direct antiglobulin test** or DAT) is used to detect if antibodies or [complement system](#) factors have bound to [RBCs](#) surface antigens *in vivo*. The DAT is not currently required for pre-transfusion testing but may be included by some laboratories.^{[[medical citation needed](#)][[clarification needed](#)]}

Examples of diseases that give a positive direct Coombs test^[edit]

The direct Coombs test is used clinically when immune-mediated [hemolytic anemia](#) (antibody-mediated destruction of RBCs) is suspected. A positive Coombs test indicates that an immune mechanism is attacking the patient's [RBCs](#). This mechanism could be [autoimmunity](#), [alloimmunity](#) or a drug-induced immune-mediated mechanism.

Examples of alloimmune hemolysis^[edit]

- [Hemolytic disease of the newborn](#) (also known as HDN or erythroblastosis fetalis)
 - [Rh D hemolytic disease of the newborn](#) (also known as Rh disease)
 - [ABO hemolytic disease of the newborn](#) (the direct Coombs test may only be weakly positive)
 - [Anti-Kell hemolytic disease of the newborn](#)
 - [Rh c hemolytic disease of the newborn](#)
 - [Rh E hemolytic disease of the newborn](#)
 - Other blood group incompatibility (RhC, Rhe, Kidd, Duffy, MN, P and others)
- Alloimmune hemolytic [transfusion reactions](#)

Examples of autoimmune hemolysis/immunohemolytic hemolysis^[edit]

- [Warm antibody autoimmune hemolytic anemia](#)
 - Idiopathic
 - [Systemic lupus erythematosus](#)
 - [Evans' syndrome](#) (antiplatelet antibodies and hemolytic antibodies)
- [Cold antibody immunohemolytic anemia](#)
 - [Idiopathic cold hemagglutinin syndrome](#)
 - [Waldenström's macroglobulinemia](#)
 - [Infectious mononucleosis](#)

- o [Paroxysmal cold hemoglobinuria](#) (rare)

Drug-induced immune-mediated hemolysis[\[edit\]](#)

- [Methyldopa](#) (IgG mediated type II hypersensitivity)
- [Penicillin](#) (high dose)
- [Quinidine](#) (IgM mediated activation of classical complement pathway and Membrane attack complex, MAC)

(A [memory device](#) to remember that the *DAT* tests the [RBCs](#) and is used to test infants for *haemolytic disease of the newborn* is: *Rh Disease*; **R** = RBCs, **D** = DAT.)

Laboratory[\[edit\]](#)

The patient's [RBCs](#) are washed (removing the patient's own [serum](#)) and then centrifuged with [antihuman globulin](#) (also known as Coombs reagent). If [immunoglobulin](#) or complement factors have been fixed on to the RBC surface [in-vitro](#), the antihuman globulin will [agglutinate](#) the RBCs and the direct Coombs test will be positive. (A visual representation of a positive direct Coombs test is shown in the upper half of the schematic).

Indirect Coombs test[\[edit\]](#)

The indirect Coombs test (also known as the **indirect antiglobulin test** or IAT) is used to detect [in-vitro](#) antibody-antigen reactions. It is used to detect very low concentrations of antibodies present in a patient's plasma/serum prior to a blood transfusion. In antenatal care, the IAT is used to screen pregnant women for antibodies that may cause [haemolytic disease of the newborn](#). The IAT can also be used for [compatibility testing](#), antibody identification, RBC phenotyping, and titration studies.

Examples of clinical uses of the indirect Coombs test[\[edit\]](#)

Blood transfusion preparation[\[edit\]](#)

Main articles: [blood transfusion](#) and [cross-matching](#)

The indirect Coombs test is used to screen for antibodies in the preparation of [blood](#) for [blood transfusion](#). The donor's and recipient's blood must be [ABO](#) and Rh D compatible. Donor blood for transfusion is also screened for infections in separate processes.

- Antibody screening

A blood sample from the recipient and a blood sample from every unit of [donor](#) blood are screened for antibodies with the indirect Coombs test. Each sample is incubated against a wide range of RBCs that together exhibit a full range of surface antigens (i.e. [blood types](#)).

- Cross matching

The indirect Coombs test is used to test a sample of the recipient's serum for antibodies against a sample of the [blood donor's](#) RBCs. This is sometimes called [cross-matching](#) blood.

Antenatal antibody screening[\[edit\]](#)

The indirect Coombs test is used to screen pregnant women for [IgG antibodies](#) that are likely to pass through the [placenta](#) into the fetal blood and cause [haemolytic disease of the newborn](#).

Laboratory method[\[edit\]](#)

The IAT is a two-stage test. (A cross match is shown visually in the lower half of the schematic as an example of an indirect Coombs test).

First stage[\[edit\]](#)

Washed test [red blood cells](#) (RBCs) are incubated with a known human serum. If the serum contains [antibodies](#) to antigens on the RBC surface, the antibodies will bind onto the surface of the RBCs.

Second stage^[edit]

The RBCs are washed three or four times with isotonic saline solution and then incubated with antihuman globulin. If antibodies have bound to RBC surface antigens in the first stage, RBCs will agglutinate when incubated with the [antihuman globulin](#) (also known Coombs reagent) in this stage, and the indirect Coombs test will be positive.

Titration^[edit]

By diluting a serum containing antibodies the quantity of the antibody in the serum can be gauged. This is done by using doubling dilutions of the serum and finding the maximum dilution of test serum that is able to produce agglutination of relevant RBCs.

Coombs reagent^[edit]

Coombs reagent (also known as **Coombs antiglobulin** or **antihuman globulin**) is used in both the direct Coombs test and the indirect Coombs test. Coombs reagent is antihuman [globulin](#). It is made by injecting human globulin into animals, which produce [polyclonal antibodies](#) specific for human immunoglobulins and human complement system factors. More specific Coombs reagents or [monoclonal antibodies](#) can be used.

Enhancement media^[edit]

Both [IgM](#) and [IgG antibodies](#) bind strongly with their complementary antigens. [IgG](#) antibodies are most reactive at 37 °C. IgM antibodies are easily detected in [saline](#) at [room temperature](#) as IgM antibodies are able to bridge between RBC's owing to their large size, efficiently creating what is seen as [agglutination](#). IgG antibodies are smaller and require assistance to bridge well enough to form a visual [agglutination](#) reaction. Reagents used to enhance IgG detection are referred to as potentiators. RBCs have a net negative charge called zeta potential which causes them to have a natural repulsion for one another. Potentiators reduce the zeta potential of RBC membranes. Common potentiators include low ionic strength solution (LISS), [albumin](#), [polyethylene glycol](#) (PEG), and proteolytic [enzymes](#).

History^[edit]

The Coombs test was first described in 1945 by [Cambridge immunologists Robin Coombs](#) (after whom it is named), [Arthur Mourant](#) and [Rob Race](#).^[a] Historically, it was done in [test tubes](#). Today, it is commonly done using [microarray](#) and gel technology.

In popular culture^[edit]

In [Arthur Hailey's](#) 1959 book *The Final Diagnosis*, the wife of a hospital staff member is pregnant; she is [Rh negative](#) and he is Rh positive. He mentions this, and recommends an indirect Coombs test to the new assistant pathologist, who presumes this to be obvious, and even signs the requisition form to order Coombs serum. Due to extreme penny-pinching by the chief [pathologist](#), he overrides the other doctor's decision to order Coombs serum, and refuses to perform an indirect Coombs test, instead deciding that protein and saline tests alone are sufficient. This decision ends up causing tragic consequences.

References^[edit]
