

Table 1. Electronic Fetal Monitoring Definitions

Pattern	Definition
Baseline	• The mean FHR rounded to increments of 5 beats per minute during a 10-minute segment, excluding: —Periodic or episodic changes —Periods of marked FHR variability —Segments of baseline that differ by more than 25 beats per minute • The baseline must be for a minimum of 2 minutes in any 10-minute segment, or the baseline for that time period is indeterminate. In this case, one may refer to the prior 10-minute window for determination of baseline. • Normal FHR baseline: 110–160 beats per minute • Tachycardia: FHR baseline is greater than 160 beats per minute • Bradycardia: FHR baseline is less than 110 beats per minute
Baseline variability	• Fluctuations in the baseline FHR that are irregular in amplitude and frequency • Variability is visually quantitated as the amplitude of peak-to-trough in beats per minute. —Absent—amplitude range undetectable —Minimal—amplitude range detectable but 5 beats per minute or fewer —Moderate (normal)—amplitude range 6–25 beats per minute —Marked—amplitude range greater than 25 beats per minute
Acceleration	• A visually apparent abrupt increase (onset to peak in less than 30 seconds) in the FHR • At 32 weeks of gestation and beyond, an acceleration has a peak of 15 beats per minute or more above baseline, with a duration of 15 seconds or more but less than 2 minutes from onset to return. • Before 32 weeks of gestation, an acceleration has a peak of 10 beats per minute or more above baseline, with a duration of 10 seconds or more but less than 2 minutes from onset to return. • Prolonged acceleration lasts 2 minutes or more but less than 10 minutes in duration. • If an acceleration lasts 10 minutes or longer, it is a baseline change.
Early deceleration	• Visually apparent usually symmetrical gradual decrease and return of the FHR associated with a uterine contraction • A gradual FHR decrease is defined as from the onset to the FHR nadir of 30 seconds or more. • The decrease in FHR is calculated from the onset to the nadir of the deceleration. • The nadir of the deceleration occurs at the same time as the peak of the contraction. • In most cases the onset, nadir, and recovery of the deceleration are coincident with the beginning, peak, and ending of the contraction, respectively.
Late deceleration	• Visually apparent usually symmetrical gradual decrease and return of the FHR associated with a uterine contraction • A gradual FHR decrease is defined as from the onset to the FHR nadir of 30 seconds or more. • The decrease in FHR is calculated from the onset to the nadir of the deceleration. • The deceleration is delayed in timing, with the nadir of the deceleration occurring after the peak of the contraction. • In most cases, the onset, nadir, and recovery of the deceleration occur after the beginning, peak, and ending of the contraction, respectively.
Variable deceleration	• Visually apparent abrupt decrease in FHR • An abrupt FHR decrease is defined as from the onset of the deceleration to the beginning of the FHR nadir of less than 30 seconds. • The decrease in FHR is calculated from the onset to the nadir of the deceleration. • The decrease in FHR is 15 beats per minute or greater, lasting 15 seconds or greater, and less than 2 minutes in duration. • When variable decelerations are associated with uterine contractions, their onset, depth, and duration commonly vary with successive uterine contractions.
Prolonged deceleration	• Visually apparent decrease in the FHR below the baseline • Decrease in FHR from the baseline that is 15 beats per minute or more, lasting 2 minutes or more but less than 10 minutes in duration. • If a deceleration lasts 10 minutes or longer, it is a baseline change.
Sinusoidal pattern	• Visually apparent, smooth, sine wave-like undulating pattern in FHR baseline with a cycle frequency of 3–5 per minute which persists for 20 minutes or more.

Abbreviation: FHR, fetal heart rate.

Macones GA, Hankins GD, Spong CY, Hauth J, Moore T. The 2008 National Institute of Child Health and Human Development workshop report on electronic fetal monitoring: update on definitions, interpretation, and research guidelines. *Obstet Gynecol* 2008;112:661–6.