



MNA Study Guide – Test #4

Chapters 25-35

| | |
|--|--|
| Histamine (H2)- Receptor Antagonists | Administration of Ophthalmic Agents |
| Coating Agents | Natacyn |
| Metoclopramide (Reglan) | Ocufen, Profenal, Voltaren |
| Antacids | Drug therapy for Glaucoma |
| Gastro-Intestinal Prostaglandins | Signs & Symptoms Osmotic Agent for Glaucoma |
| Anti-Spasmadic Agents | Cholinergic Agent for Glaucoma |
| Anti-Cholinergic Agents | Prostaglandin Agonists for Glaucoma |
| Laxatives | Artificial Tears |
| Anti-Thyroid Agents | Side Effects from Chemotherapy |
| Hypoglycemia | Dantrolene (Dantrium) |
| Hyperglycemia | Use and side effects from centrally acting skeletal muscle relaxants |
| Repaglinide (Prandin) Parameters | Baclofen |
| Insulin Guidelines | Allopurinol |
| Sulfonylurea Oral Hypo-Glycemic Agents | Action of centrally acting skeletal muscle relaxants |
| Meglitinide Oral Hypo-Glycemic Agents | Action of direct-acting skeletal muscle relaxants |
| Glucocorticoid | Gout medications and their side effects |
| Estrogen | Neo-Fradin side effects |
| Progestins | Zithromax |
| Testosterone | Flagyl |
| Corticosteroid Therapy | Tamiflu |
| Oral Contraceptive Package | Cipro |
| Trans-Dermal Patch Application | Ethambutol (Myambutol) |
| Seasonale- Oral Contraceptive | Rifampin |
| Benign Prostatic Drug Therapy | Ancobon |
| Hyperplasia | Percutaneous endoscopic gastrostomy (PEG) tube |
| Uroxatral | Griseofulvin (Fulvicin and Grifulvin) |
| Macrodantin & Furadantin | |