

Ana DaCosta

Unit 6 AS2: Choosing the Appropriate Hypothesis Test

SECTION I:

SCENARIO #1

In a classic study of problem solving, Duncker (1945) asked subjects to mount a candle on a wall in an upright position so that it would burn normally. One group of subject was given a candle, a book of matches, and a box of tacks. A second group was given the same items, except the tacks and the box were presented separately as two distinct items. The solution to this problem involves using the tacks to mount the box on the wall, which creates a shelf for the candle. Duncker reasoned that the first group of subjects would have trouble seeing a “new” function for the box (a shelf) because it was already serving a function (holding tacks). For each subject, the amount of time to solve the problem was recorded.

What is the appropriate hypothesis test to analyze these data? **Independent Measures t-test.**

Why did you choose this test? **Because it is a Between-Subject research design. The researcher is comparing the control group (placebo or in this case no tacks) to the experimental group (box of tacks)**

SCENARIO #2

A researcher for a cereal company wanted to demonstrate the health benefits of eating oatmeal. A sample of 9 volunteers was obtained and each subject ate a fixed diet without any oatmeal for 30 days. At the end of the 30-day period, cholesterol was measured for each individual. Then the subjects began a second 30-day period in which they repeated exactly the same diet except for 2 cups of oatmeal each day. After the second 30-day period, cholesterol levels were measured again.

What is the appropriate hypothesis test to analyze these data? **Dependent Measures t-test**

Why did you choose this test? **Because the research is a within-subject design and participants are measured 2 times, before and after the IV.**

SCENARIO #3

Suppose that scores on the Scholastic Achievement Test (SAT) form a normal distribution with a mean of 500 and a standard deviation of 100. A high school counselor has developed a special course designed to boost SAT scores. A random sample of $n = 16$ students is selected to take the course and then the SAT. The sample has a mean of 554. Was the course worth taking?

What is the appropriate hypothesis test to analyze these data? **Z-test**

Why did you choose this test? **Because the research is to compare the sample mean to the population mean.**

SCENARIO #4

Students often better remember information if they review their notes before going to bed. A researcher wished to empirically demonstrate this phenomenon. A sample of $n=40$ students participated in this study. After being given a list of words to memorize before going to bed, all participants were tested the next day at 8a.m. Half of his participants studied up to 1 hour prior to going to bed and the other half studied up to 3 hours prior to going to bed.

What hypothesis test did the researcher likely use to analyze the data? **Independent Measure t-test.**

Why did you choose this hypothesis test? **Because it is a two group between subject research design with 2 groups.**

SCENARIO #5

A clinical psychologist has noted that autistic children seem to respond to treatment better if they are in a familiar environment. To evaluate the influence of environment, the psychologist selects a group of $n=50$ autistic children who are currently in treatment and randomly assigns the sample to (a) standard clinical treatment, (b) at home clinical treatment, and (c) half standard and half home clinical treatment.

What hypothesis test did the researchers most likely use to analyze the data? **One way Independent measures ANOVA**

Why did you choose this hypothesis test? **Because it is a between-subject research design with more than 2 groups.**

SCENARIO #6

Researchers wanted to investigate whether soccer players (who can sometimes get hit in the head with the ball) suffered any neurological deficits. These researchers measured neurological deficits in soccer players and compared the soccer player data to the data of non-soccer players believed to not engage in any other activity that could deliver blows to the head. The researchers did find significant differences between the soccer players and the non-soccer players.

What hypothesis test should the researcher use to analyze her data? **A Z-test**

Why did you choose this hypothesis test? **Because the research is comparing the sample result from soccer players to the population of non-soccer players**

SECTION II:

This section requires that you choose the appropriate hypothesis test for each data set, run the test on SPSS, and write up your conclusions in APA format.

1. Analyze the following data from a randomized two group experiment, using the appropriate hypothesis test:

Control	Experimental
2, 3, 5, 7, 4, 3, 2, 1, 5,	4, 6, 5, 7, 8, 9, 7, 6, 8

Cut and paste your SPSS results here:

Group Statistics					
	VAR0000			Std.	Std. Error
	2	N	Mean	Deviation	Mean
VAR0000	1.00	9	6.6667	1.58114	.52705
1	2.00	9	3.5556	1.87824	.62608

Independent Samples Test

		Levene's Test for Equality of Variances		t-test for Equality of Means						
		F	Sig.	t	df	Sig. (2-tailed)	Mean Difference	Std. Error Difference	95% Confidence Interval of the Difference	
									Lower	Upper
VAR0001	Equal variances assumed	.324	.577	3.802	16	.002	3.11111	.81838	1.37621	4.84601
	Equal variances not assumed			3.802	15.548	.002	3.11111	.81838	1.37211	4.85012

Write up your results here:

Probability value .002. Reject the Null hypotheses.

The Mean for the control group was 3.55 with SD=1.87. The mean for the experimental group was 6.66 with SD=1.58. The data reached significance there is significant difference between the groups. $t(16)=2.802, p<.05$

2. Analyze the following data measuring heart rate before and after watching a horror film:

Before	After
65	70
73	74
74	78
82	83
71	80
62	65
64	72

Cut and Paste your SPSS results here:

Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	VAR00001	74.5714	7	6.21442	2.34883
	VAR00002	70.1429	7	7.01020	2.64961

Paired Samples Test

		Paired Differences							
		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference		t	df	Sig. (2-tailed)
					Lower	Upper			
Pair 1	VAR00001 - VAR00002	4.42857	3.15474	1.19238	1.51092	7.34622	3.714	6	.010

Write up your research results here:

Reject the Null hypotheses. Probability value .010

The mean for before IV group was 70.14 with a SD 7.01. The mean for after IV group was 74.57 with a SD 6.21. Our data support that the IV have an effect on data, there was significant difference between groups. $t(6) = 3.714, p < .05$

3. Analyze the following data from a randomized experiment that examined the effect of hours of sleep on happiness.

6 hours	8 hours	10 hours
10	10	9
15	11	9
11	10	9
9	11	8
8	13	9
12	13	7

Cut and paste your SPSS results, including the post-hoc test here:

Descriptives

VAR00001

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
6.00	6	10.8333	2.48328	1.01379	8.2273	13.4394	8.00	15.00
8.00	6	11.3333	1.36626	.55777	9.8995	12.7671	10.00	13.00
10.00	6	8.5000	.83666	.34157	7.6220	9.3780	7.00	9.00
Total	18	10.2222	2.04524	.48207	9.2051	11.2393	7.00	15.00

ANOVA

VAR00001

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	27.444	2	13.722	4.714	.026
Within Groups	43.667	15	2.911		
Total	71.111	17			

Multiple Comparisons

Dependent Variable: VAR00001

Scheffe

(I)	(J)	Mean Difference (I-J)	Std. Error	Sig.	95% Confidence Interval	
VAR00002	VAR00002				Lower Bound	Upper Bound
6.00	8.00	-.50000	.98507	.880	-3.1733	2.1733
	10.00	2.33333	.98507	.092	-.3399	5.0066
8.00	6.00	.50000	.98507	.880	-2.1733	3.1733
	10.00	2.83333*	.98507	.037	.1601	5.5066
10.00	6.00	-2.33333	.98507	.092	-5.0066	.3399
	8.00	-2.83333*	.98507	.037	-5.5066	-.1601

*. The mean difference is significant at the 0.05 level.

Write the statistical statement for your results: $F(2,15) = 4.714, p = .026$

Were the data significant? Yes No

If yes, where was the significance found? The data indicates a significant difference between the 8 hour group and the 10 hour group. However, there is no statistically significant difference between the 6 hour group and 8 hour group, nor between the 10 hour group and the 6 hour group. $F(2,15) = 4.714, p = .026$