

Unit 6 AS2: Choosing the Appropriate Hypothesis Test

SECTION I:

SCENARIO #1

In a classic study of problem solving, Duncker (1945) asked subjects to mount a candle on a wall in an upright position so that it would burn normally. One group of subjects was given a candle, a book of matches, and a box of tacks. A second group was given the same items, except the tacks and the box were presented separately as two distinct items. The solution to this problem involves using the tacks to mount the box on the wall, which creates a shelf for the candle. Duncker reasoned that the first group of subjects would have trouble seeing a “new” function for the box (a shelf) because it was already serving a function (holding tacks). For each subject, the amount of time to solve the problem was recorded.

What is the appropriate hypothesis test to analyze these data? ___ independent measures t test _____

Why did you choose this test? _____ I chose the independent measures t test because we are given two groups and are comparing the mean difference from both groups. _____

SCENARIO #2

A researcher for a cereal company wanted to demonstrate the health benefits of eating oatmeal. A sample of 9 volunteers was obtained and each subject ate a fixed diet without any oatmeal for 30 days. At the end of the 30-day period, cholesterol was measured for each individual. Then the subjects began a second 30-day period in which they repeated exactly the same diet except for 2 cups of oatmeal each day. After the second 30-day period, cholesterol levels were measured again.

What is the appropriate hypothesis test to analyze this data? ___ Dependent measures t-test _____

Why did you choose this test? ___ I chose the dependent measures t-test because this study used the same sample of 9 volunteers. They tested the subjects for 30 days, then tested the same subjects again for another 30 days comparing the data. _____

SCENARIO #3

Suppose that scores on the Scholastic Achievement Test (SAT) form a normal distribution with a mean of 500 and a standard deviation of 100. A high school counselor has developed a special course designed to boost SAT scores. A random sample of $n = 16$ students is selected to take the course and then the SAT. The sample has a mean of 554. Was the course worth taking?

What is the appropriate hypothesis test to analyze this data? ___ Z-test _____

Why did you choose this test? _____ I chose z-test because we are given the population standard deviation and are also given the mean. _____

SCENARIO #4

Students often better remember information if they review their notes before going to bed. A researcher wished to empirically demonstrate this phenomenon. A sample of $n=40$ students participated in this study. After being given a list of words to memorize before going to bed, all participants were tested the next day at 8a.m. Half of his participants studied up to 1 hour prior to going to bed and the other half studied up to 3 hours prior to going to bed.

What hypothesis test did the researcher likely use to analyze the data? Independent measures t test

Why did you choose this hypothesis test? I chose the independent measures t test because we are comparing the means of 2 different groups.

SCENARIO #5

A clinical psychologist has noted that autistic children seem to respond to treatment better if they are in a familiar environment. To evaluate the influence of the environment, the psychologist selects a group of $n=50$ autistic children who are currently in treatment and randomly assigns the sample to (a) standard clinical treatment, (b) at home clinical treatment, and (c) half standard and half home clinical treatment.

What hypothesis test did the researchers most likely use to analyze the data? One way independent ANOVA test

Why did you choose this hypothesis test? I chose this hypothesis test because there are 3 different groups being tested here. The standard clinical treatment, at home clinical treatment and half standard/half at home clinical treatment.

SCENARIO #6

Researchers wanted to investigate whether soccer players (who can sometimes get hit in the head with the ball) suffered any neurological deficits. These researchers measured neurological deficits in soccer players and compared the soccer player data to the data of non-soccer players believed to not engage in any other activity that could deliver blows to the head. The researchers did find significant differences between the soccer players and the non-soccer players.

What hypothesis test should the researcher use to analyze her data? Independent samples t-test

Why did you choose this hypothesis test? I chose the independent samples t test because because we are comparing two different groups, one that participates in soccer that takes blows to the head and another group that does not.

SECTION II:

This section requires that you choose the appropriate hypothesis test for each data set, run the test on SPSS, and write up your conclusions in APA format.

1. Analyze the following data from a randomized two group experiment, using the appropriate hypothesis test:

Control	Experimental
2, 3, 5, 7, 4, 3, 2, 1, 5,	4, 6, 5, 7, 8, 9, 7, 6, 8

Cut and paste your SPSS results here:

Paired Samples Statistics

		Mean	N	Std. Deviation	Std. Error Mean
Pair 1	VAR00001	74.5714	7	6.21442	2.34883
	VAR00002	70.1429	7	7.01020	2.64961

Paired Samples Correlations

		N	Correlation	Sig.
Pair 1	VAR00001 & VAR00002	7	.893	.007

Paired Samples Test

		Mean	Std. Deviation	Std. Error Mean	95% Confidence Interval of the Difference		t	df	Sig. (2-tailed)
					Lower	Upper			
Pair 1	VAR00001 - VAR00002	4.42857	3.15474	1.19238	1.51092	7.34622	3.714	6	.010

Write up your research results here:

_____ The mean for measuring heart rate before a horror film was $M=70.1429$ with a $SD=7.01$. The mean for measuring heart rate after a horror film was $M=74.5714$ with a $SD=6.2$. Our data supports that watching a horror film did have an effect on heart rate, $t(6)=3.714$, $p<.05$ _____

3. Analyze the following data from a randomized experiment that examined the effect of hours of sleep on happiness.

6 hours	8 hours	10 hours
10	10	9
15	11	9
11	10	9
9	11	8
8	13	9
12	13	7

Cut and paste your SPSS results, including the post-hoc test here:

Descriptives

VAR00001

	N	Mean	Std. Deviation	Std. Error	95% Confidence Interval for Mean		Minimum	Maximum
					Lower Bound	Upper Bound		
1.00	6	10.8333	2.48328	1.01379	8.2273	13.4394	8.00	15.00
2.00	6	11.3333	1.36626	.55777	9.8995	12.7671	10.00	13.00
3.00	6	8.5000	.83666	.34157	7.6220	9.3780	7.00	9.00
Total	18	10.2222	2.04524	.48207	9.2051	11.2393	7.00	15.00

ANOVA

VAR00001

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	27.444	2	13.722	4.714	.026
Within Groups	43.667	15	2.911		
Total	71.111	17			

