

COMPLICATIONS OF HEART DISEASE

By R.Thomas

Heart Failure #1

- ❖ Cardiovascular disease is the leading cause of death in the United States
- ❖ Heart disease remains a chronic and often progressive condition, associated with serious comorbidities, such as heart failure
- ❖ Heart failure (HF) is a clinical syndrome resulting from structural or functional cardiac disorders that impair the ability of a ventricle to fill or eject blood; the heart is unable to pump enough blood to meet the body's metabolic demands or needs



Copyright © 2022 Wolters Kluwer - All Rights Reserved

Heart Failure #2

- ❖ The term heart failure indicates myocardial disease, in which there is a problem with the contraction of the heart (systolic dysfunction) or filling of the heart (diastolic dysfunction) may cause pulmonary or systemic congestion
- ❖ Some cases are reversible depending on the cause
- ❖ Most HF is a chronic, progressive condition managed with lifestyle changes and medications



Copyright © 2022 Wolters Kluwer - All Rights Reserved

NEW YORK HEART ASSOCIATION CLASSIFICATION

- Class I

- Class II

- Class II

- Class IV

HEART FAILURE

- Left-sided heart failure
 - Systolic heart failure
 - EF ↓ 40%, pulmonary and systemic congestion
 - Diastolic heart failure
 - "stiffening"

- Right sided heart failure
 - ↓ right ventricle output
 - Systemic venous congestion

RISK FACTORS

- Left side heart failure
 - HTN
 - CAD, angina, MI
 - Valvular disease

- Right side heart failure
 - Left side failure
 - Right ventricle MI
 - Pulmonary disease (COPD, ARDS)

DIAGNOSTIC TEST

- **BNP**
- **Hemodynamic monitoring**
- **Echocardiogram- 2D or 3D**
 - **Ejection fraction**
 - Left ventricle 55- 70%
 - Right ventricle 45-60%
- **CXR**
- **ECG**
- **Cardiac Enzymes, CBC, Chemistry**

Clinical Manifestations of Heart Failure

Right Sided

- ❖ Viscera and peripheral congestion
- ❖ Jugular venous distention (JVD)
- ❖ Dependent edema
- ❖ Hepatomegaly
- ❖ Ascites
- ❖ Weight gain

Left Sided

- ❖ Pulmonary congestion, crackles
- ❖ S₃ or “ventricular gallop”
- ❖ Dyspnea on exertion (DOE)
- ❖ Low O₂ sat
- ❖ Dry, nonproductive cough initially
- ❖ Oliguria



Copyright © 2022 Wolters Kluwer - All Rights Reserved

SIGNS & SYMPTOMS

Left Sided HF


- Dyspnea, orthopnea, nocturnal dyspnea
- Fatigue
- Displaced apical
- S3
- Pulmonary congestion
- Frothy sputum
- AMS
- S/S of organ failure
- Hemodynamic findings:
 - CVP/ right atrial pressure
 - PAP
 - PAWP
 - Cardiac Output

Right Sided HF

- JVD
- Dependent edema
- Abdominal distention, ascites
- Fatigue, weakness
- Nausea & anorexia
- Polyuria
- Hepatomegaly & tenderness
- Weight gain
- Hemodynamic findings:
 - CVP/right arterial pressure

LEFT SIDED ♥ FAILURE

- PAROXYSMAL NOCTURNAL DYSPNEA
- RESTLESSNESS
- CONFUSION
- ELEVATED PULMONARY CAPILLARY WEDGE PRESSURE
- COUGH
- CRACKLES
- WHEEZES
- BLOOD TINGED SPUTUM
- ORTHOPNEA
- TACHYCARDIA
- EXERTIONAL DYSPNEA
- CYANOSIS



© 2000 Nursing Education Consultants

RIGHT SIDED ♥ FAILURE

(Cor Pulmonale)

- Fatigue
- May be secondary to chronic pulmonary problems
- ↑ Peripheral Venous Pressure
- Distended Jugular Veins
- Ascites
- Anorexia & Complaints of GI Distress
- Enlarged Liver & Spleen
- Weight Gain
- Dependent Edema



© 2007 Nursing Education Consultants, Inc.

ASSESSMENTS

- O2 sat
- Vital signs
- Heart rhythm
- Lung sounds
- Dyspnea
- Electrolytes
- Daily weight
- LOC
- I & O
- Sx of drug toxicity
- Support systems

Medical Management of the Patient with Heart Failure

- ❖ Vary according to the severity of the patient's condition, comorbidities, and cause
- ❖ Treatment may include
 - Oral and IV medications
 - Lifestyle modifications
 - Supplemental O₂
 - Surgical interventions: ICD and heart transplant
- ❖ Comprehensive education and counseling to patient and family is needed

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Medications Used to Treat HF #1

- ❖ Diuretics: decreases fluid volume, monitor serum electrolytes
- ❖ Angiotensin-converting enzyme (ACE) inhibitors: vasodilation; diuresis; decreases afterload; monitor for hypotension, hyperkalemia, and altered renal function; cough
- ❖ Angiotensin II receptor blockers: prescribed as an alternative to ACE inhibitors; work similarly
- ❖ Beta-blockers: prescribed in addition to ACE inhibitors; may be several weeks before effects seen; use with caution in patients with asthma

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Medications Used to Treat HF #2

- ❖ Ivabradine: decreases rate of conduction through the SA node; observe for decrease HR and BP
- ❖ Hydralazine and isosorbide dinitrate: alternative to ACE inhibitors; observe for decreased BP
- ❖ Digitalis: improves contractility; monitor for digitalis toxicity especially if patient is hypokalemic

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Medications Used to Treat HF #3

- ❖ IV medications: indicated for hospitalized patients admitted for acute decompensated HF
 - Dopamine: vasopressor to increase BP and myocardial contractility; adjunct with loop diuretics
 - Dobutamine: used for patients with left ventricular dysfunction; increases cardiac contractility and renal perfusion
 - Milrinone: decreases preload and afterload; causes hypotension and increased risk of dysrhythmias
 - Vasodilators: IV nitro, nitroprusside, nesiritide; enhance symptom relief

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Gerontologic Considerations

- ❖ May present with atypical signs and symptoms such as fatigue, weakness, and somnolence
- ❖ Decreased renal function can make older patients resistant to diuretics and more sensitive to changes in volume
- ❖ Administration of diuretics to older men requires nursing surveillance for bladder distention caused by urethral obstruction from an enlarged prostate gland

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

INTERVENTIONS

- O2
- High fowler's
- Bedrest till stable
- Dietary restrictions- NA an fluid
- Medications
 - Diuretics
 - To decrease preload
 - ↓ afterload
 - ACE inhibitors, beta blockers, ARB's
 - Inotropic Agents
 - ↑ contractility → ↑ cardiac output
 - Vasodilators
 - Anticoagulants

COMPLICATIONS OF HEART FAILURE

- Pulmonary Edema
- Cardiogenic Shock
- Pericardial Effusion/ Pericardial tamponade
- Systemic and pulmonary emboli (Thromboembolism)
- Organ failure
- Cardiac arrest

Assessment of the Patient with Heart Failure

- ❖ Focus
 - Effectiveness of therapy
 - Patient's self-management
 - S&S of increased HF
 - Emotional or psychosocial response
- ❖ Health history
- ❖ PE
 - Mental status; lung sounds: crackles and wheezes; heart sounds: S₃; fluid status or signs of fluid overload; daily weight and I&O; assess responses to medications

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Collaborative Problems and Potential Complications of the Patient with HF

- ❖ Pulmonary edema
- ❖ Hypotension, poor perfusion, and cardiogenic shock (see Chapter 11)
- ❖ Arrhythmias (see Chapter 22)
- ❖ Thromboembolism (see Chapter 26)
- ❖ Pericardial effusion

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Planning and Goals for the Patient With Heart Failure

❖ Goals

- Promote activity and reduce fatigue
- Relieve fluid overload symptoms
- Decrease anxiety or increase the patient's ability to manage anxiety
- Encourage the patient to verbalize his or her ability to make decisions and influence outcomes
- Educate the patient and family about management of the therapeutic regimen

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Nursing Interventions for the Patient with Heart Failure #1

❖ Promote activity tolerance

- Bed rest for acute exacerbations
- Encourage regular physical activity; build up to about 30 minutes daily
- Exercise training
- Pacing of activities; wait 2 hours after eating for physical activity
- Avoid activities in extreme hot, cold, or humid weather
- Modify activities to conserve energy
- Positioning; elevation of the head of bed to facilitate breathing and rest, support of arms

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Nursing Interventions for the Patient with Heart Failure #2

❖ Manage fluid volume

- Assess for symptoms of fluid overload
- Daily weight
- I&O
- Diuretic therapy; timing of meds
- Fluid intake; fluid restriction
- Maintenance of sodium restriction

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Patient Education for the Patient with Heart Failure

- ❖ Medications
- ❖ Diet: low-sodium diet and fluid restriction
- ❖ Monitor for signs of excess fluid, hypotension, and symptoms of disease exacerbation, including daily weight
- ❖ Exercise and activity program
- ❖ Stress management
- ❖ Prevention of infection
- ❖ Know how and when to contact health care provider
- ❖ Include family in education

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Pulmonary Edema

- ❖ Acute event reflecting a breakdown of physiologic compensatory mechanisms
- ❖ As LV begins to fail, blood backs up into the pulmonary circulation, causing pulmonary interstitial edema
- ❖ Results in hypoxemia, often severe
- ❖ Clinical manifestations: restlessness, anxiety, tachypnea, dyspnea, cool and clammy skin, cyanosis, weak and rapid pulse, cough, lung congestion (moist, noisy respirations), increased sputum production (sputum may be frothy and blood tinged), decreased level of consciousness

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

PULMONARY EDEMA

- Fluid in the lungs
- Etiology
- Risk Factors
- S/S
 - Persistent cough, pink frothy sputum, tachypnea, dyspnea, orthopnea, restless, anxiety, unable to sleep, hypoxemia, cyanosis (late sx), crackles, tachycardia, confusion, stupor, S3
- Dx- ABG, CXR, hemodynamic monitoring
- Assessment
- Treatment

Management of Pulmonary Edema

- ❖ Easier to prevent than to treat
- ❖ Early recognition: monitor lung sounds and for signs of decreased activity tolerance and increased fluid retention
- ❖ Minimize exertion and stress
- ❖ Oxygen; nonrebreather
- ❖ Medications
 - Diuretics (furosemide), vasodilators (nitroglycerin)

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Nursing Interventions for the Patient with Pulmonary Edema

- ❖ Positioning the patient to promote circulation
 - Positioned upright with legs dangling
- ❖ Providing psychological support
 - Reassure patient and provide anticipatory care
- ❖ Monitoring medications
 - I&O

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

End-of-Life Considerations

- ❖ HF is a chronic and often progressive condition
 - Need to consider issues related to the end of life
 - When palliative or hospice care should be considered

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Cardiogenic Shock

- ❖ A life-threatening condition with a high mortality rate
- ❖ Decreased CO leads to inadequate tissue perfusion and initiation of shock syndrome
- ❖ Commonly occurs following acute MI when a large area of myocardium becomes ischemic and hypokinetic
- ❖ Can occur as a result of end-stage HF, cardiac tamponade, pulmonary embolism (PE), cardiomyopathy, and arrhythmias

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

CARDIOGENIC SHOCK

- | | |
|--|---|
| <ul style="list-style-type: none"> ■ Pump failure/ heart failure ■ Etiology ■ S/S <ul style="list-style-type: none"> ▪ Initial ▪ Compensatory ▪ Progressive ▪ Refractory ■ Dx <ul style="list-style-type: none"> ▪ ECG, echo, CT, cardiac catheterization, CXR, cardiac enzymes | <ul style="list-style-type: none"> ■ Assessments <ul style="list-style-type: none"> ▪ VS, output, LOC, cardiac rhythm, skin color, temperature, moisture, capillary refill, turgor, hypoxia, hypotension, tachycardia, weak thready pulse ■ Treatment/ Management <ul style="list-style-type: none"> ▪ Decrease afterload ▪ Inotropics ▪ vasopressors |
|--|---|

Thromboembolism

- ❖ Patients with cardiovascular disorders are at risk for the development of arterial thromboemboli and venous thromboemboli (VTE)
- ❖ Decreased mobility and circulation increase the risk for thromboembolism in patients with cardiac disorders, including those with HF
- ❖ Intracardiac thrombi can form in patients with atrial fibrillation because the atria do not contract forcefully, and increasing the likelihood of thrombus formation
- ❖ Pulmonary embolism: blood clot from the legs moves to obstruct the pulmonary vessels

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Pericardial Effusion and Cardiac Tamponade

- ❖ Pericardial effusion is the accumulation of fluid in the pericardial sac
- ❖ Cardiac tamponade is the restriction of heart function because of this fluid, resulting in decreased venous return and decreased CO
- ❖ Clinical manifestations: ill-defined chest pain or fullness, pulsus paradoxus, engorged neck veins, labile or low BP, shortness of breath
- ❖ Cardinal signs of cardiac tamponade: falling systolic BP, narrowing pulse pressure, rising venous pressure, distant heart sounds

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

PERICARDIAL EFFUSION/ CARDIAC TAMPONADE

- Fluid in pericardial sac
- Etiology
- S/S
 - JVD, chest fullness, SOB, low bp, pulsus paradoxus
 - Cardiac tamponade- falling systolic bp, narrowing pulse pressure, JVD, distant or muffled heart sounds
- Dx
 - echo, CXR
- Treatment/Management
 - Pericardiocentesis, pericardiomy

Medical Management of Pericardial Effusion and Cardiac Tamponade

- ❖ Pericardiocentesis
 - Puncture of the pericardial sac to aspirate pericardial fluid
- ❖ Pericardiomy
 - Under general anesthesia, a portion of the pericardium is excised to permit the exudative pericardial fluid to drain into the lymphatic system

 Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

Sudden Cardiac Death or Cardiac Arrest

❖ Emergency management: cardiopulmonary resuscitation

- **A:** airway
- **B:** breathing
- **C:** circulation
- **D:** defibrillation for VT and VF

Wolters Kluwer

Copyright © 2022 Wolters Kluwer - All Rights Reserved

CARDIAC ARREST

- Ventricular fibrillation (V-fib)
- Asystole
- Pulseless Ventricular Tachycardia (Vtach),
- Pulseless Electrical Activity (PEA)
- S/S
- Treatment
 - IV fluid
 - Medications
 - Epinephrine
 - Vasopressin
 - NaHCO₃
 - Magnesium
 - Antiarrhythmics