

AS1 (Assignment 1, Unit 4) Logic of Hypothesis Testing

Please write "true" if the statement is true and "false" if the statement is false in the space provided:

- _____ 1. A statistical hypothesis test tests the null hypothesis rather than the researcher's hypothesis
- _____ 2. In most research situations, the goal is to reject the researcher's hypothesis
- _____ 3. A decision to reject the null hypothesis means that the data do provide evidence of a treatment effect: the independent variable did have an effect on the dependent variable
- _____ 4. If the data provide convincing evidence that the treatment does have an effect, then the correct statistical decision is to reject the null hypothesis
- _____ 5. For a hypothesis test, the critical region is defined as the set of test statistics that are very unlikely (very low probability) of being obtained if the null hypothesis is true
- _____ 6. If the obtained sample data (test statistic value) is inside the critical region, then we have provided support for the null hypothesis
- _____ 7. When the Z-test statistic, obtained from the sample data, falls inside the critical region, we reject the null hypothesis
- _____ 8. If the obtained sample data (test statistic value) are not in the critical region, the correct statistical decision is "fail to reject the null hypothesis."
- _____ 9. If you fail to reject the null hypothesis, it means that the data provide sufficient evidence to say that the treatment has an effect: the independent variable had an effect on the dependent variable
- _____ 10. Whenever the statistical decision is to reject the null hypothesis, there is a probability that the decision is incorrect and this probability is known as Type I error
- _____ 11. The probability of committing a Type I error is equal to alpha
- _____ 12. A Type II error occurs when a treatment actually does have an effect but the effect fails to show up in a research study
- _____ 13. One way to reduce the risk of Type I error, saying there is an effect when in fact there is not an effect, is to lower the alpha level from .05 to .01
- _____ 14. Type II error is to fail to reject a null hypothesis that is actually false
- _____ 15. In a research report, the term "statistically significant" is used to indicate that the null hypothesis was rejected
- _____ 16. In a research report, the notation $p < .05$ is used when the null hypothesis is rejected and the IV is shown to have a significant effect using an alpha level = .05

- _____ 17. Alpha (α) is the probability of committing a Type II error
- _____ 18. In a Type II error, the experimenter concludes that there is evidence for an effect when in fact an effect does not exist
- _____ 19. Changing the level of significance from .01 to .05 increases the risk of Type I error
- _____ 20. When a researcher report demonstrates a significant treatment effect at the .05 alpha level, you can be more confident that the effect is real than if the researcher had reported a significant effect with an alpha level of only .01
- _____ 21. There is always a possibility that the decision in a hypothesis test is incorrect
- _____ 22. To be more confident that a treatment actually does have a real effect, a researcher should use a large value for alpha
- _____ 23. The following hypothesis: "Nicotine should increase memory ability" requires a one-tail, directional test with the entire critical region located on the right side of the distribution
- _____ 24. The following hypothesis: "Nicotine should decrease memory ability" requires a one-tail, directional test with the entire critical region located on the left side of the distribution
- _____ 25. The following hypothesis: "Nicotine will have an effect on memory" requires a two-tail, non-directional test with the critical region divided between both tails of the distribution
- _____ 26. In the following hypothesis: "Nicotine will have an effect on memory," the dependent variable is nicotine
- _____ 27. In the following hypothesis: "Nicotine will have an effect on memory," the independent variable is memory
- _____ 28. Researchers collect data by measuring the dependent variable in their studies
- _____ 29. Independent variables are the variables that researchers manipulate in their studies
- _____ 30. In a directional (one-tail) hypothesis test, the entire critical region is located in either one of the other tail of the distribution, but not in both tails of the distribution
- _____ 31. The reason for computing Cohen's d is that a hypothesis test, per se, does not measure the size of the effect
- _____ 32. Cohen's d or any other " d " measures effect size
- _____ 33. Standard error is the discrepancy or difference, on average, that you should expect between your sample mean (M) and the population mean (μ)
- _____ 34. All hypotheses tests analyze (or compare) two main differences. These differences are the differences between or among means due to the IV, and difference between or among means due to chance.
35. What mean differences are you comparing when you compute a Z-test statistic?

36. Name the four steps of a hypothesis test:

1. _____

2. _____

3. _____

4. _____