

# AS1 (Assignment 1): Unit 1

## Introduction to Terminology, Scales of Measurement, Notation and Basic Computation

1. A researcher investigates the effects of amphetamine on memory. The researcher selects a random sample of 50 undergraduate students from a major New Jersey University. Half (25) of the students selected receive amphetamine while studying a word list and half (25) of the students receive a placebo. For this study
  - a. Identify the population: study all people, that how the amphetamine influence people's memory
  - b. Identify the sample: the 50 undergraduate students from a major New Jersey University, where 25 receive amphetamine and other half receive placebo.
2. Define the terms:
  - a. Population The entire set of a certain groups (categories) that researcher wants to study
  - b. Sample It is a small parts or individual of that population which researcher are interested in studying or analyzing, whereas, that selected sample are represent to analyze the population.
  - c. Parameter it is the summarized data and number describing from a population.
  - d. Statistic it is the summarized data that are selected from sample.
3. Name 3 descriptive statistics methods
  - 1) histogram
  - 2) calculating measures
  - 3) bar graph

Please place a T if the statement is true and an F if the statement is falsa in the space provided.

- T 4. A researcher calculates a mean from a population. Her mean is an example of a parameter.
- T 5. A researcher calculates a mean from a sample set of data. His mean is a statistic.
- F 6. The entire group of alcoholics in the state of NJ is an example of a population.
- F 7. A researcher who selects a sample from a population should expect no difference between the sample mean statistic and the true population parameter.
- T 8. The participants in a research study are classified as high, medium, or low in self-esteem. This classification involves measurement on an ordinal scale.
- F 9. A continuous variable must be measured on a nominal or an ordinal scale.
- F 10. Students in an introductory art class are classified as art majors and non-art majors. This is an example of measurement on an ordinal scale.
- F 11. Men's shirt sizes are classified as small, medium, large, and extra-large. This is an example of measurement on an interval scale.

\_\_T\_\_ 12. A researcher records the number of errors a rat makes running a maze. This is an example of measurement on a ratio scale.

\_\_F\_\_ 13. To determine how much difference there is between two individuals, you must use either a nominal or interval scale of measurement.

\_\_F\_\_ 14. If a researcher measures two individuals on a nominal scale, it is possible to determine which individual has the smaller score.

\_\_F\_\_ 15. Recording the number of students who are absent each day at a high school would be an example of measuring a continuous variable.

\_\_F\_\_ 16. A track coach records how much time each runner takes to complete the 100-yard dash. This is an example of measuring a discrete variable.

\_\_T\_\_ 17. Gender is a discrete variable while age is a continuous variable.

\_\_F\_\_ 18. A data set is described as consisting of  $n = 15$  scores. Based on the notation being used, the data set is a population.

\_\_F\_\_ 19. To compute  $(\Sigma X)^2$ , you have to remember to first square all of the scores.

\_\_T\_\_ 20. For the following scores, 1, 3, 5,  $\Sigma X^2 = 35$ .

\_\_F\_\_ 21. For the following scores, 4, 4, 2, 0,  $(\Sigma X)^2 = 36$ .

22. Statistical techniques are classified into two major categories: Name the categories and differentiate one major difference between them:

There are Descriptive statistics and Inferential statistics; descriptive statistic help to simplify and summarizing the data in which research studies, while inferential statistic can explain and analyze the date, such as its relationship in-between or its cause and effects.

23. You (1) select 30 people from a population, (2) measure the IQ of those 30 people, and (3) calculate a mean statistic = 110. What should you expect as far as the mean parameter? That is, would you expect the corresponding mean parameter to also be 110?

Since it is the selected sample from the population of the studies (mean of the IQ score), Which represented as M that it samples average is a statistic rather than parameter of population. That is, the some deviation of mean statistic and mean parameter will be expected.

24. Calculate each value requested for the following set of scores. Scores: 1, 7, 6, 4, 3, 0, 0, 1

$$N = \underline{8} \quad \Sigma X = \underline{22} \quad \Sigma X^2 = \underline{112} \quad (\Sigma X)^2 = \underline{484}$$

25. For the following set of scores, find the value of each expression: 10, 11, 10, 9, 7

$$\Sigma X = \underline{47}$$

$$\Sigma X^2 = \underline{451}$$

$$(EX)^2 = \underline{2209}$$

26. For the following set of scores, find the value of each expression

X

4

0

-3

a.  $\Sigma X = \underline{1}$

b.  $\Sigma X^2 = \underline{7}$

c.  $(EX)^2 = \underline{1}$

27. For the following set of scores, find the value of each expression:

X

6

-2

3

-1

$$n = \underline{4}$$

$$EX = \underline{6}$$

$$EX^2 = \underline{40}$$

$$(EX)^2 = \underline{36}$$

28. Two scores, X and Y, are recorded for each of  $n = 4$  subjects. For these scores, find the value of each expression.

<u>Subject</u>	<u>X</u>	<u>Y</u>
A	4	3
B	0	7
C	-1	5
D	3	2

$$\Sigma X = \underline{\underline{6}}$$

$$\Sigma Y = \underline{\underline{17}}$$

29. For each set of scores at the right, find the value of each expression.

x  
2  
4  
0  
-2

$$n = \underline{\underline{4}}$$

$$EX = \underline{\underline{4}}$$

$$EX^2 = \underline{\underline{16}}$$

$$(EX)^2 = \underline{\underline{16}}$$