

Diagnosis: DVT/Pain

Diastasis recti:  absent  present Hemorrhoids N/A  
 Incision:  None  
 Type:  transverse  vertical  umbilical  
 Closure:  staples  sutures  steri-strips  
 Condition:  approximated  redness  swelling  discharge  other ~~\_\_\_\_\_~~  
 Dressing: \_\_\_\_\_  
 fundal assessment:  firm  boggy  firm with massage  
<sup>2 fingers below umbilicus</sup>  midline  displaced:  right  left  
 Fingerwidths/ fingerbreaths: \_\_\_\_\_

**Genitalia**  
 Perineum:  intact  episiotomy  laceration/ repaired:  1°  2°  3°  4° Extension \_\_\_\_\_  
 Condition:  approximated  redness  swelling  discharge  other tender  
 Lochia: color scarlet amount serous odor N/A  
 Note: \_\_\_\_\_

**Extremities**  
 Varicosities N/A pedal pulses +2 homan's sign + Homan's sign  
 Edema:  none  non-pitting  pitting:  1+  2+  3+  4+  
 thrombophlebitis

**Elimination**  
 Voiding pattern:  normal  incontinence  distended  catheter type: \_\_\_\_\_  
 Last BM 11/26/13  constipation  diarrhea  hemorrhoid

**Psychological**  
 Stage:  taking in  taking hold  letting go  
 postpartum blues  history of depression  history of psychiatric diagnosis  
 Edinberg depression scale score \_\_\_\_\_  
**Bonding**  
 Describe interaction between mother and infant  
not much interaction w/ the baby.

Client Education Topic	Patient verbalize or demonstrate understanding or needs reinforcement	Additional information
educate w/ about breast feeding, the more they breastfeed, the more milk they produce		

Additional Comments:  
 \_\_\_\_\_  
 \_\_\_\_\_  
 \_\_\_\_\_