

Enoxaparin (Lovenox)

Enoxaparin (Lovenox)	
Classification:	Indication:
low molecular weight heparin	to treat or prevent a type of blood clot called deep vein thrombosis (DVT
Side effects/ adverse reactions:	Nursing Considerations:
<ul style="list-style-type: none">• Bleeding gums.• coughing up blood.• difficulty with breathing or swallowing.• increased menstrual flow or vagina bleeding.• nosebleeds.• paralysis.• prolonged bleeding from cuts.• red or black, tarry stools.	<ul style="list-style-type: none">• Contraindicated in pork hypersensitivity• Monitor for signs of bleeding• Administer in subcutaneous tissue• DO NOT eject air bubble prior to injection• DO NOT aspirate or massage site

Ferrous sulfate (Feosol)

Classification:	Indication:
a type of iron.	Prevention and treatment of iron-deficiency anemia
Side effects/ adverse reactions:	Nursing Considerations:
<ul style="list-style-type: none"> • feeling or being sick (nausea or vomiting), stomach pain or heartburn. • loss of appetite. • constipation. • diarrhea. • dark or black poo. • black stained teeth 	<p>May cause seizures, hypotension, constipation, epigastric pain, diarrhea, skin staining, anaphylaxis</p> <ul style="list-style-type: none"> • Assess nutritional status, bowel function • Monitor hemoglobin, hematocrit, iron levels • May cause elevated liver enzymes • Take on an empty stomach to increase absorption/vitamin c helps with absorption • Use z-track for IM injections