

Nyack College SON- Concept Map

Student- **Minho Kim**
 Submission date- **02/16/23**

Nursing Dx Risk for bleeding related to boggy fundus
Expected Outcome Patient will be free of hemorrhage until the discharge. (02/11/23)

1. Monitor vital signs every 4 hours
 2. Educate patient about signs and symptoms of bleeding
 3. Educate patient to massage fundus twice a day
 4. Educate patient to avoid constipation
- Evaluation-** until the day of discharge, patient remained free of hemorrhage. (02/22/23)

Nursing Dx Risk for impaired parenting related to dependence on substance and history of depression
Expected Outcome Patient will demonstrate understanding of negative effect of second-hand smoking of marijuana to children by the end of the shift (2/10/23)

Interventions

1. Assess for signs of postpartum depression
2. Educate patient about the negative effect of second hand smoking of marijuana
3. Closely monitor interactions with newborn
4. Provide information about social help system

Evaluation- Patient verbalized understanding of negative

Past Medical History

Obesity, depression,

Assessment (Mother): AAO x4, Vitals: within normal range, pain 3/10, Vaginal delivery, linea negra, striae present, soft breast, fundus firm with massage, midline 3 fingers down, rubra scent lochia, earthy odor, 2+ pedal pulse bilaterally, taking-in phase
Assesment(baby): Vitals within normal range, breastfed, weigh, length, head size AGA 10th to 90th percentile, New ballard score of 40, all reflexes positive, symmetrical, circumcised, parents refused administration of any medications

Medications

Mother
 - Acetaminophen 650mg every 6 hours PRN
 - ferrous sulfate 325mg 2times a day
 - benzocaine 20% topical 2time a day
 - dextrose 10% in water 750ml over 20min
 - docusate sodium 100mg PRN
 - glucagon 1mg PRN every 15 min
 - glycerin witch haze

Baby
 - No medication given

Nursing Dx Risk for bleeding related to circumcision and insufficient vitamin K.
Expected Outcome Newborn patient will maintain free of bleeding complications until discharge (02/11/23)

Interventions

1. Assess and monitor vital signs every 1 hour
2. Educate parents about risk for bleeding
3. Obtain informed consent for circumcision and risk that follows when vitamin K is not given
4. Assess

Evaluation- Newborn patient maintained clear airway until the discharge

Nursing Dx

Expected Outcome

4.