

High Risk Labor & Birth

Dr. Rexi Thomas



Dysfunctional Labor

- ▶ Problem with the “Powers”
 - Hypertonic Uterine Dysfunction
 - Hypotonic Uterine Dysfunction
- ▶ Risk factors
- ▶ Ultrasound to R/O CPD



Hypertonic Uterine Dysfunction

- Nonproductive, uncoordinated ctx
- Frequent and prolong duration
- Problem with uterine relaxation
- Latent phase
- Uteroplacental insufficiency
- Maternal exhaustion
- ▶ Treatment



Hypotonic Uterine Dysfunction (FTP)

- › Active phase
- › Ineffective or absent ctx
- › Associated factors
- › Assessment
- › Treatment
 - Position changes
 - Ambulate
 - Induction/augmentation of labor
 - AROM
 - Pitocin

Malpositions/ Malpresentation

- › Malposition– other than OA
- › Malpresentation– other than vertex
- › Occipit posterior most common
- › C/S may be necessary
- › Nursing Interventions
 - Change positions
 - Counterpressure
 - May need operative vaginal delivery

Breech Presentations

- › Complete breech
- › Frank breech
- › Footling breech

- › External Version
- › C/S delivery
- › May pass meconium

Risk Factors for Dystocia #1

- ❖ Epidural analgesia/excessive analgesia
- ❖ Multiple gestation
- ❖ Hydramnios
- ❖ Maternal exhaustion
- ❖ Ineffective maternal pushing technique
- ❖ Occiput posterior position

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Risk Factors for Dystocia #2

- ❖ Longer first stage of labor
- ❖ Nulliparity, short maternal stature
- ❖ Fetal birth weight over 8.8 lb
- ❖ Shoulder dystocia
- ❖ Abnormal fetal presentation or position
- ❖ Fetal anomalies

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Causes of Dystocia #1

- ❖ Problems with powers
 - Hypertonic uterine dysfunction
 - Hypotonic uterine dysfunction
 - Protracted disorders
 - Arrest disorders
 - Precipitate labor
- ❖ Problems with the passageway
 - Pelvic contraction
 - Obstructions in maternal birth canal

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Causes of Dystocia #2

- ❖ Problems with passenger
 - Occiput posterior position
 - Breech presentation
 - Multifetal pregnancy
 - Macrosomia and CPD
 - Structural abnormalities (see Table 21.1)
- ❖ Problems with psyche: psychological distress

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved


Shoulder Dystocia

- Fetal Risks
 - Asphyxia, nerve damage, clavicle fx, CNS injury, death
- Maternal Risks
 - Postpartum hemorrhage, extensive lacerations, uterine rupture, infection, fistulas, bladder injury
- McRoberts maneuver
- Suprapubic pressure



Copyright © 2010, 2008, 2002, 1996, 1994 by Saunders, an imprint of Elsevier Inc.

Shoulder Dystocia



Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Precipitate Labor

- › Labor < 3 hours
- › Maternal Risks
- › Fetal Risks
- › Management

- › Precipitous delivery



Preterm Labor

- › Incidence
- › Premature rupture of membranes (PROM)
- › Preterm premature rupture of membranes (PPROM)



Risks Factors for Preterm labor

- › Dehydration
- › Multiple gestation
- › Fetal abnormalities
- › Hx of preterm birth
- › Underweight
- › BV, UTI
- › DM,
- › ↑ blood pressure
- › Peridontal disease
- › Poor weight gain
- › Drug use
- › Smoking
- › Hx of cervical surgeries
- › DES exposure

Predicting Preterm Births

- › Cervical length
- › Previous hx etiology of PROM
- › + fetal fibronectin (FFN) screening

Management of Preterm Labor

- › Identify cause
- › Laboratory and diagnostic testing: CBC, UA, amniotic fluid analysis, fetal fibronectin, cervical length via transvaginal ultrasound, salivary estriol, home uterine activity monitoring
- › Corticosteroid therapy for fetal lung maturity
 - Celestone
 - Betamethasone
 - Surfactant
- › Limit activity/ Bedrest
- › ↑ hydration
- › Tocolytics
 - Terbutaline (Brethine)
 - Indomethacin
 - Magnesium Sulfate
 - < 25 weeks
 - < 34 weeks

Postterm Labor

- ❖ Pregnancy continuing past end of 42 weeks' gestation
- ❖ Unknown etiology
- ❖ Maternal risks: cesarean birth, dystocia, birth trauma, postpartum hemorrhage, and infection
- ❖ Fetal risks: macrosomia, shoulder dystocia, brachial plexus injuries, low Apgar scores, postmaturity syndrome, and cephalopelvic disproportion

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Postterm Pregnancy: Assessment and Management

- ❖ Nursing assessment: estimated date of birth; daily fetal movement counts, nonstress tests twice weekly, amniotic fluid analysis, weekly cervical examinations, client understanding, anxiety, and coping ability
- ❖ Nursing management: fetal surveillance; decision for labor induction; support; education, intrapartal care

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Labor Induction and Augmentation

- ❖ Induction: stimulating contractions via medical or surgical means
- ❖ Augmentation: enhancing ineffective contractions after labor has begun
- ❖ Indications: prolonged gestation, prolonged premature rupture of the membranes, gestational hypertension, cardiac disease, renal disease, chorioamnionitis, dystocia, intrauterine fetal demise, isoimmunization, and diabetes

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Labor Induction: Therapeutic Management

- ❖ Cervical ripening (Bishop score, see Table 21.2)
- ❖ Herbal agents
- ❖ Castor oil, hot baths, enemas
- ❖ Sexual intercourse with breast stimulation
- ❖ Mechanical methods and surgical methods
- ❖ Pharmacologic agents (see Drug Guide 21.2)
- ❖ Oxytocin

Wolters Kluwer

Copyright © 2021 Wolters Kluwer All Rights Reserved

Induction/ Augmentation of Labor

- ▶ Bishop's Scale
 - Dilation
 - Effacement
 - Consistency
 - Station
 - Position
- ▶ Prostaglandin therapy
- ▶ Amniotomy/ AROM
- ▶ Oxytocin Infusion



Intrauterine Fetal Demise

- ❖ Numerous causes
- ❖ Devastating effects on family and staff
- ❖ Nursing assessment
 - Inability to obtain fetal heart sounds
 - Ultrasound to confirm absence of fetal activity
 - Labor induction
- ❖ Nursing management
 - Assistance with grieving process
 - Referrals

Wolters Kluwer

Copyright © 2021 Wolters Kluwer All Rights Reserved

Uterine Rupture

- ❖ Obstetric emergency; onset marked by sudden fetal bradycardia
- ❖ Nursing assessment
 - Risk factors
 - Onset of sudden fetal distress; other signs
- ❖ Nursing management
 - Preparation for urgent cesarean birth
 - Continuous maternal and fetal monitoring

Wolters Kluwer

Copyright © 2021 Wolters Kluwer All Rights Reserved

Amniotic Fluid Embolism

- ❖ Obstetric emergency
- ❖ Sudden onset of hypotension, hypoxia, and coagulopathy due to breakage in barrier between maternal circulation and amniotic fluid
- ❖ Nursing assessment: difficulty breathing, hypotension, cyanosis, seizures, tachycardia, coagulation failure, DIC, pulmonary edema, uterine atony with subsequent hemorrhage, ARDS, cardiac arrest
- ❖ Nursing management: supportive measures to maintain oxygenation and hemodynamic function and to correct coagulopathy; critical care monitoring

Wolters Kluwer

Copyright © 2021 Wolters Kluwer All Rights Reserved

Amniotic Fluid Embolism

- Treatment
 - CPR
 - Intubation
 - Hemodynamics
 - Central venous line
 - Immediate delivery



Amnioinfusion

- ❖ Indications
 - Severe variable decelerations due to cord compression
 - Oligohydramnios due to placental insufficiency
 - Postmaturity or rupture of membranes
 - Preterm labor with premature rupture of membranes
 - Thick meconium fluid
- ❖ Nursing management: teaching, maternal and fetal assessment, preparation for possible cesarean birth

Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved

Forceps- or Vacuum-Assisted Birth #1

- ❖ Application of traction to fetal head
- ❖ Indications: prolonged second stage of labor, nonreassuring FHR pattern, failure of presenting part to fully rotate and descend, limited sensation or inability to push effectively, presumed fetal jeopardy or fetal distress, maternal heart disease, acute pulmonary edema, intrapartum infection, maternal fatigue, infection
- ❖ Risk of tissue trauma to mother and newborn
- ❖ Prevention as key

Wolters Kluwer


Copyright © 2021 Wolters Kluwer. All Rights Reserved

Forceps- or Vacuum-Assisted Birth #2



Wolters Kluwer

Copyright © 2021 Wolters Kluwer. All Rights Reserved



Cesarean Birth

- ❖ Classic or low transverse incision (see Figure 21.8)
- ❖ Major surgical procedure with accompanying risks
- ❖ Nursing assessment: history and physical examination for maternal and fetal indications
- ❖ Nursing management
 - Preoperative care
 - Postoperative care

Wolters Kluwer


Copyright © 2021 Wolters Kluwer. All Rights Reserved

Cesarean Birth: Technique


- Preparation
 - Anesthesia
 - Medication
 - Laboratory studies
 - Prophylactic antibiotics
 - Skin prep
 - Foley catheter

Copyright © 2011, 2006, 2002, 1998, 1994, by Saunders, an imprint of Elsevier Inc.

Skin Incisions



Uterine Incisions




Vaginal Birth after Cesarean (VBAC)

- ❖ Controversy related to risk of uterine rupture and hemorrhage
- ❖ Contraindications
- ❖ Special areas of focus: consent, documentation, surveillance, and readiness for emergency
- ❖ Nurses as advocates for clients; expertise in reading fetal monitoring tracings to identify nonreassuring pattern and instituting measures for emergency delivery

Version: Indications

- ▶ External version
 - Change the fetal position from a breech, shoulder (transverse lie), or oblique presentation to cephalic
- ▶ Internal version
 - Change the position of a second twin in a vaginal birth

External Version



Internal Version

