

Valentine Sanon
NUR 203

41. ibuprofen (Advil, Caldolor, Motrin)	
Classification:	Indication:
NSAID/Analgesic	Treatment of mild to moderate pain ,fever ,migraine headache ,primary dysmenorrhea. Treatment of osteoarthritis, rheumatoid arthritis ,treatment of juvenile arthritis ,
Side effects/ adverse reactions:	Nursing Considerations:
Agranulocytosis Bronchospasm, dizziness, G.I. bleeding , edema, eye changes, GI pain, Heart failure, insomnia, nausea, somnia, stomatitis rash.	Ensure hydration with IV use Stop if eye changes renal impairment Hepatic impairment Give me food if G.I. upset Patient should avoid OTC products that may contain ibuprofen Report ankle swelling Report black Tarry stool report vision changes

42. acetaminophen (Tylenol)

42. acetaminophen (Tylenol)	
Classification: analgesics and antipyretics	Indication: Temporary reduction of fever Temporary relief of minor aches and pains close by common cold and influenza; headache, sore throat, toothache, backache, menstrual cramps, minor arthritis pain, muscle aches.
Side effects/ adverse reactions: Headache, hemolytic anemia, renal dysfunction, skin rash and fever , Hepatotoxicity usually associated with chronic use and overdose	Nursing Considerations: Do not exceed recommended dosage Avoid combining many different products contain acetaminophen Administer IV over 15 minutes Do not recommend for longer than 10 days Have Acetylcysteine for overdose The nurse should closely monitor AST and ALT levels and BUN and creatinine

43. baclofen (Gablofen, Lioresal)

Classification:	Indication:	
Centrally acting skeletal muscle relaxants	Alleviation of s&sx of spasticity from MS or spinal cord injuries Spinal cord injuries. Other spinal cord disease	
Side effects/ adverse reactions:	Nursing Considerations:	
confusion dizziness weakness headache nausea Constipation Urinary frequency insomnia fatigue	Use caution if spasticity needed to stay upright Taper slowly if Discontinuing not for use in pregnancy Patient should avoid OTC sleeping drugs and alcohol monitor the patient's bowel actions to ensure regularity	

44. methadone (Dolophine, methadose)	
Classification:	Indication:
Opioid agonist analgesic	Relief of severe pain unresponsive to nonopioid analgesics Detoxification temporary maintain treatment of opioid addiction
Side effects/ adverse reactions:	Nursing Considerations:
Apnea Cardiac arrest Circulatory depression Itchy skin Heavy sweating Constipation Weight gain.	Have opioid antagonist, equipment for assisted or controlled respiration on hand during parenteral administration Not for use in pregnancy We poured difficulty breathing severe nausea and vomiting

45. morphine (AVINza, Kadian)	
Classification:	Indication:
Opioid agonist and Analgesic	Relief of motivate to severe pain and analgesic adjunct during anesthesia , preop medication Relief of intractable pain
Side effects/ adverse reactions:	Nursing Considerations:
Apnea Laryngospasm drowsiness stomach pain and cramps dry mouth sweating mood changes headache	Patient should lie during IV use Monitor patient's respiratory rate prior to administration. Reassess pain after administration of morphine. Monitor for hypotension frequently up to 24 hours after

nervousness dizziness	
--------------------------	--

46. naloxone (Narcan)	
Classification:	Indication:
opioid antagonist	emergency tx for opioid overdose or suspected opioid overdose.
Side effects/ adverse reactions:	Nursing Considerations:
tachycardia Sweating runny nose weakness nervousness restlessness	Monitor continually After use Maintain open airway provide life support as needed patient should report sweating tremors

irritability diarrhea nausea vomiting stomach cramps hypertension Pulmonary Edema	
---	--

47. dopamine (Intropin)	
Classification:	Indication:
Dopaminergic, Sympathomimetic	Correction of hemodynamic imbalance, low cardiac output hypotension.
Side effects/ adverse reactions:	Nursing Considerations:

<p>Angina dyspnea ectopic beats nausea vomiting hypotension palpitations tachycardia</p>	<p>Use extreme caution in calculating doses Base dosing on patient response Give in large vein; avoid extravasation monitor urine output monitor cardiac output blood pressure during infusion</p>
--	--

48. hydrocortisone (Cortef, Pandel, Hydrocort)	
<p>Classification: corticosteroids</p>	<p>Indication: Replacement treatment in adrenocortical insufficiency Allergic states inflammatory disorders, Hematology disorder, ulcerative colitis.</p>
<p>Side effects/ adverse reactions: Anaphylactoid reaction Amenorrhea Heart failure Headache fluid retention immunosuppression infection irregular menses local pain irritation at application site muscle weakness pancreatitis Vertigo</p>	<p>Nursing Considerations: Rotate IM injection site Taper doses when discontinuing high-dose or long-term treatment Patient should report infection swelling difficulty breathing Monitor signs of hypersensitivity reactions or anaphylaxis, pulmonary symptoms</p>

49. fentanyl (Abstral, Actiq, Fentora)

Classification:		Indication:	
Opioid agonist analgesic		Control of postop pain, Tachypnea emergency delirium, Management of chronic pain in patient requiring continuous opioid analgesia over extended period. Treatment of breakthrough pain in cancer patients treated with and tolerant opioids.	
Side effects/ adverse reactions:		Nursing Considerations:	
Apnea Cardiac arrest Clamminess Confusion Constipation Dizziness Headache Lethargy Sweating vertigo		Adjust dose as needed tolerated for pain Patient should avoid grapefruit juice Remove or change the old patch before applying new one Report difficulty breathing	

50. oxycodone (OxyContin)

50. oxycodone (OxyContin)	
Classification:	Indication:
Opioid agonist analgesic	Relief of moderate to moderately severe pain Treatment of breakthrough pain Management of moderate to severe pain when continuous around the clock analgesic needed for extended period
Side effects/ adverse reactions:	Nursing Considerations:
Bronchospasm Cardiac arrest Constipation dizziness drowsiness flushing laryngospasm lightheadedness nausea vomiting respiratory arrest sedation Shock sweating vision changes	Have opioid antagonist Patient should swallow CR form whole and not cut, crush or chew it Use laxative for constipation report difficulty breathing Report fainting