References

Hishimoto, A., Fukutake, M., Mouri, K., Nagasaki, Y., Asano, M.,Ueno,Y.,Nishiguchi, N., & Shirakawa, O. (2010). Alcohol and aldehyde dehydrogenase polymorphisms and risk for suicide: a preliminary observation in the Japanese male population. *Genes, Brain and Behavior*, 498-502

Kawaguchi, H., Koike, S. (2016). Association between the Density of Physicians and Suicide Rates in Japan: Nationwide Ecological Study Using a Spatial Bayesian Model. *PLoS ONE, Vol. 11 Issue 2,* 1-9.

Kitanaka, J. (2008). Diagnosing Suicides of Resolve: Psychiatric Practice in Contemporary Japan. *Culture, Medicine & Psychiatry. Jun2008, Vol. 32 Issue 2*, 152-176.

Ozawa-de Silva, C. (2008). Too Lonely to Die Alone: Internet Suicide Pacts and Existential Suffering in Japan. *Culture, Medicine & Psychiatry, Vol. 32 Issue 4*, 516-551.

Picone, M. (2012). Suicide and the afterlife: Popular religion and the standardisation of 'culture' in Japan. *Cult Med Psychiatry*, 36:391-408.

Russel, R., Metraux, D. & Tohen, M. (2016). Cultural influences on suicide in Japan. *Psychiatry and Clinical Neurosciences. Jan 2017, Vol. 71 Issue 1,* 2-5.