

Nyack College SON- Concept Map

Student- Lakeisha P. Beasley
 Submission date- 3/20/23

Nursing Dx Risk for impaired skin integrity related to fecal and urinary incontinence
Expected Outcome
 Patient will not develop a pressure injury within the next 8 hours

Interventions

1. Perform skin assessment each shift.
2. Use pillows or foam mattress to prevent direct contact with mattress to alternate pressure
3. maintain clean, dry and moisturized skin
4. Educate the client on how to call for assistance when incontinent care is required.

Evaluation- Goal met. Patient did not develop a pressure injury during 8 hours shift

Nursing Dx Ineffective airway clearance related to endotracheal intubation as evidenced by excessive secretions
Expected Outcome
 The patient will maintain open airways evidenced by normal breath sounds after suctioning

Interventions

1. Assess airway patency
2. Educate the parent or caregiver on the suctioning procedure and the desired outcome
3. monitor ET tube placement
4. Institute airway suctioning as indicated for adventitious lung sounds or audible respirations

Evaluation-
 Goal not met. Patient had minimal secretions so no suction was indicated at this time.

Past Medical History
 Dental Caries
 Airway compromised
 cerebral Palsy
 Contractures involving both knees
 Developmental delay
 Dysphagia
 Fracture history
 Hypoxic ischemic encephalopathy
 wrist
 optic atrophy
 seizure
 spasticity
 Tracheostomy in place

Arteriovenous malformation

Medications (see back)
 Acetaminophen (Tylenol) 650mg PO q4h PRN
 Baclofen (Lioresal) 15mg PO TID @ 0830, 1500, 2030 SCH
 Benzoyl Peroxide Gel 0gm TP Bedtime SCH
 Chlorhexidine Gluconate (Peridex) 0ml 4mm BID SCH
 Enoxaparin Sodium (Lovenox) 30mg Subq Daily @ 0500 SCH
 Erythromycin (Ery 2% Pads) 1 each TP Daily SCH
 Famotidine (Pepicol) 20mg PO BID SCH
 Fentanyl Sulfate (Doseol 300mg/5mg Udc*) 300 mg q 4 BID
 Fluticasone Propionate (Flovent HFA Inhaler) 1H BID
 Dose: 2 puffs both times daily
 Levamisole HCl (Expensol) 0.63mg 1H q3H PRN
 Levethracetam (Keppra) 2000mg PO BID SCH
 melatonin long PO bedtime PRN
 midazolam HCl (versed) 10mg both times q 15 PRN

Nursing Dx Constipation related to use of medication
Expected Outcome
 Within 24 hours of nursing treatment the patient will have 1 bowel movement.

Interventions

1. Encourage fluid intake
2. Administer stool softeners per provider's order
3. ~~educate patient~~ Educate patient on desired effects of stool softeners and laxatives.
4. monitor intake and output each shift.

Evaluation- Goal met. Patient had 1 large bowel movement within 24 hours.

Nursing Dx Risk for aspiration related to impaired swallow
Expected Outcome
 The patient will not aspirate during breakfast and lunch.

Interventions

1. Eliminate any environmental stimuli during mealtimes to reduce risk for aspiration.
2. Ensure patient is sitting in an upright position with head, liquid or medication is being administered.
3. Verify all foods on meal tray are pureed per provider's order.
4. Allow adequate time for patient to eat and drink to ensure proper chewing and swallowing

Evaluation-
 Goal met. Patient did not aspirate during breakfast or lunch.