

References

- Hishimoto, A., Fukutake, M., Mouri, K., Nagasaki, Y., Asano, M., Ueno, Y., Nishiguchi, N., & Shirakawa, O. (2010). Alcohol and aldehyde dehydrogenase polymorphisms and risk for suicide: a preliminary observation in the Japanese male population. *Genes, Brain and Behavior*, 498-502
- Kawaguchi, H., Koike, S. (2016). Association between the density of physicians and suicide rates in Japan: Nationwide Ecological Study Using a Spatial Bayesian Model. *PLoS ONE*, Vol. 11 Issue 2, 1-9.
- Kitanaka, J. (2008). Diagnosing suicides of resolve: Psychiatric practice in contemporary Japan. *Culture, Medicine & Psychiatry*. Jun2008, Vol. 32 Issue 2, 152-176.
- Ozawa-de Silva, C. (2008). Too lonely to die alone: Internet suicide pacts and existential suffering in Japan. *Culture, Medicine & Psychiatry*, Vol. 32 Issue 4, 516-551.
- Picone, M. (2012). Suicide and the afterlife: Popular religion and the standardisation of 'culture' in Japan. *Cult Med Psychiatry*, 36:391-408.
- Russel, R., Metraux, D. & Tohen, M. (2016). Cultural influences on suicide in Japan. *Psychiatry and Clinical Neurosciences*. Jan 2017, Vol. 71 Issue 1, 2-5.