

DRUG CARD TEMPLATE:

GI Generic name (brand name) Hydrochlorozide/ microsside	
Classification:	Indication:
Diuretic	Hydrochloktoobiapide is used to treat high blood pressure
Side effects/ adverse reactions:	Nursing Considerations:
Back , leg or stomach Black tarry stools Blue lips in the urine Chest pain or tightness	Give with food or milk GI upset occurs Reduce dosage of other anti hypertension b6 at least 50%

DRUG CARD TEMPLATE:

62		Generic name (brand name) Spironolactone / aldosterone	
Classification: Aldosterone Receptor antagonist		Indication: Heart failure; hypertension nephrotic syndrome and primary hyperaldosteronism	
Side effects/ adverse reactions: Feeling dizzy Feeling on being sick Muscle leg or cramps Feeling tired low or in energy		Nursing Considerations: Mark calendars of edema outpatients as reminders of alternate day or 3- to 5-day/wk therapy. Give daily doses early so that increased urination does not interfere with sleep. Make suspension as follows: Tablets may be pulverized and given in cherry syrup for young children.	

DRUG CARD TEMPLATE:

63	
Generic name (brand name) Mannitol / osmitrol	
Classification:	Indication:
Diuretics, Osmotic Agents.	Mannitol has approval for the reduction of intracranial pressure and brain mass.[1] Mannitol is approved to reduce intraocular pressure if this is not achievable treat the oliguric phase before irreversible damage.[3]
Side effects/ adverse reactions:	Nursing Considerations:
Common side effects may include: increased urination; nausea, vomiting; fever, chills, headache, runny nose; chest pain; rash; or dizziness, blurred vision.	Nursing care of the patient receiving mannitol requires vigilant monitoring of electrolytes and overall fluid balance, and observation for the development of cardiopulmonary complications in addition to neurologic assessment.

DRUG CARD TEMPLATE:

64		Generic name (brand name) Oxybutynin chloride Ditropan XL
Classification:	Indication:	
anticholinergics/ antimuscarinics.	Oxybutynin is an anticholinergic medication that is FDA-approved and indicated in patients with overactive bladder or symptoms of detrusor overactivity, including urinary frequency and urgency.	
Side effects/ adverse reactions:		Nursing Considerations:
Common side effects of oxybutynin dry mouth. headache. feeling dizzy, sleepy, or a spinning sensation (vertigo) diarrhoea or being sick (vomiting) constipation. farting and burping (wind) stomach pain. dry eyes.		Oxybutynin may cause anxiety, confusion, irritability, sleepiness or unusual drowsiness, or hallucinations (seeing, hearing, or feeling things that are not there). These symptoms are more likely to occur when you begin taking this medicine, or when the dose is increased.

DRUG CARD TEMPLATE:

65	
Generic name (brand name) Phenazopyridine HCl (Probid, Pyridiate, Baridium, Uricalm, Urodine, and UTI Relief)	
Classification:	Indication:
urinary tract analgesics.	Phenazopyridine HCl is indicated for the symptomatic relief of pain, burning, urgency frequency,
Side effects/ adverse reactions:	Nursing Considerations:
<p>Image result for phenazopyridine side effect swelling, rapid weight gain; confusion, loss of appetite, pain in your side or lower back; fever, pale or yellowed skin, stomach pain, nausea and vomiting; or. blue or purple appearance of your skin.</p> <p>... Common side effects of phenazopyridine may include: headache; dizziness; or. upset stomach.</p>	<p>It should be avoided while breastfeeding, especially with an infant under 1 month of age or with glucose-6-phosphate dehydrogenase (G6PD)</p>

DRUG CARD TEMPLATE:

66	
Generic name (brand name) Finasteride / proscar and prapecia	
Classification:	Indication:
5-alpha reductase inhibitors.	Finasteride is indicated for the treatment of symptomatic benign prostatic hyperplasia (BPH) in men with an enlarged prostate to improve symptoms, reduce the risk of acute urinary retention,
Side effects/ adverse reactions:	Nursing Considerations:
<p>Finasteride can give you side effects such as not being able to get an erection (impotence) and having less interest in sex. These side effects usually pass after a while. If these side effects do not go away, are worrying you or affecting your sex life, speak to your doctor.</p>	<p>Finasteride may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away: inability to have or maintain an erection. decreased sexual desire. problems with ejaculation (including decreased volume of ejaculate) pain in the testicles. depression.</p>

DRUG CARD TEMPLATE:

67		Generic name (brand name) Fluoxetine / Prozac.
Classification:	Indication:	
selective serotonin reuptake inhibitors (SSRIs)	Fluoxetine is indicated for both acute and maintenance treatment of major depressive disorder, obsessive compulsive disorder, and bulimia nervosa; however, it is only indicated for acute treatment of panic disorder independent of whether agoraphobia is present.	
Side effects/ adverse reactions:		Nursing Considerations:
What side effects can this medication cause? nervousness. anxiety. difficulty falling asleep or staying asleep. nausea. diarrhea. dry mouth. heartburn. yawning.		Keep this drug, and all medications, out of the reach of children. You may experience these side effects: Dizziness, drowsiness, nervousness, insomnia (avoid driving or performing hazardous tasks); nausea, vomiting, weight loss (eat small frequent meals; monitor your weight loss); sexual dysfunction; flulike symptoms.

DRUG CARD TEMPLATE:

68	
Generic name (brand name) Amitriptyline Elavil	
Classification:	Indication:
tricyclic antidepressants.	indications are anxiety, post-traumatic stress disorder, insomnia, chronic pain (diabetic neuropathy, fibromyalgia), irritable bowel syndrome, interstitial cystitis (bladder pain syndrome), migraine prophylaxis, postherpetic neuralgia, and sialorrhea
Side effects/ adverse reactions:	Nursing Considerations:
What side effects can this medication cause? nausea. vomiting. drowsiness. weakness or tiredness. nightmares. headaches. dry mouth. constipation.	Amitriptyline can increase the risk of suicidal thoughts and behavior, especially in children, adolescents, and young adults. People of all ages who are started on antidepressant therapy should be watched closely for signs of changes in behavior or worsening depression.

DRUG CARD TEMPLATE:

69	
Generic name (brand name) Lithium Priadel, Camcolit, Liskonium, Li-Liquid.	
Classification:	Indication:
antimanic agents	Lithium is indicated in the treatment of manic episodes of Bipolar Disorder.
Side effects/ adverse reactions:	Nursing Considerations:
Common side effects feeling sick (nausea) diarrhoea. a dry mouth and/or a metallic taste in the mouth. feeling thirsty and needing to drink more and pee more than usual. slight shaking of the hands (mild tremor) feeling tired or sleepy. weight gain (this is likely to be very gradual)	Take this drug exactly as prescribed, after meals or with food or milk. ... Eat a normal diet with normal salt intake; maintain adequate fluid intake (at least 2.5 quarts/day). Arrange for frequent checkups, including blood tests. ... Use contraception to avoid pregnancy.

DRUG CARD TEMPLATE:

70		Generic name (brand name) Haloperidol Haldol, Haldol decanoate and Serenace.
Classification:	Indication:	
conventional antipsychotics.	Haloperidol is indicated for the treatment of the manifestations of several psychotic disorders including schizophrenia	
Side effects/ adverse reactions:		Nursing Considerations:
Haloperidol may cause side effects. Tell your doctor if any of these symptoms are severe or do not go away: dry mouth. increased saliva. blurred vision. loss of appetite. constipation. diarrhea. heartburn. nausea.		This medicine may cause some people to become dizzy, drowsy, or may cause trouble with thinking or controlling body movements, which may lead to falls, fractures or other injuries. Even if you take haloperidol at bedtime, you may feel drowsy or less alert on arising.